

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

Draft Genome Sequences of Three Capnocytophaga canimorsus Strains Isolated from Healthy Canine Oral Cavities

Manfredi, Pablo; Renzi, Francesco; Cornelis, Guy

Published in:
Genome Announcements

DOI:
[10.1128/genomeA.00199-15](https://doi.org/10.1128/genomeA.00199-15)

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (HARVARD):
Manfredi, P, Renzi, F & Cornelis, G 2015, 'Draft Genome Sequences of Three Capnocytophaga canimorsus Strains Isolated from Healthy Canine Oral Cavities', *Genome Announcements*.
<https://doi.org/10.1128/genomeA.00199-15>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Draft Genome Sequences of Three *Capnocytophaga canimorsus* Strains Isolated from Healthy Canine Oral Cavities

Pablo Manfredi,^a Francesco Renzi,^{a,b} Guy R. Cornelis^{a,b}

Biozentrum der Universität Basel, Basel, Switzerland^a; University of Namur, Namur, Belgium^b

Here, we present the draft genome sequences of three strains of *Capnocytophaga canimorsus*, each isolated from a different dog's mouth. Genome analysis provided evidence that these organisms may belong to a different nonpathogenic subtype of *C. canimorsus*.

Received 12 February 2015 Accepted 28 April 2015 Published 28 May 2015

Citation Manfredi P, Renzi F, Cornelis GR. 2015. Draft genome sequences of three *Capnocytophaga canimorsus* strains isolated from healthy canine oral cavities. *Genome Announc* 3(3):e00199-15. doi:10.1128/genomeA.00199-15.

Copyright © 2015 Manfredi et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Guy R. Cornelis, guy.cornelis@unamur.be.

Capnocytophaga canimorsus is a capnophilic gliding Gram-negative bacterium from the *Bacteroidetes* phylum that is part of the normal flora of a dog's mouth (1, 2). It is not reported to cause infections in dogs, but it causes severe infections in humans who are in contact with dogs (3, 4). It has been estimated by culture-based methods that more than every other dog carries *C. canimorsus* in its normal oral flora (5). Because of the very specific culture conditions required by *C. canimorsus* strains, their prevalence has often been underestimated. A recent study using PCR-based methods reported that up to 74% of dogs carry *C. canimorsus* in their mouths (6). Interestingly, despite such a high prevalence, human infections remain very rare. Considering that in barely half of the reported cases, patients do not show any obvious risk factors (7), it is possible that different strains are not equally virulent and that only a tiny proportion is actually pathogenic for humans.

The three strains CcD38, CcD93, and CcD95 were isolated from canine oral swabs and partially characterized (5). The strains were selected as potential representatives of a subgroup of *C. canimorsus*, as suggested by 16S rRNA phylogenetics and various phenotyping assays (5, 8). Genomic DNA was extracted using the Genomic-tip 500/G DNA extraction kit (catalog no. 10262; Qiagen), according to the manufacturer's instructions, followed by an additional phenol-chloroform purification step. Sequencing was performed at LGC Genomics, Berlin, Germany, on one Illumina HiSeq 2000 channel and generated between 8.6 and 9.8 million 100-bp single reads per strain. Due to low mapping efficiency on the *C. canimorsus* 5 reference genome (9), *de novo* assembly was performed with Velvet with optimized *k*-mers (10). The sizes of the draft assemblies ranged from 2.56 to 2.74 Mb, with 191 (CcD95), 216 (CcD38), and 293 (CcD93) contigs. Genome annotation and preliminary analyses were performed by the LABGeM, France Génomique (11). The global G+C content (35.56% ± 0.07%) differs slightly but significantly from that of clinical isolates of *C. canimorsus* (36.16% ± 0.08%; *P* < 0.0002). A total of 2,325 to 2,618 coding sequences (CDSs) were identified per genome, giving a core genome of 1,896 orthologous groups. When considering the whole *Capnocytophaga* pangenome, 290 genes

were conserved and exclusively found in the three genomes presented here. This is similar to the number of *Capnocytophaga cynodegmi*-specific genes (341) (12) and supports the idea of a speciation event within the *C. canimorsus* taxon. Besides 247 clusters of unknown function, 16 genes implicated in basal energy metabolism, 10 in polysaccharide synthesis, 11 in integral outer membrane proteins, and 5 in ABC transporters formed predominant functional classes of the species-specific core. In addition, a consensual phylogenetic tree based on 771 individual phylogenies of orthologous proteins conserved among all *Capnocytophaga* spp. exhibited a very clear discrimination between clinical (clustering together in 677 trees) and canine oral (clustering together in 729 trees) isolates of *C. canimorsus*, even compared to a different species, such as *C. cynodegmi* (clustering together in 686 trees). This is again consistent with a potential speciation event within the *C. canimorsus* taxon.

Nucleotide sequence accession numbers. These whole-genome shotgun projects have been deposited in ENA under the accession numbers [CD0I00000000](https://ena.ebi.ac.uk/ena/record/CD0I00000000) (CcD38), [CDOL00000000](https://ena.ebi.ac.uk/ena/record/CDOL00000000) (CcD93), and [CDOH00000000](https://ena.ebi.ac.uk/ena/record/CDOH00000000) (CcD95). The versions described in this paper are the initial versions.

ACKNOWLEDGMENTS

This work was supported by grants 3100A0-128659 from the Swiss National Science Foundation and ERC 2011-ADG 20110310 from the European Research Council.

F.R. is a postdoctoral research fellow (chargé de recherche) of the Belgian Fonds National de la Recherche Scientifique (FNRS).

REFERENCES

- Brenner DJ, Hollis DG, Fanning GR, Weaver RE. 1989. *Capnocytophaga canimorsus* sp. nov. (formerly CDC group DF-2), a cause of septicemia following dog bite, and *C. cynodegmi* sp. nov., a cause of localized wound infection following dog bite. *J Clin Microbiol* 27:231–235.
- Shin H, Mally M, Meyer S, Fiechter C, Paroz C, Zaehring U, Cornelis GR. 2009. Resistance of *Capnocytophaga canimorsus* to killing by human complement and polymorphonuclear leukocytes. *Infect Immun* 77:2262–2271. <http://dx.doi.org/10.1128/IAI.01324-08>.
- Bobo RA, Newton EJ. 1976. A previously undescribed Gram-negative bacillus causing septicemia and meningitis. *Am J Clin Pathol* 65:564–569.

4. Butler T, Weaver RE, Ramani TK, Uyeda CT, Bobo RA, Ryu JS, Kohler RB. 1977. Unidentified Gram-negative rod infection. A new disease of man. *Ann Intern Med* 86:1–5. <http://dx.doi.org/10.7326/0003-4819-86-1-1>.
5. Mally M, Paroz C, Shin H, Meyer S, Soussoula LV, Schmiediger U, Saillen-Paroz C, Cornelis GR. 2009. Prevalence of *Capnocytophaga canimorsus* in dogs and occurrence of potential virulence factors. *Microbes Infect* 11:509–514. <http://dx.doi.org/10.1016/j.micinf.2009.02.005>.
6. Suzuki M, Kimura M, Imaoka K, Yamada A. 2010. Prevalence of *Capnocytophaga canimorsus* and *Capnocytophaga cynodegmi* in dogs and cats determined by using a newly established species-specific PCR. *Vet Microbiol* 144:172–176. <http://dx.doi.org/10.1016/j.vetmic.2010.01.001>.
7. Lion C, Escande F, Burdin JC. 1996. *Capnocytophaga canimorsus* infections in human: review of the literature and cases report. *Eur J Epidemiol* 12:521–533. <http://dx.doi.org/10.1007/BF00144007>.
8. Manfredi P, Lauber F, Renzi F, Hack K, Hess E, Cornelis GR. 2015. New iron acquisition system in *Bacteroidetes*. *Infect Immun* 83:300–310. <http://dx.doi.org/10.1128/IAI.02042-14>.
9. Manfredi P, Pagni M, Cornelis GR. 2011. Complete genome sequence of the dog commensal and human pathogen *Capnocytophaga canimorsus* strain 5. *J Bacteriol* 193:5558–5559. <http://dx.doi.org/10.1128/JB.05853-11>.
10. Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
11. Vallenet D, Belda E, Calteau A, Cruveiller S, Engelen S, Lajus A, Le Fèvre F, Longin C, Mornico D, Roche D, Rouy Z, Salvignol G, Scarpelli C, Thil Smith AA, Weiman M, Médigue C. 2013. MicroScope—an integrated microbial resource for the curation and comparative analysis of genomic and metabolic data. *Nucleic Acids Res* 41:D636–D647. <http://dx.doi.org/10.1093/nar/gks1194>.
12. Manfredi P, Renzi F, Cornelis GR. 2015. Draft genome sequences of three *Capnocytophaga cynodegmi* strains isolated from the oral cavity of healthy dogs. *Genome Announc* 3(3):e00200-15. <http://dx.doi.org/10.1128/genomeA.00200-15>.