

# **RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE**

### **Indol-2-yl ethanones as novel indoleamine 2,3-dioxygenase (IDO) inhibitors**

Dolušić, Eduard; Larrieu, Pierre; Blanc, Sébastien; Sapunaric, Frédéric; Norberg, Bernadette; Moineaux, Laurence; Colette, Delphine; Stroobant, Vincent; Pilotte, Luc; Colau, Didier; Ferain, Thierry; Fraser, Graeme; Galeni, Moreno; Frre, Jean Marie; Masereel, Bernard; Van Den Eynde, Benoît; Wouters, Johan; Frédérick, Raphaël Published in:

Bioorganic and Medicinal Chemistry

DOI: [10.1016/j.bmc.2010.12.032](https://doi.org/10.1016/j.bmc.2010.12.032)

Publication date: 2011

Document Version Early version, also known as pre-print

#### [Link to publication](https://researchportal.unamur.be/en/publications/549ce448-cb96-4da7-a8d4-0fdc0f746cb2)

Citation for pulished version (HARVARD):

Dolušić, E, Larrieu, P, Blanc, S, Sapunaric, F, Norberg, B, Moineaux, L, Colette, D, Stroobant, V, Pilotte, L, Colau, D, Ferain, T, Fraser, G, Galeni, M, Frre, JM, Masereel, B, Van Den Eynde, B, Wouters, J & Frédérick, R 2011, 'Indol-2-yl ethanones as novel indoleamine 2,3-dioxygenase (IDO) inhibitors', *Bioorganic and Medicinal* Chemistry, vol. 19, no. 4, pp. 1550-1561.<https://doi.org/10.1016/j.bmc.2010.12.032>

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

### AUTHOR QUERY FORM

<span id="page-1-0"></span>

Dear Author,

Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list.

For correction or revision of any artwork, please consult http://www.elsevier.com/artworkinstructions.

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Click on the 'Q' link to go to the location in the proof.



## BMC 8927 No. of Pages 1, Model 5G No. of Pages 1, Model 5G

#### Graphical abstract

#### Indol-2-yl ethanones as novel indoleamine 2,3-dioxygenase (IDO) inhibitors pp xxx–xxx

Eduard Dolušic´, Pierre Larrieu, Sébastien Blanc, Frédéric Sapunaric, Bernadette Norberg, Laurence Moineaux, Delphine Colette, Vincent Stroobant, Luc Pilotte, Didier Colau, Thierry Ferain, Graeme Fraser, Moreno Galeni, Jean-Marie Frère, Bernard Masereel, Benoît Van den Eynde, Johan Wouters, Raphaël Frédérick<sup>\*</sup>



**T** 

[Bioorganic & Medicinal Chemistry xxx \(2010\) xxx–xxx](http://dx.doi.org/10.1016/j.bmc.2010.12.032)

Contents lists available at [ScienceDirect](http://www.sciencedirect.com/science/journal/09680896)

Bioorganic & Medicinal Chemistry

journal homepage: [www.elsevier.com/locate/bmc](http://www.elsevier.com/locate/bmc)



### <span id="page-3-0"></span>Indol-2-yl ethanones as novel indoleamine 2,3-dioxygenase (IDO) inhibitors

Eduard Dolušić <sup>a</sup>, Pierre Larrieu <sup>b</sup>, Sébastien Blanc <sup>c</sup>, Frédéric Sapunaric <sup>d</sup>, Bernadette Norberg <sup>a</sup>, Laurence Moineaux <sup>a</sup>, Delphine Colette <sup>c</sup>, Vincent Stroobant <sup>b</sup>, Luc Pilotte <sup>b</sup>, Didier Colau <sup>b</sup>, Thierry Ferain <sup>c</sup>, Graeme Fraser <sup>c</sup>, Moreno Galeni <sup>d</sup>, Jean-Marie Frère <sup>d</sup>, Bernard Masereel <sup>a</sup>, Benoît <mark>Van den</mark> Eynde <sup>b</sup>, Johan Wouters <sup>a</sup>, Raphaël Frédérick <sup>a,</sup>\*

<sup>a</sup> Drug design and Discovery Center, University of Namur, 61 Rue de Bruxelles, 5000 Namur, Belgium <sup>b</sup> Ludwig Institute for Cancer Research, Université Catholique de Louvain, 74 Avenue Hippocrate, 1200 Brussels, Belgium  $\epsilon$  Euroscreen SA, 47 Rue Adrienne Boland, 6041 Gosselies, Belgium <sup>d</sup> Centre d'Ingénierie des Protéines, Université de Liège, Allée du 6 août, 4000 Liège, Belgium

#### article info

Article history: Received 1 October 2010 Revised 7 December 2010 Accepted 13 December 2010 Available online xxxx

Keywords: Indoleamine 2,3-dioxygenase IDO Anticancer Immunotherapy

#### **ABSTRACT**

Indoleamine 2,3-dioxygenase (IDO) is a heme dioxygenase which has been shown to be involved in the pathological immune escape of diseases such as cancer. The synthesis and structure–activity relationships (SAR) of a novel series of IDO inhibitors based on the indol-2-yl ethanone scaffold is described. In vitro and in vivo biological activities have been evaluated, leading to compounds with  $IC_{50}$  values in the micromolar range in both tests. Introduction of small substituents in the 5- and 6-positions of the indole ring, indole N-methylation and variations of the aromatic side chain are all well tolerated. An iron coordinating group on the linker is a prerequisite for biological activity, thus corroborating the virtual screening results.

© 2010 Published by Elsevier Ltd.

 $\overline{01}$ 

Immunotherapy is a promising novel strategy for cancer treatment. It consists of therapeutic vaccination of cancer patients in order to stimulate their (natural) immune system against cancer cells. This approach, however, showed limited efficacy in vivo. Indeed, cancer cells are able to develop enzymatic mechanisms allowing tumors to resist or escape immune rejection. Among the enzymes involved, the indoleamine 2,3-dioxygenase (IDO; EC 1.13.11.52) represents a crucial actor.<sup>1,2</sup>

IDO is a  $\sim$ 45 kDa monomeric extrahepatic cytosolic heme dioxygenase which catalyzes the initial and rate-limiting step in the catabolism of the essential amino acid tryptophan (Trp) along the kynurenine pathway, the de novo biosynthetic route leading to nicotinamide adenine dinucleotide  $(NAD)$ .<sup>3,4</sup> IDO is a more promiscuous enzyme than the catalytically related hepatic enzyme tryptophan 2,3-dioxygenase (TDO; EC 1.13.11.11). Apart from both tryptophan enantiomers, IDO substrates are indoleamines such as serotonin, melatonin, and tryptamine.<sup>[5](#page-14-0)</sup> In order for IDO to be catalytically active, it is essential to maintain the heme iron ion in the ferrous ( $Fe^{2+}$ ) state. As the enzyme is susceptible to autoxidation, a reductant is necessary for preserving the enzymatic activity. Cytochrome  $b_5$  has been demonstrated to play this role in vivo.<sup>6,7</sup>

By depleting Trp locally, IDO blocks the proliferation of T lymphocytes, which are sensitive to Trp shortage. $8$  The downstream catabolites of the kynurenine pathway exacerbate this immunosuppressive effect<sup>1</sup> and are also involved in other pathological conditions, particularly in the nervous system. $9-11$ 

Many human tumors constitutively express IDO.<sup>[12](#page-14-0)</sup> It has been shown that an increased level of IDO expression in tumor cells is correlated with poor prognosis for survival in cancer.<sup>[13,14](#page-14-0)</sup> This led to the hypothesis that IDO inhibition might enhance the efficacy of cancer treatments. Indeed, results from in vitro and in vivo studies have suggested an improvement of the efficacy of therapeutic vaccination or chemotherapy by concomitant administration of 70 an IDO inhibitor thus highlighting IDO as an attractive target[.12,13,15–17](#page-14-0)

Until recently, the best known IDO inhibitors<sup>[18](#page-14-0)</sup> displayed affinities in the micromolar range and comprised mainly Trp derivatives such as 1-methyltryptophan  $(1MT)^{19}$  $(1MT)^{19}$  $(1MT)^{19}$  (K<sub>i</sub> = 37 µM) or  $\beta$ -carbolines ( $K_i \sim 0.12$  mM).<sup>20</sup> Other indole-based IDO inhibitors, such as  $br>{bra}$  $binom{21,22}{1}$  and methylthiohydantoin-tryptophan  $(MTH-Trp)^{13}$  $(MTH-Trp)^{13}$  $(MTH-Trp)^{13}$  have been described. The best inhibitor of the brassinin class has a  $K_i$  of 12  $\mu$ M but is not Lipinski-compliant due to high hydrophobicity. $22$ 

In 2006, potent IDO inhibitors were isolated from marine invertebrate extracts, $23,24$  leading to two new series of uncompetitive nanomolar inhibitors.[26,27](#page-14-0) However, some of the most active compounds of the original series revealed to be inactive in a

0968-0896/\$ - see front matter © 2010 Published by Elsevier Ltd. doi[:10.1016/j.bmc.2010.12.032](http://dx.doi.org/10.1016/j.bmc.2010.12.032)

<sup>⇑</sup> Corresponding author. Tel.: +32 81 72 42 90; fax: +32 81 72 42 38. E-mail address: [Raphael.frederick@fundp.ac.be](mailto:Raphael.frederick@fundp.ac.be) (R. Frédérick).

2 E. Dolušić et al. / Bioorg. Med. Chem. xxx (2010) xxx–xxx

yeast-based cellular assay, suggesting poor cell penetration.<sup>[7](#page-14-0)</sup> This test nevertheless provided two new micromolar inhibitors with unknown modes of action.

In 1989, 4-phenylimidazole (PIM) was identified as a modestly potent IDO inhibitor.[4,19](#page-14-0) Despite the uncompetitive inhibition kinetics, the authors showed through spectroscopic studies that PIM binds into the active site of IDO. Moreover, they discovered a preferred binding of PIM to the inactive ferric ( $Fe<sup>3+</sup>$ ) form of the enzyme, which may explain this apparent discrepancy. The first crystal structure of human IDO complexed with the inhibitor reported later<sup>25</sup> confirmed the binding of PIM in the active site. A recently undertaken structure-based study<sup>26</sup> of PIM-derived molecules prompted by Sugimoto's work confirmed the antecedent findings while providing access to novel inhibitors with low micromolar potencies.

<sup>100</sup> Recent developments in the field include hypoxic-neoplastic cells-targeting 1MT-tirapazamine hybrids,<sup>27</sup> novel inhibitors obtained from marine sediments (IC<sub>50</sub>  $\approx$  2  $\mu$ M), $^{28}$  potent competitive inhibitors including a hydroxyamidine chelating motif  $(IC_{50}$  values down to 60 nM), $^{29}$  strong inhibition by modification of cysteine residues by ebselen,<sup>30</sup> sub-micromolar S-benzylisothiourea-based inhibitors active in cells,  $31$  various potent inhibitors (IC<sub>50</sub> values down to 0.2  $\mu$ M) discovered by a combination of a docking-based pharmacophore model development and fragment-based drug design (FBDD). $32$ 

110 Despite all these efforts, modestly potent **1MT**  $(K_i = 34 \mu M)^{19}$  $(K_i = 34 \mu M)^{19}$  $(K_i = 34 \mu M)^{19}$ remains the only compound to have been passed into clinical trials as an IDO inhibitor so  $far.^{33}$  $far.^{33}$  $far.^{33}$  This prompted us to search for new scaffolds that could be further optimized with the objective of identifying potent and selective IDO inhibitors. Briefly, a collection of  $\sim$  62,000 structures was downloaded from the ZINC website using Lipinski-style rules for fragments (molecular weight  $\leq$  250,  $Log P \le 2.5$ , number of rotatable bonds  $\le 5$ ).<sup>34</sup> These compounds were docked inside the IDO active site and ranked according to their best potential binding. From the 39 compounds that were se-<sup>120</sup> lected, bought and assayed on IDO, six derivatives displayed an inhibitory potency  $>30\%$  at a concentration of 100 µM. Among these, compound  $7a$ , characterized by an indole nucleus substituted in the 2-position by a 2-pyridinyl-ethanone function, was our best hit (IDO IC<sub>50</sub>  $\sim$  65 µM), with an inhibitory potency in the same range as the reference compound 1MT. (Table 1). Herein we report the synthesis and structure–activity study of compound 7a.

An initial docking study of 7a within the IDO binding cleft was performed by means of the automated GOLD program<sup>35</sup> and re-<sup>130</sup> vealed essential features stabilizing 7a in the IDO binding cleft and particularly that (i) the oxygen atom of the ketone function, being situated  $\sim$ 2 Å above the plane of the heme, coordinates to the heme iron,  $(i)$  the indole ring is found to be stabilized in the so-called lipophilic A-pocket of IDO.<sup>32</sup> Indeed, this stabilization of the indole residue in Pocket A was already suggested by other



Inhibition of IDO activity by 1MT and 7a identified using VS



groups to explain substrate binding as well as interaction of 1MT. In this position, the 3-pyridyl group is projected towards the entrance of the active site and is stabilized in the aromatic Pocket B through T-shape interactions with Phe163 and Phe226 (Fig. 1).  $140$ 

Our structural modifications of 7a were focused on exploiting three binding interactions with IDO: (1) the interior of the active site, in particular interactions with C129 and S167 in Pocket A; (2) the active site entrance region (Pocket B) occupied by the N-cyclohexyl-2-aminoethanesulfonic acid (CHES) buffer molecule in a crystal structure; $^{25}$  $^{25}$  $^{25}$  (3) the heme iron binding group. According to the binding orientation of 7a we hoped to achieve interactions in the interior region of the active site (Pocket A) by introducing substituents on the indole ring. The substituents were chosen with a view to balance on one side the lipophilic character of this Pocket 150 introducing groups such as fluorine, bromine, chlorine, methyl; and, on the other side the possibility of forming H-bond interactions with specific amino acids with such substituents as hydroxyl, methoxy, nitro, or trifluoromethoxy. The interaction within Pocket B would be probed by replacement of the 3-pyridyl group of 7a with other (hetero)aromatics. The heme iron binding interaction would be explored by replacement of the ethanone moiety with other linkers between the indole and the aromatic side chain.

#### 2. Chemistry

Initial efforts to prepare analogues of compound **7a** included at-<br>160 tempts to react indole-2-acyl chlorides [\(Scheme 1,](#page-5-0) 3,  $R'' = Cl$ ) with reagents derived from 3-bromomethylpyridine hydrobromide under various conditions.<sup>36-38</sup> In fact, all of these approaches revealed to be unsatisfactory, the main difficulty likely being the intrinsic instability of the in situ generated free base 3-bromomethylpyridine which readily oligomerizes. $39$  The only strategy allowing access to 7a analogues was the approach originally developed by Sundberg et al., as outlined in [Scheme 1](#page-5-0). This method involves the LDA effected deprotonation of alkylarene 1 followed by quenching of the in situ generated carbanion with a reactive indole 170 carbonyl compound (3 and 6) at 0 °C. Stirring overnight while slowly attaining room temperature followed by aqueous workup, chromatographic purification and crystallization provided the targeted compounds.



Figure 1. View of the lead compound 7a inside the IDO binding cleft. (Picture made with  $Pymol<sup>34</sup>$  $Pymol<sup>34</sup>$  $Pymol<sup>34</sup>$ ).

Please cite this article in press as: Dolušić, E.; et al. Bioorg. Med. Chem. (2010), doi[:10.1016/j.bmc.2010.12.032](http://dx.doi.org/10.1016/j.bmc.2010.12.032)

### <span id="page-5-0"></span>BMC 8927 No. of Pages 13, Model 5G



Scheme 1. General synthetic scheme for indol-2-yl ethanones. Reagents and conditions: (i) LDA, THF/hexanes, -78-0 °C, 1 h; (ii) SOCl<sub>2</sub>,  $\Delta$ , 15 min; (iii) THF/hexanes, 0 °C to rt, 16-24 h; (iv) N<sub>3</sub>CH<sub>2</sub>CO<sub>2</sub>Et, NaOEt, EtOH, -10-4 ℃, 1.5-20 h; (v) hexane, 200 ℃ (µW), 5 min; (vi) MeI, K<sub>2</sub>CO<sub>3</sub>, DMF, 80 ℃, 16 h to 5 days; (vii) AlCl<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, 0 ℃ to rt, 24 h; (viii) HCO<sub>2</sub>NH<sub>4</sub>, Pd black, MeOH, rt, 1 h; (ix) H<sub>2</sub> (1 atm), 3% Pd/C, EtOH, rt, 45 min; (x) PhB(OH)<sub>2</sub>, Pd(PPh<sub>3</sub>)<sub>4</sub>, K<sub>2</sub>CO<sub>3</sub>, EtOH/toluene 1:1,  $\Delta$ , 20 h.

Accordingly, most of the compounds 7 as well as 8a, 9a, 10a, and 11a were prepared from indole esters  $(3, R'' = OMe, OEt)$  or acyl chlorides  $(3, R'' = Cl)$  when these starting material were commercially available. Otherwise, the necessary indole acyl chloride 3 was prepared by reaction of the commercially available indole-2- <sup>180</sup> carboxylic acid 2 in refluxing thionyl chloride. Non-commercially available indole esters 3 were made using the Hemetsberger–Knittel procedure starting from the corresponding  $\alpha$ -azidocinnamates 5, which were in turn made by condensation of the substituted benzaldehydes 4 with ethyl azidoacetate under Knoevenagel conditions (Scheme 1). $40,41$  It should be noted that compound 11a (Scheme 1) was prepared in 37% yield using commercial Grignard reagent in lieu of the LDA procedure. Moreover, during the course of this reaction, the tertiary alcohol 16 (inactive in the biological tests; data not shown) was also isolated in 17% yield.

<sup>190</sup> Palladium catalyzed reductions were used to convert benzyloxy-substituted 2-ketoindoles 7n and 7r to their debenzylated counterparts 7o and 7s, respectively. Bis-alcohol 17 was also isolated as a byproduct during the catalytic reduction of 7n. The 5-phenylindole derivative 7l was prepared from the corresponding 5-bromo analogue 7h by a Suzuki cross-coupling with phenylboronic acid (Scheme 1). Finally, after initial unsuccessful attempts to demethylate 8d with boron tribromide (which seemed to cause product decomposition), the 4-hydroxyindole 8e was obtained by reaction of 8d with aluminum trichloride in <sup>200</sup> dichloromethane.

For the synthesis of the N-methylated analogue 12a, the required N-methylated indole ester 6 was synthesized by dimethylation of N-unsubstituted indole-2-carboxylic acid 2 (Scheme 1).<sup>[42](#page-14-0)</sup>

The catalytic hydrogenation of 7a afforded the hydroxyl derivative 18 (Scheme 2). Compound 19 bearing an ethylene linker between the two aromatic rings was prepared from 7a by a



Scheme 2. Synthesis of compounds bearing different linker groups. Reagents and conditions: (i) HCO<sub>2</sub>NH<sub>4</sub>, Pd black, MeOH, rt, 3 days; (ii) H<sub>2</sub>NNH<sub>2</sub>, KOH, (CH<sub>2</sub>OH)<sub>2</sub>,  $\mu$ W, 1 h then aq NH<sub>4</sub>Cl; (iii) NH<sub>2</sub>OH·HCl, pyridine, EtOH,  $\mu$ W (120 °C), 30 min; (iv) SOCl<sub>2</sub>,  $\Delta$ , 15 min then 3-aminopyridine, DIPEA, THF, 0 °C $\rightarrow$ rt, 30 min.

Please cite this article in press as: Dolušić, E.; et al. Bioorg. Med. Chem. (2010), doi:[10.1016/j.bmc.2010.12.032](http://dx.doi.org/10.1016/j.bmc.2010.12.032)

## <span id="page-6-0"></span>BMC 8927 No. of Pages 13, Model 5G

4 E. Dolušic´ et al. / Bioorg. Med. Chem. xxx (2010) xxx–xxx

Wolff–Kishner reduction. Oxime 20 was obtained by reacting ketone 7a with hydroxylamine. Finally, amide 21 was made from acid 2 by reaction with thionyl chloride followed by nucleophilic <sup>210</sup> substitution with 3-aminopyridine [\(Scheme 2\)](#page-5-0).

The structure of all compounds was confirmed by means of  $^1\mathrm{H}$ ,  $13C$  and (where applicable)  $19F$  NMR, LC–MS, and elemental analysis (see Section 5). In addition, single crystal X-ray diffraction was used to corroborate the structures obtained in these series. The molecular structures of compounds 7a and 7m with displacement ellipsoids drawn at the 50% probability levels are depicted in Figure 2. It should be noted that for 7m thermal agitation of the trifluoromethoxy side chain led to large displacement ellipsoids for the fluorine atoms.

#### <sup>220</sup> 3. Results

#### 3.1. IDO inhibition through interactions in the interior of the active site (Pocket A)

Detailed analysis of the binding conformation of 7a inside the IDO binding cleft suggests that introduction of small substituents around the indole scaffold could be beneficial for IDO inhibition.

Starting from our hit **7a** (IC<sub>50</sub> = 65  $\mu$ M), we first explored the influence of substitution around the indole ring. From Table 2 it can be observed that introduction of substituents in the 3- (7b), 4- (7c–e), and 7-position (7q) does not generally afford more <sup>230</sup> potent compounds than the hit 7a. Substitutions in the 5- and 6-positions are well tolerated and more potent compounds are obtained. Particularly, introducing a trifluoromethoxy group on the 5-position of the indole ring led to a fivefold improvement in the IDO inhibitory potency. Regarding the disubstitution, introductions of substituents in the 4,5- or the 4,6-positions do not drastically affect IDO inhibition. Finally,  $N$ -methylation of the indole nucleus seems to be tolerated as 12a is in vitro twofold more potent than parent compound 7a.

#### 3.2. IDO inhibition through interactions with Pocket B of the <sup>240</sup> active site

We next explored the influence of modifying the fragment interacting with Pocket B of the IDO active site. From Table 3, it can be observed that introductions of an unsubstituted phenyl (11a), a 2-pyridyl (8a) or a 2-pyrazinyl group (10a) in lieu of the 3-pyridyl moiety of 7a slightly improve the IDO inhibitory potency ( $IC_{50}$ 's of 29, 37, and 26  $\mu$ M, respectively). On the contrary, introduction of a  $\overline{4}$ -pyridyl moiety (**9a**) afforded an inactive compound.

#### 3.3. IDO inhibition through interaction with the heme iron binding group

<sup>250</sup> Finally we appraised the importance of the ketone moiety of compound 7a in establishing an interaction with the heme iron. Indeed, reduction of this group (18) or its removal altogether (19) led to inactive compounds thus confirming the requirement for an iron coordinating group ([Table 4](#page-7-0)). Oxime 20 is endowed with an in vivo inhibitory potency similar to that of ketone 7a but surprisingly,



Figure 2. The molecular structures of (a) 7a and (b) 7m. Displacement ellipsoids are drawn at the 50% probability levels. (Pictures made with MERCURY2.3)

Table 2

IDO inhibitory potency of compounds 7a–u, 12a





⁄⁄ NT = not tested.

 $NI = no$  inhibition.

#### Table 3

IDO inhibitory potency of compounds 7a–11a



\* NI = no inhibition.

this compound is not active in vitro. This might result from a poor stability of the oxime moiety in the experimental conditions of the assay. Methylating the ethanone link  $(7v)$  or its replacement with an amide (21) afforded less potent compounds.

#### **4. Discussion and conclusion** 260

About 30 original compounds were synthesized around the indol-2-yl ethanone scaffold and their in vivo and in vitro inhibitory potency against indoleamine 2,3-dioxygenase (IDO) was evaluated. Starting from the hit (1H-indol-2-yl)-2-pyridin-3-

Please cite this article in press as: Dolušić, E.; et al. Bioorg. Med. Chem. (2010), doi[:10.1016/j.bmc.2010.12.032](http://dx.doi.org/10.1016/j.bmc.2010.12.032)

[Q3](#page-1-0)

### <span id="page-7-0"></span>BMC 8927 No. of Pages 13, Model 5G

#### E. Dolušić et al. / Bioorg. Med. Chem.  $xxx(2010)$   $xxx-xxx$  5



IDO inhibitory potency of compounds 7a, 7v, 18–21



NI = no inhibition.

yl-ethanone (7a) identified through a virtual screening procedure and possessing an  $IC_{50}$  of 65  $\mu$ M in the in vitro test, a fivefold improvement in potency ( $IC_{50} = 13 \mu M$ ) could be achieved by introducing a trifluoromethoxy group in the 5-position of the indole nucleus. Most small substituents around the indole moiety 270 led to compounds of roughly equivalent potency, indicating room for further modifications but also a probable lack of close interactions of the inhibitor molecules with the IDO Pocket A interior as described by Rohrig et al. $32$  A similar generalization can be applied to the other aromatic moiety and its putative interactions with Pocket B of the IDO. Altogether, it seems no strong H-bond interactions with any of the hydrogen bonding residues in the IDO active site could be established. The presence of an iron chelating group seems to be of crucial importance, since its deletion leads to loss of potency.

<sup>280</sup> A number of compounds investigated here possess a moderate in vivo activity. The compounds 7f and 7g (with small halogen atoms in the 5-position of the indole moiety) are both endowed with 24% IDO inhibition at a 20  $\mu$ M concentration in the cellular assay, which indicates reasonable cell penetration. A very similar value was obtained for oxime 20.

In summary, a novel series of compounds with a moderate inhibitory activity against IDO has been identified. Both in vitro and in vivo assays show promising results. Further modifications aiming to achieve more favorable interactions with the enzyme ac-<sup>290</sup> tive site are in due course.

#### 5. Experimental part

#### 5.1. Chemistry

#### 5.1.1. General experimental procedures

Chemicals were purchased from commercial suppliers (Acros, Sigma–Aldrich, Maybridge, Apollo) and used without further purification. All solvents were used as purchased from Acros, Sigma–Aldrich, Fisher Scientific and Biosolve. All reactions were performed under an inert argon (Alphagaz 2) atmosphere, unless stated otherwise. Melting points were determined in open capillaries on a Büchi B-540 capillary melting point apparatus and are 300 uncorrected.

Thin layer chromatography (TLC) was performed on silica gel plates ( $60F_{254}$ , 0.2 mm thick, Merck) eluting with gradients of ethyl acetate in cyclohexane. The spots were visualized by ultraviolet light (wavelengths 254 and 366 nm).

<sup>1</sup>H NMR spectra were recorded in  $DMSO-d_6$  solutions on a Jeol JNM EX 400 machine at 400 MHz with tetramethylsilane (TMS) as internal standard.  $^{13}C$  and  $^{19}F$  NMR spectra were recorded on the same machine in  $DMSO-d_6$  solutions at 100 MHz and 376.2 MHz, respectively. Chemical shifts  $(\delta)$  are reported in ppm 310 downfield from TMS.

Elemental analyses (C, H, N, S) were performed on a Thermo Finnigan FlashEA 1112 apparatus.

Analytical LC–MS analyses were performed on an Agilent 1100 Series LC–MSD Trap system using UV detection at 254 and 366 nm. MS data were recorded using electrospray ionization (ESI) operating in positive mode. The following two methods were applied: method 1: injection of 10  $\mu$ l of a 20  $\mu$ g/mL acetonitrile solution onto a C18-3.5 µm Zorbax SB column (100  $\times$  3 mm); separation using a gradient (flow rate:  $0.5$  mL/min) of acetonitrile in acetic  $320$ acid 0.1% ( $v/v$ ) in water: 5–95% acetonitrile during 5 min, holding for 3 min, then reversing to 5% acetonitrile during 0.1 min and holding for an additional 5.4 min. Method 2: injection of 10  $\mu$ l of a 2  $\mu$ g/mL acetonitrile solution onto a C18-5  $\mu$ m Agilent LiChrospher 100 column (250  $\times$  4 mm); separation using a gradient (flow rate: 1.0 mL/min) of acetonitrile in acetic acid 0.1% ( $v/v$ ) in water: 20–80% acetonitrile during 20 min, holding for 20 min, then reversing to 20% acetonitrile during 0.1 min and holding for an additional 5 min.

Automated flash chromatography was performed on the 330 Biotage AB (Uppsala, Sweden) SP1 system equipped with prepacked flash KP-Sil silica cartridges eluting with gradients of ethyl acetate in cyclohexane. The peaks were detected by UV absorption at 254 and 320 nm.

The yields refer to combined yields of chromatographically and spectroscopically pure (>95%) compounds and the crops obtained by evaporation of mother liquors. All new compounds were determined to be >95% pure by LC–MS and NMR.

6 E. Dolušic´ et al. / Bioorg. Med. Chem. xxx (2010) xxx–xxx

The microwave-assisted syntheses were carried out in an Initi-<sup>340</sup> ator 16 single-mode microwave instrument producing controlled irradiation at 2.450 GHz (Biotage AB). Reaction times refer to hold times at the temperatures indicated, not to total irradiation times. The temperature was measured with an IR sensor on the outside of the reaction vessel.

#### 5.1.2. Methods for preparation of indolic intermediates (3 and 6)

5.1.2.1. Indole-2-acyl chlorides  $(3, R'' = Cl)$ . Commercial indole-2-carboxylic acid was dissolved in thionyl chloride  $(\sim 1.5 \text{ mL})$ mmol) and the suspension was boiled for 15 min, during which <sup>350</sup> time typically a dark green solution was formed. Excess thionyl chloride was removed under reduced pressure and the solid residue was suspended in toluene and the suspension was evaporated to dryness (toluene treatment was repeated three times). The green-grayish solid residues were used without further purification.

5.1.2.2. Indolic esters by Hemetsberger–Knittel synthesis (3,  $R'' = OEt$ <sup>40,41</sup>. Sodium (4.0 equiv) was dissolved in absolute ethanol (0.5 mL/mmol) and the solution was cooled to  $-10$  °C in an ice/ acetone bath. A corresponding benzaldehyde (1.0 equiv) was dissolved in a commercial 25% solution of ethyl azidoacetate in ethanol (2.1 g per mmol of starting benzaldehyde, 4.0 equiv) and this solution was added dropwise to the sodium ethoxide solution during 30 min while the bath temperature was kept at  $-10$  °C. The reaction mixtures typically turned into bright yellow suspensions. The stirring was continued until TLC (developing with 15% EtOAc in cyclohexane) indicated completion or the reaction (1.5–2.5 h). If the reaction required more than 1.5 h, the temperature was allowed to rise to  $0^{\circ}$ C. In one case (5r), the reaction had to be run overnight. The mixtures were partitioned between ethyl acetate 370 and water. The aqueous phase was extracted twice with ethyl acetate and the organic layers were pooled, washed with brine, dried over magnesium sulfate, and concentrated under reduced pressure to provide crude azidoacrylic esters (5) as yellow to orange oils or liquids in a 46–100% yield. These products were suspended in nhexane (6–9 mL/mmol of crude azidoacrylic ester) and subjected to microwave heating at 200 °C for 5 min  $(P = 100-140 \text{ W}$ ,  $p = 12-13$  bar). The resulting residues were taken into ethyl acetate and evaporated with silica gel. The corresponding indolic esters 3 (yields 19–93%) were purified by chromatography. In one case, a mixture of isomers was obtained. 2-Azido-3-(3-methoxyphenyl)acrylic acid ethyl ester gave 5-methoxy-1H-indole-2-carboxylic acid ethyl ester (3j) and 7-methoxy-1H-indole-2-carboxylic acid ethyl ester (3q) in a 1:1 ratio (each in a 47% yield).

5.1.2.3. N-Methylated indolic ester; methyl 1-methyl-1H-indole-

**2-carboxylate (6).** Prepared according to a literature procedure. $42$ Methyl iodide (5.0 mL, 80.0 mmol) was added to a stirred suspension of anhydrous potassium carbonate (6.3 g, 45.6 mol) and 1Hindole-2-carboxylic acid (2; 5.3 mmol) in dry DMF (6 mL) and the mixture was stirred at 80 $^{\circ}$ C until the reaction was complete as 390 indicated by TLC and/or LC–MS (3–5 days). Upon cooling to room temperature, the suspension was partitioned between ethyl acetate and water. The aqueous phase was extracted with ethyl acetate and the combined organic layers were dried over MgSO<sub>4</sub>, concentrated under reduced pressure and dried to yield the desired N-methylated indole ester (6) as a beige solid in a 96–99% yield.

#### 5.1.3. General synthetic procedure for 1-(1H-indol-2-yl)-2-arylketones (7–10, 12) $^{43}$

An oven-dried flask was purged with argon while hot, then allowed to cool down to room temperature under argon and <sup>400</sup> charged with dry THF (2 mL/mmol of starting indole 3/6) and diiso-

ice/isopropanol bath and n-butyl lithium, 1.6 M solution in hexanes (4.0 equiv), was added to give a bright yellow solution. The mixture was stirred for 30 min below  $-60$  °C whereupon a solution of picoline or another appropriate reagent with an ionizable methyl(ene) group  $(4.0 \text{ equiv})$  in THF  $(2 \text{ mL/mm})$  of starting indole 3 or 6) was added to give a yellow to red colored mixture. During a further 30 min, the bath temperature was allowed to rise to approx.  $0^{\circ}$ C during which time the reaction mixture became a yellow to red suspension. The temperature was then kept at  $0^{\circ}C = 410$ (using an ice/water bath) for an additional 30 min. At this point, a solution of the indole-2-carbonyl compound  $3/6$  (1.0 equiv) in THF (4 mL/mmol) was added dropwise over the flask wall. The reaction mixture darkened (dark purple to dark brown color in most cases) and was then allowed to attain room temperature overnight. At 18 $h$  (unless otherwise stated) after the addition of the indole component, the reaction was quenched with saturated aqueous ammonium chloride solution (10 mL/mmol of starting compound 3/6) and stirred for several minutes. Upon partition between ethyl acetate and water, the aqueous phase was extracted 420 twice with dichloromethane and the combined organic layers were dried over MgSO4. The title products were purified by chromatography (the products typically eluted with 70–80% of ethyl acetate). The products were then precipitated by concentration of the pooled column fractions combined with the addition of cyclohexane. The precipitates formed were filtered off, washed twice with cyclohexane and dried at 40 $\degree$ C in vacuo to yield analytically pure samples. The following compounds were prepared in this way:

propylamine (4.0 equiv). The solution was cooled to  $-78$  °C in a dry

5.1.3.1. 1-(1H-Indol-2-yl)-2-pyridin-3-yl-ethanone  $(7a)^{43-45}$ . Prepared from commercial 1H-indole-2-carboxylic acid ethyl ester 3a (986 mg, 5.05 mmol) and 3-picoline. Yellowish solid (699 mg, 59%), mp 177-180 °C. LC-MS  $R_t$  5.2 min (method 1), 7.0 min (method 2);  $m/z$  [MH<sup>+</sup>] 237. <sup>1</sup>H NMR  $\delta$  11.76 (1H, br s, indole-H1), 8.53 (1H, d,  $J = 1.7$  Hz, ArH), 8.43 (1H, dd,  $J = 4.8$ , 1.7 Hz, ArH), 7.71 (1H, d,  $J = 7.8$  Hz, ArH), 7.69 (1H, d,  $J = 7.8$  Hz, ArH), 7.56 (1H, d,  $J = 1.8$  Hz, ArH), 7.41 (1H, d,  $J = 8.2$  Hz, ArH), 7.33 (1H, dd,  $J = 7.8$ , 4.8 Hz, ArH), 7.26 (1H, t,  $J = 7.6$  Hz, ArH), 7.06 (1H, t, J = 7.6 Hz, ArH), 4.36 (2H, s, CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$  190.1 (C=O), 151.1 (ArC), 148.3 (ArC), 138.6 (ArC), 137.8 (ArC), 135.4 (ArC), 131.6 (ArC), 127.4 (ArC), 126.2 (ArC), 124.0 (ArC), 123.3 (ArC), <sup>440</sup> 120.9 (ArC), 113.3 (ArC), 110.7 (ArC), 42.0 (CH<sub>2</sub>). Anal. Calcd for  $C_{15}H_{12}N_2O$ : C, 76.25; H, 5.12; N, 11.86. Found: C, 76.16; H, 5.13; N, 11.87.

5.1.3.2. 1-(3-Bromo-1H-indol-2-yl)-2-pyridin-3-yl-ethanone (7b). Prepared from commercial 3-bromo-1H-indole-2-carboxylic acid (267 mg, 1.09 mmol) and 3-picoline via corresponding acyl chloride 3b (method A). Bright yellow solid (100 mg, 29%), mp 194– 195 °C (dec). LC–MS  $R_t$  6.0 min (method 1), 12.8 min (method 2);  $m/z$  [MH<sup>+</sup>] 315; 317. <sup>4</sup>H NMR  $\delta$  12.19 (1H, br s, indole-H1), 8.49 (1H, d, J = 1.8 Hz, ArH), 8.45 (1H, dd, J =  $4.\overline{8}$  Hz, 1.6, ArH), 7.69  $450$  $(1H, d, J = 7.8$  Hz, ArH), 7.57  $(1H, d, J = 8.0$  Hz, ArH), 7.49  $(1H, d, J = 1.6)$ J = 8.2 Hz, ArH), 7.38–7.33 (2H, overlapped, 2 ArH), 7.19 (1H, t,  $J = 7.6$  Hz, ArH),  $\overline{4.53}$  (2H, s, CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$  189.3 (C=O), 151.4 (ArC), 148.4 (ArC), 138.2 (ArC), 136.5 (ArC), 132.1 (ArC), 130.8 (ArC), 127.6 (ArC), 127.3 (ArC), 123.9 (ArC), 122.0 (ArC), 121.2 (ArC), 113.7 (ArC), 96.5 (ArC), 44.4 (CH<sub>2</sub>). Anal. Calcd for  $C_{15}H_{11}BrN_2O$ : C, 57.16; H, 3.52; N, 8.89. Found: C, 58.26; H, 3.57; N, 8.48.

5.1.3.3. 1-(4-Fluoro-1H-indol-2-yl)-2-pyridin-3-yl-ethanone (7c). Prepared from 4-fluoro-1H-indole-2-carboxylic acid ethyl ester  $3c$ (method B; 184 mg, 0.89 mmol) and 3-picoline. Yellow half-solid (15 mg, 7%). LC–MS  $R_t$  5.4 min (method 1);  $m/z$  [MH<sup>+</sup>] 255. <sup>1</sup>H NMR  $\delta$  12.10 (1H, br s, indole-H1), 8.53 (1H, br s, ArH), 8.43 (1H,

Please cite this article in press as: Dolušic´, E.; et al. Bioorg. Med. Chem. (2010), doi[:10.1016/j.bmc.2010.12.032](http://dx.doi.org/10.1016/j.bmc.2010.12.032)

E. Dolušić et al. / Bioorg. Med. Chem. xxx (2010) xxx–xxx 7 7

d, J = 4.1 Hz, ArH), 7.73–7.65 (2H, overlapped, 2 ArH), 7.33 (1H, dd, J = 7.2, 5.1 Hz, ArH), 7.24 (2H, overlapped, 2 ArH), 6.85 (1H, m, ArH), 4.40 (2H, s, CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$  190.2 (C=O), 167.5 (ArC), 149.8 (d,  $J_{CF}$  = 278 Hz, indole-C4), 137.8 (ArC), 135.7 (ArC), 132.2 (d, overlapped, ArC), 132.1 (ArC), 129.2 (ArC), 126.8 (d,  $J_{CF}$  = 10 Hz, ArC), 124.0 (ArC), 116.9 (d,  $J_{CF}$  = 22 Hz, ArC), 109.9 (ArC), 106.0 470 (ArC), 105.0 (d,  $J_{CF}$  = 18 Hz, ArC), 42.0 (CH<sub>2</sub>).

5.1.3.4. 1-(4-Methoxy-1H-indol-2-yl)-2-pyridin-3-yl-ethanone (7d). Prepared from commercial 4-methoxy-1H-indole-2-carboxylic acid methyl ester 3d (268 mg, 1.29 mmol) and 3-picoline. White solid (238 mg, 69%), mp 159-160 °C (dec). LC-MS  $R_t$ 5.3 min (method 1);  $m/z$  [MH<sup>+</sup>] 267. <sup>1</sup>H NMR  $\delta$  11.75 (1H, br s, indole-H1), 8.52 (1H,  $\frac{1}{2}$  pr s, ArH), 8.42 (1H, d, J = 4.9 Hz, ArH), 7.70 (1H, d, J = 7.8 Hz, ArH), 7.57 (1H,  $\frac{\text{br}}{\text{s}}$ , ArH), 7.32 (1H, dd, J = 7.6, 4.9 Hz, ArH), 7.17 (1H, t, J = 8.1 Hz, ArH), 6.97 (1H, d, J = 8.1 Hz, ArH), 6.51 (1H, d, J = 7.6 Hz, ArH), 4.33 (2H, s, CH<sub>2</sub>), 3.88 (3H, s, OCH<sub>3</sub>). <sup>13</sup>C 480 NMR  $\delta$  189.7 (C=O), 154.9 (ArC), 151.1 (ArC), 148.3 (ArC), 139.9 (ArC), 137.7 (ArC), 134.3 (ArC), 131.8 (ArC), 127.5 (ArC), 123.9 (ArC), 118.8 (ArC), 108.3 (ArC), 106.1 (ArC), 100.0 (ArC), 55.6 (OCH<sub>3</sub>), 41.8 (CH<sub>2</sub>). Anal. Calcd for C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>2</sub>: C, 72.16; H, 5.30; N, 10.52. Found: C, 72.36; H, 5.30; N, 10.36.

5.1.3.5. 1-(4-Hydroxy-1H-indol-2-yl)-2-pyridin-3-yl-ethanone (7e). A solution of methoxyindole 7d (143 mg, 0.54 mmol) in dry dichloromethane (5 mL) was cooled to 0  $\degree$ C and anhydrous aluminum chloride (247 mg, 1.84 mmol) was added to give a yellow suspension. Room temperature was slowly attained while the color of the reaction mixture gradually changed to dark red. After 19 h, as TLC still indicated presence of the starting compound, another aliquot of aluminum chloride and 2 mL of dichloromethane were added. At  $\overline{24}$  h of total reaction time (the majority of the starting compound still being unconverted), the mixture was partitioned between ethyl acetate and water. The aqueous phase was extracted with ethyl acetate and the combined organic layers were washed with brine and dried over magnesium sulfate. Chromatographic purification (the title product eluted with 95–100% ethyl acetate) yielded alcohol ed 140-3 as an off-white solid (11 mg, 8%). LC– 500 MS  $R_t$  4.3 min (method 1);  $m/z$  [MH<sup>+</sup>] 253. <sup>1</sup>H NMR  $\delta$  11.59 (1H, br s, indole-H1), 9.88 (1H, br s, OH), 8.52 (1H, d,  $J = 1.8$  Hz, ArH), 8.42 (1H, dd, J = 4.6, 1.4 Hz, ArH), 7.70 (1H, d, J = 7.6 Hz, ArH), 7.57 (1H, d,  $J = 1.6$  Hz, ArH), 7.32 (1H, dd,  $J = 7.3$ , 4.6 Hz, ArH), 7.03 (1H, t,  $J = 8.0$  Hz, ArH), 6.82 (1H, d,  $J = 8.3$  Hz, ArH), 6.37 (1H, d, J = 7.6 Hz, ArH), 4.32 (2H, s, CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$  189.5 (C=O), 153.1 (ArC), 151.1 (ArC), 148.3 (ArC), 140.5 (ArC), 137.7 (ArC), 134.0 (ArC), 131.8 (ArC), 127.6 (ArC), 123.9 (ArC), 118.7 (ArC), 108.7 (ArC), 104.1 (ArC), 103.9 (ArC), 41.9 (CH<sub>2</sub>).

5.1.3.6. 1-(5-Fluoro-1H-indol-2-yl)-2-pyridin-3-yl-ethanone (7f). Prepared from commercial 5-fluoro-1H-indole-2-carboxylic acid (186 mg, 1.02 mmol) and 3-picoline via corresponding acyl chloride 3f (method A). Yellowish solid (62 mg, 24%), mp  $194-197$  °C. LC–MS  $R_t$  5.3 min (method 1);  $m/z$  [MH<sup>+</sup>] 255. <sup>1</sup>H NMR  $\delta$  11.88 (1H,  $\frac{1}{2}$  or  $\frac{1}{2}$ , indole-H1), 8.51 (1H,  $\frac{1}{2}$  br s, ArH), 8.43 (1H, d, J = 4.8 Hz, ArH), 7.70 (1H, d, J = 7.8 Hz, ArH), 7.53 (1H, br s, ArH), 7.46 (1H, dd,  ${}^{3}$ <sub>JHF</sub> = 9.6 Hz,  ${}^{4}$ J<sub>HH</sub> = 2.1 Hz, ArH), 7.42 (1H, dd,  ${}^{3}$ J<sub>HH</sub> = 9.1 Hz, 4.4.<br><sup>4</sup>L = 4.7 Hz, ArH), 7.33 (1H, dd, I = 7.8, 4.8 Hz, ArH), 7.14 (1H, dt,  $^{4}J_{HF}$  = 4.7 Hz, ArH), 7.33 (1H, dd, J = 7.8, 4.8 Hz, ArH), 7.14 (1H, dt,  $J = 9.2$ , 2.3 Hz, ArH), 4.36 (2H, s, CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$  190.2 (C=O), 157.8 (d,  $J_{CF}$  = 233 Hz, indole-C5), 151.1 (ArC), 148.4 (ArC), 137.8 520 (ArC), 136.8 (ArC), 135.2 (ArC), 131.4 (ArC), 127.4 (d,  $J_{CF}$  = 10 Hz, ArC), 124.0 (ArC), 115.2 (d,  $J_{CF}$  = 27 Hz, ArC), 114.7 (d,  $J_{CF}$  = 10 Hz, ArC), 110.3 (d,  $J_{CF}$  = 5 Hz, indole-C7a), 107.1 (d,  $J_{CF}$  = 23 Hz, ArC), 42.1 (CH<sub>2</sub>). <sup>19</sup>F NMR  $\delta$  -122.4. Anal. Calcd for C<sub>15</sub>H<sub>11</sub>FN<sub>2</sub>O: C, 70.86; H, 4.36; N, 11.02. Found: C, 70.61; H, 4.36; N, 10.87.

5.1.3.7. 1-(5-Chloro-1H-indol-2-yl)-2-pyridin-3-yl-ethanone (7g). Prepared from commercial 5-chloro-1H-indole-2-carboxylic acid (355 mg, 1.78 mmol) and 3-picoline via corresponding acyl chloride 3g (method A). Yellow solid (137 mg, 29%), mp 206– 208 °C (dec). LC–MS  $R_t$  5.8 min (method 1);  $m/z$  [MH<sup>+</sup>] 271. <sup>1</sup>H NMR  $\delta$  11.97 (1H, br s, indole-H1), 8.51 (1H, br s, ArH), 8.43 (1H, d,  $J = 4.8$  Hz, ArH), 7.77 (1H, br s, ArH), 7.70 (1H, d,  $J = 7.8$  Hz, ArH), 7.52 (1H,  $\frac{1}{2}$  pr s, ArH), 7.43 (1H, d, J = 8.7 Hz, ArH), 7.33 (1H, dd,  $J = 7.8$ , 4.8 Hz, ArH), 7.26 (1H, dd,  $J = 8.7$ , 1.8 Hz, ArH), 4.38 (2H, s, CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$  190.3 (C=O), 151.1 (ArC), 148.4 (ArC), 137.9 (ArC), 136.8 (ArC), 136.6 (ArC), 131.3 (ArC), 128.4 (ArC), 126.3 (ArC), 125.3 (ArC), 124.0 (ArC), 122.2 (ArC), 115.0 (ArC), 109.8 (ArC), 42.1 (CH<sub>2</sub>). Anal. Calcd for C<sub>15</sub>H<sub>11</sub>ClN<sub>2</sub>O: C, 66.55; H, 4.10; N, 10.35. Found: C, 67.48; H, 4.11; N, 9.99.

5.1.3.8. 1-(5-Bromo-1H-indol-2-yl)-2-pyridin-3-yl-ethanone (7h). Prepared from commercial 5-bromo-1H-indole-2-carboxylic acid (260 mg, 1.08 mmol) and 3-picoline via corresponding acyl chloride 3h (method A). White solid (79 mg, 23%), decomposes without melting at  $224-226$  °C. LC–MS  $R_{t_1}$  6.0 min (method 1), 13.3 min (method 2);  $m/z$  [MH<sup>+</sup>] 315; 317. <sup>1</sup>H NMR  $\delta$  11.99 (1H, br s, indole-H1), 8.51 (1H, d,  $J = 1.8$  Hz, ArH), 8.43 (1H, dd,  $J = 4.8$ , 1.2 Hz, ArH), 7.92 (1H, br s, ArH), 7.70 (1H, d,  $J = 7.8$  Hz, ArH), 7.52 (1H, d,  $J = 1.6$  Hz, ArH), 7.37 (2H, overlapped, 2 ArH), 7.33 (1H, dd,  $J = 7.8$ , 4.8 Hz, ArH), 4.38 (2H, s, CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$  190.3 (C=O), 151.2 (ArC), 148.4 (ArC), 137.9 (ArC), 137.0 (ArC), 136.4 (ArC), 131.3 (ArC), 129.1 (ArC), 128.7 (ArC), 125.3 (ArC), 124.0 (ArC), 115.4 (ArC), 113.2 (ArC), 109.6 (ArC), 42.1 (CH<sub>2</sub>). Anal. Calcd for  $C_{15}H_{11}BrN_2O$ : C, 57.16; H, 3.52; N, 8.89. Found: C, 57.00; H, 3.31; N, 8.40.

5.1.3.9. 1-(5-Methyl-1H-indol-2-yl)-2-pyridin-3-yl-ethanone (7i). Prepared from commercial 5-methyl-1H-indole-2-carboxylic acid (482 mg, 2.75 mmol) and 3-picoline via corresponding acyl chloride 3i (method A). Yellowish solid (114 mg, 17%), mp 181– 182 °C (dec). LC–MS  $R_t$  5.7 min (method 1);  $m/z$  [MH<sup>+</sup>] 251. <sup>1</sup>H NMR  $\delta$  11.64 (1H, br s, indole-H1), 8.52 (1H, br s, ArH), 8.42 (1H, d,  $J = 3.9$  Hz, ArH), 7.70 (1H, d,  $J = 7.8$  Hz, ArH), 7.46 (2H, br s, 560 2ArH), 7.34-7.29 (2H, m, 2ArH), 7.09 (1H, d, J = 8.5 Hz, ArH), 4.33 (2H, s, CH<sub>2</sub>), 2.34 (3H, s, CH<sub>3</sub>). <sup>13</sup>C NMR  $\delta$  190.0 (C=O), 151.1 (ArC), 148.3 (ArC), 137.7 (ArC), 137.1 (ArC), 135.4 (ArC), 131.7 (ArC), 129.6 (ArC), 128.2 (ArC), 127.7 (ArC), 123.9 (ArC), 122.3 (ArC), 113.0 (ArC), 110.1 (ArC), 42.0 (CH<sub>2</sub>), 21.6 (CH<sub>3</sub>). Anal. Calcd for  $C_{16}H_{14}N_2O$ : C, 76.78; H, 5.64; N, 11.19. Found: C, 77.04; H, 5.52; N, 10.90.

5.1.3.10. 1-(5-Methoxy-1H-indol-2-yl)-2-pyridin-3-yl-ethanone (7j) and 1-(7-methoxy-1H-indol-2-yl)-2-pyridin-3-yl-ethanone  $(7q)^{43}$ . Prepared from a ~1:1 mixture (method B; 208 mg, 0.95 mmol) of 5-methoxy-1H-indole-2-carboxylic acid ethyl ester (3j) and 7-methoxy-1H-indole-2-carboxylic acid ethyl ester  $(3q)$ and 3-picoline. The two final products were separated by chromatography. 7j: (Sundberg 1978) eluted with  $\sim$ 80% ethyl acetate. Yellowish solid (88 mg, 35%). LC–MS  $R_t$  5.1 min (method 1), 8.4 min (method 2);  $m/z$  [MH<sup>+</sup>] 267. <sup>1</sup>H NMR  $\delta$  11.64 (1H, br s, indole-H1), 8.52 (1H, d,  $J = 2.0$  Hz, ArH), 8.43 (1H, dd,  $J = 4.7$ , 1.5 Hz, ArH), 7.70 (1H, dt,  $J = 7.8$ , 1.8 Hz, ArH), 7.44 (1H, d,  $J = 1.8$  Hz, ArH), 7.34-7.29 (2H, overlapped, 2 ArH), 7.11 (1H, d, J = 2.4 Hz, ArH),  $\overline{6.92}$  (1H, dd, J = 8.9, 2.4 Hz, ArH), 4.33 (2H, s, CH<sub>2</sub>), 3.74 580 (3H, s, OCH<sub>3</sub>). <sup>13</sup>C NMR  $\delta$  189.8 (C=O), 154.6 (ArC), 147.4 (ArC), 135.6 (ArC), 134.0 (ArC), 132.2 (ArC), 132.1 (2 ArC), 129.2 (ArC), 127.7 (ArC), 118.0 (ArC), 114.3 (ArC), 110.1 (ArC), 102.8 (ArC), 55.8 (OCH<sub>3</sub>), 42.0 (CH<sub>2</sub>). **7q**: eluted with  $\sim$ 70% ethyl acetate. White solid (111 mg, 44%), mp 171-175 °C (dec). LC-MS  $R_t$  5.3 min

8 E. Dolušić et al. / Bioorg. Med. Chem. xxx (2010) xxx–xxx

(method 1), 8.7 min (method 2);  $m/z$  [MH<sup>+</sup>] 267. <sup>1</sup>H NMR  $\delta$  11.72 (1H,  $\frac{1}{2}$  pr s, indole-H1), 8.51 (1H, d, J = 1.6 Hz, ArH), 8.42 (1H, d,  $J = 3.4$  Hz, ArH), 7.69 (1H, d,  $J = 8.0$  Hz, ArH), 7.50 (1H, d,  $J = 1.4$  Hz, ArH), 7.32 (1H, dd,  $J = 7.7$ , 4.7 Hz, ArH), 7.25 (1H, d,  $J = 8.0$  Hz, ArH), 6.99 (1H, t,  $J = 7.9$  Hz, ArH), 6.79 (1H, d,  $J = 7.6$  Hz, ArH), 4.35 (2H, s, CH<sub>2</sub>), 3.87 (3H, s, OCH<sub>3</sub>). <sup>13</sup>C NMR  $\delta$ 189.7 (C=O), 151.1 (ArC), 148.3 (ArC), 147.4 (ArC), 137.8 (ArC), 135.6 (ArC), 131.6 (ArC), 129.5 (ArC), 128.8 (ArC), 123.9 (ArC), 121.6 (ArC), 115.3 (ArC), 111.0 (ArC), 105.6 (ArC), 55.9 (OCH3), 42.2 (CH<sub>2</sub>). Anal. Calcd for C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>2</sub>: C, 72.16; H, 5.30; N, 10.52. Found: C, 72.90; H, 5.14; N, 10.20.

5.1.3.11. 1-(5-Nitro-1H-indol-2-yl)-2-pyridin-3-yl-ethanone (7k). Prepared from commercial 5-nitro-1H-indole-2-carboxylic acid ethyl ester 3k (249 mg, 1.01 mmol) and 3-picoline. Reaction time: 600 48 h. Bright yellow solid (42 mg, 15%). LC–MS  $R_t$  5.3 min (method 1);  $m/z$  [MH<sup>+</sup>] 282. <sup>1</sup>H NMR  $\delta$  12.46 (1H, br s, indole-H1), 8.75 (1H,  $\frac{1}{2}$  pr s, ArH), 8.52 (1H,  $\frac{1}{2}$  pr s, ArH), 8.44 (1H, d, J = 4.6 Hz, ArH), 8.12 (1H, d,  $J = 9.2$  Hz, ArH), 7.82 (1H, s, ArH), 7.71 (1H, d,  $J = 7.8$  Hz, ArH), 7.57 (1H, d,  $J = 9.2$  Hz, ArH), 7.34 (1H, dd,  $J = 8.0$ , 5.0 Hz, ArH),  $4.44$  (2H, s, CH<sub>2</sub>).

5.1.3.12. 1-(5-Phenyl-1H-indol-2-yl)-2-pyridin-3-ylethanone  $(71)^{46}$ . Prepared according to a literature procedure. A suspension of tetrakis(triphenylphosphine)palladium  $(0)$  (7 mg, 6  $\mu$ mol) in a mixture of 1 M potassium carbonate (0.8 mL), ethanol (0.8 mL) and toluene <sup>610</sup> (0.8 mL) was added to a stirred suspension of bromoindole 7h (38 mg, 0.12 mmol) and phenylboronic acid (33 mg, 0.26 mmol) in a 1:1 mixture of toluene and ethanol (0.8 mL). The (initially ochre colored) suspension was heated at reflux temperature for 20 h, allowed to cool down to room temperature and diluted with ethyl acetate. Evaporation with silica and chromatographic purification afforded the title product as a bright yellow solid (12 mg, 32%). LC–MS  $R_t$  6.5 min (method 1), 10.2 min (method 2);  $m/z$ [MH<sup>+</sup>] 313. <sup>1</sup>H NMR  $\delta$  11.85 (1H, br s, indole-H1), 8.54 (1H, d,  $J = 1.8$  Hz, ArH), 8.44 (1H, dd,  $J = 4.\overline{8}$ , 1.6 Hz, ArH), 7.96 (1H, br s, 620 ArH), 7.72 (1H, dt, J = 7.8, 1.8 Hz, ArH),  $7.67-7.49$  (5H, overlapped, 5 ArH), 7.43 (2H, t,  $J = 7.7$  Hz, 2 ArH), 7.33 (1H, dd,  $J = 8.0$ , 2.7 Hz, ArH), 7.29 (1H, t, J = 7.2 Hz, ArH), 4.39 (2H, s, CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$ 190.1 (C=O), 151.1 (ArC), 148.3 (ArC), 141.6 (ArC), 138.1 (ArC), 137.8 (ArC), 136.1 (ArC), 134.6 (ArC), 132.1 (ArC), 132.0 (ArC), 129.42 (2 ArC), 129.36 (ArC), 127.3 (2 ArC), 125.9 (ArC), 124.0 (ArC), 121.0 (ArC), 113.8 (ArC), 111.0 (ArC), 42.1 (CH<sub>2</sub>). Anal. Calcd for  $C_{21}H_{16}N_2O$ : C, 80.75; H, 5.16; N, 8.97. Found: C, 80.01; H, 5.23; N, 8.49.

5.1.3.13. 2-Pyridin-3-yl-1-(5-trifluoromethoxy-1H-indol-2-yl)-  $630$  ethanone (7m). Prepared from commercial 5-trifluoromethoxy-1H-indole-2-carboxylic acid (256 mg, 1.01 mmol) and 3-picoline via corresponding acyl chloride 3m (method A). Yellowish solid (90 mg, 28%), mp 184–186 °C. LC–MS  $R_t$  6.2 min (method 1); m/z [MH<sup>+</sup>] 321. <sup>1</sup>H NMR  $\delta$  12.04 (1H, br s, indole-H1), 8.52 (1H, d,  $J = 1.6$  Hz, ArH), 8.43 (1H, dd,  $J = 4.6$ , 1.4 Hz, ArH), 7.71–7.69 (2H, overlapped, 2ArH), 7.60 (1H,  $\frac{1}{1}$  br s, ArH), 7.50 (1H, d, J = 8.9 Hz, ArH), 7.33 (1H, dd, J = 7.8, 4.8 Hz, ArH), 7.24 (1H, d, J = 8.9 Hz, ArH), 4.38 (2H, s, CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$  190.3 (C=O), 151.1 (ArC), 148.4 (ArC), 142.9 (ArC), 137.8 (ArC), 137.1 (ArC), 136.8 (ArC), 640 131.3 (ArC), 127.3 (ArC), 124.0 (ArC), 120.9 (d,  $J_{CF} = 254$  Hz, OCF<sub>3</sub>), 120.1 (ArC), 115.2 (ArC), 114.8 (ArC), 110.6 (ArC), 42.1 (CH<sub>2</sub>). <sup>19</sup>F NMR  $\delta$  -56.9. Anal. Calcd for C<sub>16</sub>H<sub>11</sub>F<sub>3</sub>N<sub>2</sub>O<sub>2</sub>: C, 60.00; H, 3.46; N, 8.75. Found: C, 59.95; H, 3.43; N, 8.45.

5.1.3.14. 1-(5-Benzyloxy-1H-indol-2-yl)-2-pyridin-3-yl-ethanone (7n). Prepared from commercial 5-benzyloxy-1H-indole-2-carboxylic acid ethyl ester 3n (499 mg, 1.69 mmol) and 3-picoline. Off-white solid (166 mg, 29%), mp 183–184 °C. LC–MS  $R_t$  6.5 min

Please cite this article in press as: Dolušić, E.; et al. Bioorg. Med. Chem. (2010), doi[:10.1016/j.bmc.2010.12.032](http://dx.doi.org/10.1016/j.bmc.2010.12.032)

(method 1), 14.8 min (method 2);  $m/z$  [MH<sup>+</sup>] 343. <sup>1</sup>H NMR  $\delta$ 11.65 (1H, br s, indole-H1), 8.52 (1H, br s, ArH), 8.42 (1H, d,  $J = 4.8$  Hz, ArH), 7.70 (1H, d, J = 7.8 Hz, ArH), 7.45–7.43 (3H, over- 650 lapped, 3ArH), 7.38–7.27 (5H, overlapped, 5ArH), 7.21 (1H, d,  $J = 1.8$  Hz, ArH), 7.00 (1H, dd,  $J = 8.9$ , 2.3 Hz, ArH), 5.08 (2H, s, benzyl CH<sub>2</sub>), 4.32 (2H, s, C(O)CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$  189.9 (C=O), 153.5 (ArC), 151.1 (ArC), 148.3 (ArC), 137.9 (ArC), 137.7 (ArC), 135.8 (ArC), 134.1 (ArC), 131.7 (ArC), 128.9 (2C, 2 phenyl C), 128.3 (ArC), 128.2 (2C, 2 phenyl C), 127.7 (ArC), 124.0 (ArC), 118.5 (ArC), 114.3 (ArC), 110.2 (ArC), 104.4 (ArC), 70.1 (benzyl CH<sub>2</sub>), 42.0 (C(O)CH<sub>2</sub>). Anal. Calcd for C<sub>22</sub>H<sub>18</sub>N<sub>2</sub>O<sub>2</sub>: C, 77.17; H, 5.30; N, 8.18. Found: C, 76.95; H, 5.01; N, 8.10.

5.1.3.15. 1-(5-Hydroxy-1H-indol-2-yl)-2-pyridin-3-yl-ethanone <sup>660</sup> (7o) and 2-(1-hydroxy-2-pyridin-3-ylethyl)-1H-indol-5-ol (17). Ammonium formate (485 mg, 7.50 mmol) and Pd black (surface area  $40-60$  m<sup>2</sup>/g, 110 mg, 1.03 mmol) were added to a stirred suspension of benzyl ether **7n** (166 mg, 0.13 mmol) in dry methanol (55 mL). The blue-greenish suspension was stirred at room temperature for 1 h and filtered through Celite 545. The filter was extensively washed with methanol and the combined filtrate and washings were concentrated under reduced pressure. Chromatographic purification yielded hydroxyindole 7o (eluted with 95% ethyl acetate; 38 mg, 31%) and byproduct  $17$  (eluted with 100% ethyl acetate; 69 mg, 56%). Shortening the reaction time to 30 min afforded 7o as the only product in a 58% yield. Analytical data: 7o, yellow solid which decomposes without melting above 200 °C. **LC-MS**  $R_t$  4.1 min (method 1);  $m/z$  [MH<sup>+</sup>] 253. <sup>1</sup>H NMR  $\delta$ 11.49 (1H, br s, indole-H1), 8.97 (1H, s, OH), 8.51 (1H, br s, ArH), 8.42 (1H, d, J = 3.6 Hz, ArH), 7.69 (1H, dt,  $^{3}$ J<sub>HH</sub> = 7.8 Hz,  $^{4}$ L<sub>H</sub> = 7.8 Hz,  $^{4}$ L<sub>H</sub> = 7.8 Hz  $^{4}$ H<sub>H</sub> = 1.8 Hz, ArH), 7.36 (1H, d, J = 1.8 Hz, ArH), 7.32 (1H, dd,  $J = 7.8$ , 4.8 Hz, ArH), 7.22 (1H, d,  $J = 8.9$  Hz, ArH), 6.93 (1H, d,  $J = 2.3$  Hz, ArH), 6.81 (1H, dd,  $J = 8.7$ , 2.3 Hz, ArH), 4.29 (2H, s, CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$  189.8 (C=O), 152.0 (ArC), 151.1 (ArC), 148.3 (ArC), 137.7 (ArC), 135.6 (ArC), 133.5 (ArC), 131.7 (ArC), 128.1 (ArC), 123.9 (ArC), 118.0 (ArC), 113.9 (ArC), 109.6 (ArC), 105.3 (ArC), 41.9 (CH<sub>2</sub>). Compound 17, yellowish solid, mp  $188-190$  °C. LC–MS  $R_t$  3.1 min (method 1);  $m/z$  [MH<sup>+</sup>] 255. <sup>1</sup>H NMR  $\delta$  10.68 (1H, br s, indole-H1), 8.50 (1H, br s, ArH), 8.33–8.32 (2H, overlapped, ArH, OH on indole), 7.55 (1H, dd, <sup>3</sup>J<sub>HH</sub> = 7.8 Hz, 4<sup>2</sup>H<sub>H</sub> = 7.8 Hz, 4<sup>2</sup>H<sub>H</sub> = 7.8 Hz  $^{4}$ J<sub>HH</sub> = 1.8 Hz, ArH), 7.22 (1H, dd, J = 7.8, 4.8 Hz, ArH), 7.08 (1H, d,  $J = 8.6$  Hz, ArH), 6.70 (1H, d,  $J = 2.0$  Hz, ArH), 6.50 (1H, dd,  ${}^{3}J_{\text{HH}}$  = 8.6 Hz,  ${}^{4}J_{\text{HH}}$  = 2.3 Hz, ArH), 6.01 (1H, d, J = 1.4 Hz, ArH), 5.44 (1H, d, J = 5.3 Hz, CHOH), 4.81 (1H, m, CHOH), 3.07 (1H, dd,  $^{2}J_{HH}$  = 13.6 Hz,  $^{3}J_{HH}$  = 5.4 Hz, CH<sub>2</sub>), 2.97 (1H, dd,  $^{2}J_{HH}$  = 13.6 Hz,  $^{3}L_{H}$  = 13.6 Hz,  $^{13}C$  NMR  $^{3}$  151.0 (ArC) 150.9 (ArC), 147.7  $^{3}J_{\text{HH}}$  = 8.0 Hz, CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$  151.0 (ArC), 150.9 (ArC), 147.7 (ArC), 143.4 (ArC), 137.3 (ArC), 134.9 (ArC), 131.0 (ArC), 128.9 (ArC), 123.6 (ArC), 111.9 (ArC), 111.2 (ArC), 104.2 (ArC), 97.7  $(ArC)$ , 68.4 (CHOH), 41.0 (CH<sub>2</sub>).

5.1.3.16. 1-(6-Fluoro-1H-indol-2-yl)-2-pyridin-3-yl-ethanone (7p). Prepared from 6-fluoro-1H-indole-2-carboxylic acid ethyl ester 3p (method B; 80 mg, 0.39 mmol) and 3-picoline. Bright yellow solid (59 mg, 60%), mp 192–193 °C (dec). LC–MS  $R_t$  5.4 min (method 1);  $m/z$  [MH<sup>+</sup>] 255.<sup>1</sup>H NMR  $\delta$  11.86 (1H, br s, indole-<br>700 H1), 8.52 (1H, d,  $J = 1.7$  Hz, ArH), 8.43 (1H, dd,  $J = 4.9$ , 1.7 Hz, ArH), 7.74 (1H, dd, overlapped, ArH), 7.71 (1H, d, overlapped, ArH), 7.60 (1H, d,  $J = 1.4$  Hz, ArH), 7.33 (1H, dd,  $J = 7.7$ , 4.9 Hz, ArH), 7.11 (1H, dd,  ${}^{3}$  $_{\text{JHF}}$  = 9.9 Hz,  ${}^{4}$  $_{\text{JHH}}$  = 1.9 Hz, ArH), 6.95 (1H, dt,  ${}^{3}$  $_{\text{J}}$  = 0.3 Hz,  ${}^{4}$  $_{\text{J}}$  = 1.9 Hz,  ${}^{4}$  $_{\text{A}}$ rH), 4.34 (2H, s, CH, ),  ${}^{13}$ C, NMR  $_{\text{S}}$  1.89.8  $J = 9.3$  Hz,  $^{4}J = 1.9$  Hz, ArH), 4.34 (2H, s, CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$  189.8  $(C=0)$ , 161.6 (d,  $J<sub>CF</sub>$  = 219 Hz, indole-C6), 151.1 (ArC), 148.4 (ArC), 138.6 (d,  $J_{CF}$  = 13 Hz, ArC), 137.8 (ArC), 136.3 (d,  $J_{CF}$  = 3 Hz, indole-C3a), 131.5 (ArC), 125.1 (d,  $J_{CF}$  = 11 Hz, ArC), 124.4 (ArC), 124.0 (ArC), 110.9 (ArC), 110.4 (d,  $J_{CF}$  = 26 Hz, ArC), 98.6 (d,  $J_{CF}$  = 26 Hz, ArC), 41.9 (CH<sub>2</sub>). <sup>19</sup>F NMR  $\delta$  -115.0. Anal. Calcd for C<sub>15</sub>H<sub>11</sub>FN<sub>2</sub>O: 710 C, 70.86; H, 4.36; N, 11.02. Found: C, 71.12; H, 4.72; N, 10.81.

E. Dolušić et al. / Bioorg. Med. Chem. xxx (2010) xxx–xxx  $\qquad \qquad$  9

5.1.3.17. 1-(4-Benzyloxy-5-methoxy-1H-indol-2-yl)-2-pyridin-3 **vl-ethanone (7r).** Prepared from 4-benzyloxy-5-methoxy-1H-indole-2-carboxylic acid ethyl ester 3r (method B; 282 mg, 0.87 mmol) and 3-picoline. Yellow solid (84 mg, 26%), mp 137– 139 °C. LC–MS  $R_t$  6.3 min (method 1), 15.7 min (method 2);  $m/z$ [MH<sup>+</sup>] 373. <sup>1</sup>H NMR δ 11.65 (1H, br s, indole-H1), 8.51 (1H, br s, ArH), 8.43 (1H, d,  $J = 4.1$  Hz, ArH), 7.69 (1H, d,  $J = 7.8$  Hz, ArH), 7.54–7.50 (3H, overlapped, 3ArH), 7.38–7.30 (4H, overlapped,  $720$  4ArH), 7.14 (1H, d, J = 8.8 Hz, ArH), 7.07 (1H, d, J = 8.8 Hz, ArH), 5.22 (2H, s, benzyl CH<sub>2</sub>), 4.36 (2H, s, C(O)CH<sub>2</sub>), 3.79 (3H s, OCH<sub>3</sub>). <sup>13</sup>C NMR  $\delta$  190.0 (C=O), 151.1 (ArC), 148.3 (ArC), 144.8 (ArC), 141.5 (ArC), 138.4 (ArC), 137.8 (ArC), 135.6 (ArC), 131.6 (ArC), 128.8 (2 ArC), 128.6 (2 ArC), 128.4 (ArC), 124.0 (ArC), 122.9 (ArC), 117.4 (ArC), 108.4 (ArC), 107.8 (ArC), 74.6 (benzyl CH<sub>2</sub>), 58.5 (OCH<sub>3</sub>), 41.9 (C(O)CH<sub>2</sub>). Anal. Calcd for C<sub>23</sub>H<sub>20</sub>N<sub>2</sub>O<sub>3</sub>: C, 74.18; H, 5.41; N, 7.52. Found: C, 74.98; H, 5.29; N, 7.42.

5.1.3.18. 1-(4-Hydroxy-5-methoxy-1H-indol-2-yl)-2-pyridin-3 yl-ethanone (7s). Benzyl ester 7r (59 mg, 0.16 mmol) was dis-<sup>730</sup> solved in absolute ethanol (15 mL) to give a clear yellow solution to which 3% Pd on activated carbon (60 mg,  $17 \mu$ mol) was added. The flask was evacuated and back-filled with hydrogen gas (repeated three times). The suspension was stirred at room temperature for 45 min (until TLC indicated reaction completion) and filtered through Celite 545. The filter was extensively washed with ethyl acetate and the combined filtrate and washings were concentrated under reduced pressure. Chromatographic purification (the title product eluted with 80–90% ethyl acetate) afforded the title hydroxyindole 7s as a light brown half-solid in a 53% (24 mg) yield. 740 LC-MS  $R_t$  4.3 min (method 1);  $m/z$  [MH<sup>+</sup>] 283. <sup>1</sup>H NMR  $\delta$  11.47 (1H, br s, indole-H1), 9.26 (1H, s, OH), 8.52 (1H, br s, ArH), 8.42 (1H, d,

 $J = 3.9$  Hz, ArH), 7.70 (1H, d,  $J = 7.6$  Hz, ArH), 7.54 (1H, d,  $J = 1.4$  Hz, ArH), 7.33 (1H, dd, J = 7.8, 4.8 Hz, ArH), 7.04 (1H, d, J = 8.7 Hz, ArH), 6.78 (1H, d, J = 8.7 Hz, ArH), 4.31 (2H, s, CH<sub>2</sub>), 3.72 (3H s, OCH<sub>3</sub>). <sup>13</sup>C NMR δ 189.6 (C=O), 151.1 (ArC), 148.3 (ArC), 141.2 (ArC), 139.5 (ArC), 137.7 (ArC), 136.0 (ArC), 134.9 (ArC), 131.8 (ArC), 124.0 (ArC), 119.8 (ArC), 117.8 (ArC), 108.4 (ArC), 103.4 (ArC), 59.0  $(OCH<sub>3</sub>)$ , 41.9 (CH<sub>2</sub>).

5.1.3.19. 1-(4,6-Dichloro-1H-indol-2-yl)-2-pyridin-3-yl-ethanone (7t). Prepared from commercial 4,6-dichloro-1H-indole-2-carbonyl chloride 3t (647 mg, 2.47 mmol) and 3-picoline. Off-white solid (198 mg, 26%), mp 223–225 °C. LC–MS  $R_t$  6.5 min (method 1);  $m/z$ [MH<sup>+</sup>] 305. <sup>1</sup>H NMR  $\delta$  12.26 (1H, br s, indole-H1), 8.52 (1H, d,  $J = 1.7$  Hz, ArH), 8.44 (1H, dd,  $J = 4.\overline{8}$ , 1.7 Hz, ArH), 7.70 (1H, dt,  $J = 7.8$ , 1.7 Hz, ArH), 7.62 (1H, br s, ArH), 7.41 (1H, dd,  $J = 1.6$ , 0.9 Hz, ArH), 7.33 (1H, dd, J = 7.8, 4.8 Hz, ArH), 7.27 (1H, d, J = 1.8 Hz, ArH), 4.44 (2H, s, CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$  190.3 (C=0), 151.2 (ArC), 148.4 (ArC), 138.7 (ArC), 137.9 (ArC), 136.8 (ArC), 131.2 (ArC), 130.4 (ArC), 128.2 (ArC), 125.2 (ArC), 124.0 (ArC), 120.7 (ArC), 112.0 (ArC), 108.1  $760$  (ArC), 42.0 (CH<sub>2</sub>). Anal. Calcd for C<sub>15</sub>H<sub>10</sub>Cl<sub>2</sub>N<sub>2</sub>O: C, 59.04; H, 3.30; N, 9.18. Found: C, 59.79; H, 3.26; N, 9.21.

5.1.3.20. 1-(4,6-Dimethoxy-1H-indol-2-yl)-2-pyridin-3-yl-ethanone (7u). Prepared from commercial 4,6-dimethoxy-1H-indole-2-carboxylic acid methyl ester 3u (241 mg, 1.01 mmol) and 3-picoline. Off-white solid (240 mg, 80%), mp 195-197 °C. LC-MS  $R_t$ 5.3 min (method 1);  $m/z$  [MH<sup>+</sup>] 297. <sup>1</sup>H NMR  $\delta$  11.60 (1H, s, indole-H1), 8.52 (1H, d,  $J = 1.6$  Hz, ArH), 8.41 (1H, dd,  $J = 4.8$ , 1.6 Hz, ArH), 7.69 (1H, dt, J = 7.8, 0.9 Hz, ArH), 7.51 (1H, d, J = 1.8 Hz, ArH), 7.31 (1H, dd,  $J = 7.8$ , 4.8 Hz, ArH), 6.40 (1H, d,  $J = 1.5$  Hz, 770 ArH), 6.17 (1H, d, J = 1.5 Hz, ArH), 4.25 (2H, s, CH<sub>2</sub>), 3.84 (3H, s, OCH<sub>3</sub>), 3.73 (3H, s, OCH<sub>3</sub>). <sup>13</sup>C NMR  $\delta$  188.6 (C=O), 160.8 (ArC), 155.4 (ArC), 151.0 (ArC), 148.2 (ArC), 140.6 (ArC), 137.6 (ArC), 133.5 (ArC), 132.0 (ArC), 123.9 (ArC), 113.8 (ArC), 109.2 (ArC), 93.1 (ArC), 87.0 (ArC), 55.83 (OCH<sub>3</sub>), 55.76 (OCH<sub>3</sub>), 41.5 (CH<sub>2</sub>). Anal.

Calcd for  $C_{17}H_{16}N_2O_3$ : C, 68.91; H, 5.44; N, 9.45. Found: C, 69.65; H, 5.17; N, 9.40.

5.1.3.21. 1-(1H-Indol-2-yl)-2-pyridin-3-yl-propan-1-one (7v). Prepared from commercial 1H-indole-2-carboxylic acid ethyl ester 3a (983 mg, 5.04 mmol) and 3-ethylpyridine. Yellow half-solid (220 mg, 17%). LC–MS  $R_t$  5.7 min (method 1), 7.3 min (method 780 2);  $m/z$  [MH<sup>+</sup>] 251. <sup>1</sup>H NMR  $\delta$  11.74 (1H, br s, indole-H1), 8.65 (1H, d, J = 1.7 Hz, ArH), 8.39 (1H, dd, J = 4.8, 1.7 Hz, ArH), 7.75  $(1H, dt, J = 7.8, 1.7 Hz, ArH), 7.64 (1H, d, J = 8.2 Hz, ArH), 7.53$  $(1H, d, J = 1.4 Hz, ArH), 7.37 (1H, d, J = 8.5 Hz, ArH), 7.31 (1H, dd,$  $J = 7.9$ , 4.7 Hz, ArH), 7.23 (1H, t,  $J = 7.7$  Hz, ArH), 7.03 (1H, t,  $J = 7.4$  Hz, ArH), 4.89 (1H, q,  $J = 7.0$  Hz, CHCH<sub>3</sub>), 1.46 (3H, d,  $J = 7.0$  Hz, CH<sub>3</sub>). <sup>13</sup>C NMR  $\delta$  193.0 (C=O), 149.7 (ArC), 148.6 (ArC), 138.6 (ArC), 137.7 (ArC), 135.5 (ArC), 134.7 (ArC), 127.3 (ArC), 126.3 (ArC), 124.3 (ArC), 123.3 (ArC), 120.9 (ArC), 113.3 (ArC), 110.7 (ArC), 44.5 (CHCH<sub>3</sub>), 19.1 (CH<sub>3</sub>).  $2^{90}$ 

5.1.3.22. 1-(1H-Indol-2-yl)-2-pyridin-2-yl-ethanone (8a). Prepared from commercial 1H-indole-2-carboxylic acid ethyl ester 3a (218 mg, 1.12 mmol) and 2-picoline. Yellow solid (237 mg, 90%), mp  $140-141$  °C. LC–MS  $R_t$  5.5 min (method 1), 9.7 min (method 2);  $m/z$  [MH<sup>+</sup>] 237. <sup>1</sup>H NMR  $\delta$  11.75 (1H, br s, indole-H1), 8.45 (1H, dd,  $J = 4.8$ , 0.9 Hz, ArH), 7.72 (1H, dd,  $J = 7.6$ , 1.8 Hz, ArH), 7.67 (1H, d, J = 8.0 Hz, ArH), 7.44 (1H, d, J = 1.6 Hz, ArH), 7.42–7.38 (2H, overlapped, 2 ArH), 7.26–7.21 (2H, overlapped, 2 ArH), 7.04 (1H, t, J = 7.7 Hz, ArH),  $\overline{4.43}$  (2H, s, CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$  190.0 (C=O), 156.4 (ArC), 149.7 (ArC), 138.4 (ArC), 137.2 (ArC), 135.8 (ArC), 127.4 (ArC), 126.1 (ArC), 124.9 (ArC), 123.3 (ArC), 122.4 (ArC), 120.8 (ArC), 113.3 (ArC), 110.8 (ArC), 48.0 (CH<sub>2</sub>). Anal. Calcd for C<sub>15</sub>H<sub>12</sub>N<sub>2</sub>O: C, 76.25; H, 5.12; N, 11.86. Found: C, 76.09; H, 5.06; N, 11.74.

5.1.3.23. 1-(1H-Indol-2-yl)-2-pyridin-4-yl-ethanone  $(9a)^{47}$ . Prepared from commercial 1H-indole-2-carboxylic acid ethyl ester 3a (220 mg, 1.13 mmol) and 4-picoline. Bright yellow solid (240 mg, 90%), mp  $169-170$  °C. LC–MS  $R_t$  5.0 min (method 1), 8.1 min (method 2);  $\overline{m}/z$  [MH<sup>+</sup>] 237. <sup>1</sup>H NMR  $\delta$  11.79 (1H, br s, indole-H1), 8.48 (2H, d, J = 6.0 Hz, 2 ArH), 7.68 (1H, d, J = 8.0 Hz, ArH),  $810$ 7.55 (1H, d,  $J = 1.1$  Hz, ArH), 7.41 (1H, dd,  $J = 8.3$ , 0.8 Hz, ArH), 7.34  $(2H, d, J = 6.0$  Hz, 2 ArH), 7.26  $(1H, dt, J = 7.7, 1.1$  Hz, ArH), 7.06  $(1H,$ dt, J = 7.5, 0.8 Hz, ArH), 4.35 (2H, s, CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$  189.5 (C=O), 150.0 (2 ArC), 144.8 (ArC), 138.6 (ArC), 135.4 (ArC), 127.4 (ArC), 126.3 (2 ArC), 125.6 (ArC), 123.3 (ArC), 120.9 (ArC), 113.3 (ArC), 110.9 (ArC), 44.2 (CH<sub>2</sub>). Anal. Calcd for C<sub>15</sub>H<sub>12</sub>N<sub>2</sub>O: C, 76.25; H, 5.12; N, 11.86. Found: C, 75.68; H, 5.09; N, 11.73.

5.1.3.24. 1-(1H-Indol-2-yl)-2-pyrazin-2-yl-ethanone (10a). Prepared from commercial 1H-indole-2-carboxylic acid ethyl ester 3a (195 mg, 1.00 mmol) and 2-methylpyrazine. Yellow solid 820 (97 mg, 41%), mp 175–176 °C. LC–MS  $R_t$  5.5 min (method 1), 11.0 min (method 2);  $m/z$  [MH<sup>+</sup>] 238. <sup>1</sup>H NMR  $\delta$  11.80 (1H, br s, indole-H1), 8.69 (1H, d,  $J = 1.6$  Hz, ArH), 8.56 (1H, dd,  $J = 2.5$ , 1.6 Hz, ArH), 8.51 (1H, d,  $J = 2.5$  Hz, ArH), 7.69 (1H, d,  $J = 8.0$  Hz, ArH), 7.51 (1H, d, J = 1.6 Hz, ArH), 7.41 (1H, d, J = 8.3 Hz, ArH), 7.26 (1H, dt,  $J = 7.7$ , 0.9 Hz, ArH), 7.06 (1H, dt,  $J = 7.6$ , 0.7 Hz, ArH), 4.57 (2H, s, CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$  189.3 (C = 0), 152.4 (ArC), 146.4 (ArC), 144.7 (ArC), 143.4 (ArC), 138.5 (ArC), 135.5 (ArC), 127.4 (ArC), 126.3 (ArC), 123.3 (ArC), 120.9 (ArC), 113.3 (ArC), 110.9 (ArC), 45.1 (CH<sub>2</sub>). Anal. Calcd for C<sub>14</sub>H<sub>11</sub>N<sub>3</sub>O: C, 70.87; H, 4.67; N, 830 17.71. Found: C, 70.03; H, 4.63; N, 17.63.

5.1.3.25. 1-(1H-Indol-2-yl)-2-phenylethanone  $(11a)^{36,48}$  and 2-(1H-indol-2-yl)-1,3-diphenylpropan-2-ol (16). An oven-dried flask was purged with argon while hot, then allowed to cool down to room temperature under argon and charged with dry THF (1 mL)

Please cite this article in press as: Dolušić, E.; et al. Bioorg. Med. Chem. (2010), doi:[10.1016/j.bmc.2010.12.032](http://dx.doi.org/10.1016/j.bmc.2010.12.032)

10 **E. Dolušić et al. / Bioorg. Med. Chem. xxx (2010) xxx–xxx** 

and benzyl magnesium chloride, 2.0 M solution in THF (1 mL, 2.0 mmol) to give a clear grayish–yellowish solution. The solution was cooled in an ice/water bath and a solution of 1H-indole-2-carboxylic acid ethyl ester 3a (104 mg, 0.53 mmol) in dry THF (2 mL) was added dropwise over the flask wall. Clear yellowish solution was allowed to attain room temperature overnight. After  $22 h$  of total reaction time (the mixture turned greenish), the reaction was quenched with saturated aqueous ammonium chloride solution (5 mL). Upon stirring for several minutes, the mixture was partitioned between ethyl acetate and water. The aqueous phase was extracted with ethyl acetate and the combined organic layers were washed with brine and dried over MgSO<sub>4</sub>. Chromatographic purification yielded the two products: **11a**: eluted with  $\sim$ 7% ethyl acetate. Yellowish solid (46 mg, 37%), mp 125-133 °C. LC-MS  $R_t$  $_{850}$  6.8 min (method 1), 16.6 min (method 2);  $m/z$  [MH<sup>+</sup>] 236. <sup>1</sup>H NMR  $\delta$  11.73 (1H, br s, indole-H1), 7.67 (1H, d, J = 8.0 Hz, ArH), 7.53 (1H, m, ArH), 7.40 (1H, d, J = 8.2 Hz, ArH), 7.33-7.20 (6H, overlapped, 6 ArH), 7.05 (1H, dt, J = 7.6, 0.9 Hz, ArH), 4.25 (2H, s, CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$  190.8 (C=O), 138.5 (ArC), 136.0 (ArC), 135.6 (ArC), 130.0 (2 ArC), 128.9 (2 ArC), 127.4 (ArC), 127.1 (ArC), 126.1 (ArC), 123.2 (ArC), 120.8 (ArC), 113.3 (ArC), 110.7 (ArC), 45.1 (CH<sub>2</sub>). Anal. Calcd for  $C_{16}H_{13}$ NO: C, 81.68; H, 5.57; N, 5.95. Found: C, 80.67; H, 6.16; N, 5.51. **16**: eluted with  $\sim 9\%$  ethyl acetate. White solid (30 mg, 17%), mp 160–163 °C. LC–MS  $R_t$  7.2 min (method 1), 19.0 min (method  $\frac{\epsilon_{860}}{2}$ ;  $\frac{\hat{m}}{z}$  [MH<sup>+</sup>] 328. <sup>1</sup>H NMR  $\delta$  10.75 (1H, br s, indole-H1), 7.31 (1H, d, J = 7.8 Hz, ArH), 7.27 (1H, d, J = 8.0 Hz, ArH),  $7.09 - 7.03$ (6H, overlapped, 6 ArH), 6.98–6.96 (4H, overlapped, 4 ArH), 6.92 (1H, dt, J = 7.5, 1.0 Hz, ArH), 6.84 (1H, dt, J = 7.3, 0.9 Hz, ArH), 6.04 (1H, d,  $J = 1.4$  Hz, ArH), 5.18 (1H, s, OH), 3.17 (2H, d,  $J_{gem}$  = 13.5 Hz, CH<sub>2</sub>), 3.04 (2H, d,  $J_{gem}$  = 13.5 Hz, CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$ 144.7 (ArC), 138.1 (2C, ArC), 136.2 (ArC), 131.0 (4C, ArC), 128.3 (ArC), 127.9 (4C, ArC), 126.3 (2C, ArC), 120.5 (ArC), 119.9 (ArC), 118.9 (ArC), 111.6 (ArC), 98.9 (ArC), 75.0 (COH), 48.0 (2C, CH2). Anal. Calcd for C<sub>23</sub>H<sub>21</sub>NO: C, 84.37; H, 6.46; N, 4.28. Found: C, <sup>870</sup> 83.84; H, 6.82; N, 3.91.

5.1.3.26. 1-(1-Methyl-1H-indol-2-yl)-2-pyridin-3-yl-ethanone  $(12a)^{43}$ . Prepared from 1-methyl-1H-indole-2-carboxylic acid methyl ester 6a (method C; 544 mg, 2.88 mmol) and 3-picoline. Bright yellow solid (243 mg, 34%), mp  $75-77$  °C. LC–MS  $R_t$ 5.9 min (method 1);  $m/z$  [MH<sup>+</sup>] 251. <sup>1</sup>H NMR  $\delta$  8.51 (1H, s, ArH), 8.43 (1H, d,  $J = 4.6$  Hz, ArH),  $7.74 - 7.69$  (3H, overlapped, 3 ArH), 7.55 (1H, d,  $J = 8.5$  Hz, ArH),  $\overline{7.37} - 7.31$  (2H, overlapped, 2 ArH), 7.13 (1H, t, J = 7.5 Hz, ArH),  $\overline{4.39}$  (2H, s, CH<sub>2</sub>), 3.95 (3H, s, CH<sub>3</sub>). <sup>13</sup>C NMR  $\delta$  191.2 (C=O), 151.1 (ArC), 148.3 (ArC), 140.4 (ArC), <sup>880</sup> 137.8 (ArC), 134.5 (ArC), 131.7 (ArC), 126.5 (ArC), 125.9 (ArC), 123.9 (ArC), 123.4 (ArC), 121.3 (ArC), 113.0 (ArC), 111.5 (ArC), 43.3 (CH<sub>2</sub>), 32.5 (CH<sub>3</sub>). Anal. Calcd for C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O: C, 76.78; H, 5.64; N, 11.19. Found: C, 77.64; H, 5.68; N, 11.24.

5.1.3.27. 1-(1H-Indol-2-yl)-2-pyridin-3-ylethanol  $(18)^{49}$ . Ammonium formate (845 mg, 13.4 mmol) and Pd black (surface area 40–60 m $^2$ /g, 38 mg, 0.36 mmol) were added to a stirred solution of ketone 7a (203 mg, 0.86 mmol) in dry methanol (100 mL). As no significant amount of alcohol 18 was detected by TLC after 24 h, more Pd black (163 mg, 1.54 mmol) was added. Three days <sup>890</sup> later, as TLC indicated complete consumption of starting material, the reaction mixture was filtered through Celite 545 and the filter was extensively washed with methanol. Concentration under reduced pressure, dilution with a small volume of ethyl acetate and precipitation with cyclohexane yielded the title product as an dark orange solid (183 mg, 89%), mp  $144-151$  °C with decomposition. LC–MS  $R_t$  4.4 min (method 1);  $\overline{m}/z$  [MH<sup>+</sup>] 239. <sup>1</sup>H NMR  $\delta$ 11.06 (1H, br  $\overline{s}$ , indole-H1), 8.35–8.32 (2H, overlapped, 2 ArH), 7.56 (1H,  $d, J = 7.8$  Hz, ArH), 7.39 (1H, d, J = 7.8 Hz, ArH), 7.30  $(1H, d, J = 8.0 Hz, ArH), 7.23 (1H, dd, J = 7.8, 4.8 Hz, ArH), 6.98$ 

(1H, t,  $J = 7.4$  Hz, ArH), 6.89 (1H, t,  $J = 7.4$  Hz, ArH), 6.18 (1H, br s, ArH), 5.59 (1H, br s, OH), 4.88 (1H, m, CHOH), 3.10 (1H, dd,  $^{2}J_{HH}$  = 13.7 Hz,  $^{3}J_{HH}$  = 5.5 Hz, CH<sub>2</sub>), 3.01 (1H, dd,  $^{2}J_{HH}$  = 13.7 Hz,  $^{3}J_{HH}$  = 13.7 Hz,  $^{3}J_{HH}$  = 13.7 Hz,  $^{3}$ H<sub>H</sub> = 7.9 Hz, CH<sub>2</sub>). <sup>13</sup>C NMR  $\delta$  151.0 (ArC), 147.7 (ArC), 143.1 (ArC), 137.3 (ArC), 136.5 (ArC), 134.8 (ArC), 128.2 (ArC), 123.7 (ArC), 121.0 (ArC), 120.2 (ArC), 119.2 (ArC), 111.7 (ArC), 98.4  $(ArC)$ , 74.3 (CHOH), 41.0 (CH<sub>2</sub>).

5.1.3.28. 2-(2-Pyridin-3-ylethyl)-1H-indole  $(19)^{50}$  $(19)^{50}$  $(19)^{50}$ . Prepared according to a literature procedure.<sup>51</sup> A suspension of indole ketone **7a** (111 mg, 0.47 mmol), hydrazine hydrate (47  $\mu$ l, 0.97 mmol) and potassium hydroxide pellets (130 mg, 2.32 mmol) 910 in ethylene glycol (5 mL) was subjected to microwave heating at 195 °C for 1 h ( $P \sim 40$  W,  $p = 1-3$  bar). The resulting clear dark orange solution was partitioned between dichloromethane and saturated aqueous ammonium chloride. The aqueous phase was extracted twice with dichloromethane and the combined organic layers were dried over MgSO<sub>4</sub>. Concentration of the filtered solution combined with the addition of cyclohexane yielded the title product as an ochre solid (87 mg, 84%), mp 139-142 °C. LC–MS  $R_{\rm t}$ 5.2 min (method 1);  $m/z$  [MH<sup>+</sup>] 223. <sup>1</sup>H NMR  $\delta$  10.95 (1H, br s, indole-H1), 8.41 (1H, s, ArH), 8.35 (1H, d,  $J = 4.8$  Hz, ArH), 7.61 (1H, 920 dd,  $J = 7.8$ , 1.6 Hz, ArH), 7.35 (1H, d,  $J = 7.8$  Hz, ArH),  $7.27 - 7.24$ (2H, overlapped, 2 ArH),  $6.95$  (1H, t,  $J = 7.5$  Hz, ArH),  $6.88$  (1H, t,  $J = 7.5$  Hz, ArH), 6.09 (1H, s, ArH), 3.00 (4H, overlapped, 2 CH<sub>2</sub>).<br><sup>13</sup>C NMR  $\delta$  150.1 (ArC), 147.8 (ArC), 139.4 (ArC), 137.2 (ArC), 136.5 (ArC), 136.3 (ArC), 128.8 (ArC), 123.9 (ArC), 120.7 (ArC), 119.7 (ArC), 119.1 (ArC), 111.2 (ArC), 99.2 (ArC), 32.2 (CH<sub>2</sub>), 29.6 (CH<sub>2</sub>). Anal. Calcd for C<sub>15</sub>H<sub>14</sub>N<sub>2</sub>: C, 81.05; H, 6.35; N, 12.60. Found: C, 80.33; H, 6.13; N, 12.15.

5.1.3.29. 1-(1H -Indol-2-yl)-2-(pyridin-3-yl)ethanone oxime (20). A suspension of ketone 7a (52 mg, 0.22 mmol), hydroxylamine hydrochloride (17 mg, 0.24 mmol) and pyridine (20  $\mu$ l, 0.25 mmol) in absolute ethanol (1 mL) was subjected to microwave heating at 120 $\degree$ C for 30 min. The resulting clear yellow solution was evaporated with silica and oxime 20 was purified by automated flash chromatography to yield a white solid (13 mg, 24%), mp 187–192 °C. LC–MS  $R_t$  5.2 min (method 1);  $m/z$  [MH<sup>+</sup>] 252. <sup>1</sup>H  $NMR \delta 11.43$  (1H, br s), 11.11 (1H, br s), 8.55 (1H, d, J = 1.8 Hz, ArH), 8.34 (1H, dd,  $J = 4.\overline{8}$ , 1.6 Hz, ArH),  $\overline{7}$ .65 (1H, d,  $J = 7.8$  Hz, ArH), 7.47  $(1H, d, J = 8.0$  Hz, ArH), 7.34  $(1H, d, J = 8.2$  Hz, ArH), 7.26  $(1H, dd,$  $J = 7.9$ , 4.8 Hz, ArH), 7.07 (1H, t,  $J = 7.6$  Hz, ArH), 6.93 (1H, t,  $J = 7.6$  Hz, ArH), 6.84 (1H, d,  $J = 2.1$  Hz, ArH), 4.08 (2H, s, CH<sub>2</sub>). <sup>13</sup>C NMR δ 150.3 (C), 149.5 (C), 147.9 (C), 137.7 (ArC), 136.5 (ArC), 134.2 (ArC), 134.0 (ArC), 128.1 (ArC), 124.1 (ArC), 123.2 (ArC), 120.9 (ArC), 119.7 (ArC), 112.2 (ArC), 103.7 (ArC), 28.8 (CH<sub>2</sub>).<sup>44</sup>

5.1.3.30. 1H-Indole-2-carboxylic acid pyridin-3-ylamide (21). A yellow suspension of 1H-indole-2-carboxylic acid (482 mg, 2.95 mmol) in thionyl chloride (5 mL) was boiled for 15 min, during which time the mixture became a clear dark brown solution. This was concentrated under reduced pressure and the solid residue was evaporated three times with toluene. 3-Aminopyridine (306 mg, 3.22 mmol) and diisopropylethylamine (785  $\mu$ l, 4.43 mmol) were dissolved in dry THF (2 mL) to give a clear yellowish solution which was cooled to  $0^{\circ}$ C. A solution of the above 1H-indole-2-carbonyl chloride residue in THF (3 mL) was added carefully to give a red-brown mixture. The ice/water bath was removed after 5 min and the reaction mixture was stirred for a further 25 min and then partitioned between ethyl acetate and water. The aqueous layer is extracted twice with ethyl acetate and the combined organic layers were washed with brine and dried over MgSO<sub>4</sub>. Chromatographic purification (the title product 960 eluted with 50–60% ethyl acetate) afforded amide 21 as a yellow solid (372 mg, 53%), decomposes without melting above 258  $\degree$ C.

LC–MS  $R_{\rm t}$  5.1 min (method 1);  $m/z$  [MH<sup>+</sup>] 238.  $^1$ H NMR  $\delta$  11.80 (1H, br s, indole-H1), 10.40 (1H, s, CONH), 8.95 (1H, d,  $J = 2.3$  Hz, ArH), 8.29 (1H, dd,  $I = 4.6$ , 1.2 Hz, ArH), 8.19 (1H, dd,  $I = 8.4$ , 1.6 Hz, ArH), 7.66 (1H, d, J = 8.0 Hz, ArH), 7.45-7.42 (2H, overlapped, 2 ArH), 7.38 (1H, dd, J = 8.3, 4.8 Hz, ArH), 7.21 (1H, t, J = 7.5 Hz, ArH), 7.05 (1H, t, J = 7.5 Hz, ArH). <sup>13</sup>C NMR  $\delta$  160.6 (CONH), 145.0 (ArC), 142.2 (ArC), 137.5 (ArC), 136.2 (ArC), 131.5 (ArC), 127.6 <sup>970</sup> (ArC), 127.5 (ArC), 124.6 (ArC), 124.2 (ArC), 122.4 (ArC), 120.6 (ArC), 113.0 (ArC), 104.9 (ArC). Anal. Calcd for  $C_{14}H_{11}N_3O$ : C, 70.87; H, 4.67; N, 17.71. Found: C, 70.66; H, 4.57; N, 17.76.

#### 5.2. Biology

#### 5.2.1. Protein expression and purification of recombinant human IDO

The coding region for human IDO (Ala2-Gly403) was cloned into a derivative of plasmid pET9 (Novagen). The recombinant plasmid, pETIDO, encodes a histidine tag at the N-terminus of IDO. Bacterial strain BL21 AI (Invitrogen) was used for overexpression of IDO and <sup>980</sup> transformed with the pETIDO plasmid. The transformed cells were grown on a rotary shaker at 37  $\degree$ C and 220 rpm to an OD600 of 1.2, in LB medium supplemented with 25  $\mu$ g/mL kanamycin, 50  $\mu$ g/mL  $L$ -tryptophan and 10  $\mu$ M bovine hemin (Sigma). The culture was cooled in a water/ice bath and supplemented again with  $50 \mu g$ / mL  $L$ -tryptophan and 10  $\mu$ M bovine hemin. The expression of Histagged IDO was induced by the addition of  $1\%$  (w/v) arabinose. Induced cells were grown at 20 $\degree$ C and 60 rpm for 20 h. Cells (1 L culture) were collected by centrifugation, resuspended in 40 mL of 25 mM Mes, 150 mM KCl, 10 mM imidazole, and protease inhibitors (complete EDTA free, Roche Applied Science) (pH 6.5), and disrupted with a French press. The extract was clarified by centrifugation and filtration on a 0.22 µm filter. The enzyme was purified by IMAC using Ni2þ as a ligand and an IMAC HITRAP column (5 mL, GE Healthcare). Briefly, the extract was loaded on the column with

### 25 mM Mes, 150 mM KCl, and 10 mM imidazole (pH 6.5). The column was washed with 50 mL of the same buffer with the imidazole concentration adjusted to 100 mM. Finally, the protein was eluted with 25 mM Mes, 150 mM KCl, and 50 mM EDTA (pH 6.5). The buffer was then exchanged into 25 mM Mes and 150 mM KCl (pH 6.5) <sup>1000</sup> using a HITRAP desalting column (GE Healthcare). The purity of the enzyme was estimated to be >95% based on the SDS–PAGE gel and Coomassie blue staining. The ratio of absorbance at 404 nm to that at 280 nm of the protein was around 1.9.

#### 5.2.2. Assay buffer

Assay buffer was composed of Phosphate Buffer 50 mM (pH 6.5), Methylene Blue 5  $\mu$ M (Sigma–Aldrich, M9140), and Ascorbic Acid 40 mM (Sigma–Aldrich, A5960). L-Tryptophan substrate (Sigma-Aldrich, T0254) is diluted at 100  $\mu$ M in Assay Buffer. IDO enzyme was diluted in Phosphate Buffer 50 mM at 0.05 mg/ml. 1010 Hydrogen peroxide solution (Sigma–Aldrich, H1009) is prepared at a concentration of 10 mM in water.

#### 5.2.3. Enzymatic assay

Enzymatic Assay was performed in 384 Wells microplates (Greiner bio one, 781108). Ninety five microliters of reaction buffer with  $L$ -Tryptophan substrate at 100  $\mu$ M and compound to be tested are added in the microplate. The optical density at 320 nm of this mixture is first recorded as a baseline with a Spectramax plus spectrophotometer (Molecular Devices). The hIDO enzyme (0.375 µg in  $5$  µl) is then added to the mixture. After 60 min incubation at room 1020 temperature, the reaction is stopped by the addition of hydrogen peroxide (1 mM) and a second optical reading at 320 nm is performed. The difference between the two density values reflects the level of tryptophan degradation and is used to calculate the percentage of IDO inhibition relative to the reference inhibitor 1-MT. Dose–response inhibition of IDO is analyzed using GraphPad Prism 4 program (GraphPad Software Inc.).

#### 5.3. Cellular tests

#### 5.3.1. Cell line

A plasmid construct encoding murine IDO was transfected into mouse mastocytoma line P815B. Clone P185B-mIDO clone 6 (Uyt- 1030 tenhove et al, 2003), which overexpresses IDO, was selected and used for the cellular assay. Mouse IDO shares a 62% sequence identity with human IDO, and the active site residues are 100% conserved.

#### 5.3.2. Assay

The assay was performed in 96-well flat bottom plates seeded with  $2 \times 10^5$  cells in a final volume of 200 µl. To determine whether compounds were significant IDO, the cells were incubated overnight at 37 °C in HBSS (Hanks Balanced Salt Solution, Invitrogen) supplemented with 80  $\mu$ M L-Tryptophan and 20  $\mu$ M of the 1040 compound. The plates were then centrifuged 10 min at 300 g, and 150 µl of the supernatant were collected. The supernatant was analyzed by HPLC to measure the concentration of tryptophan and kynurenine, based on the retention time and the UV absorption (280 nm for tryptophan, 360 nm for kynurenine). For the HPLC analysis, 50  $\mu$ l of supernatant were mixed with 500  $\mu$ l acetonitrile to precipitate the proteins. After centrifugation, the supernatant was collected, concentrated on a speedvac, resuspended in a final volume of  $100 \mu l$  water and injected in the HPLC (C18 column). In those conditions, about 50% of the initial amount of tryptophan  $1050$ was degraded in the absence of inhibitor, and an equimolar amount of kynurenine was produced. The percentages of inhibition of tryptophan degradation and kynurenine production by the compounds were calculated in reference to this maximal activity. The initial wells containing the cells in the remaining volume of 50  $\mu$ l were used to estimate cell viability in a classical MTT assay. To that end, 50 µl of culture medium (Iscove medium with 10% FCS and amino acids) were added to the wells together with  $50 \mu l$  of MTT. After 3–4 h of incubation at 37 °C, 100 µl of SDS/DMF were added to dissolve the crystals of formazan blue and the absorbance 1060 at  $570/650$  nm was measured after overnight incubation at 37 °C.

#### 5.4. X-ray determination

X-ray measurements were performed on a Gemini Ultra R system (4-circle kappa platform, Ruby CCD detector) using  $M_0K_0$ radiation ( $\lambda$  = 0.71073 Å) for the two compounds. After mounting and centering of the single crystals on the diffractometer, cell parameters were estimated from a pre-experiment run and full datasets collected at room temperature. Software for data reduc-tion was CrysAliPro.<sup>[52](#page-14-0)</sup> Structures were solved by direct methods from  $\frac{\text{SHELXS-97}}{1070}$  program<sup>[53](#page-14-0)</sup> and then refined on  $F^2$  using  $\frac{\text{SHELXL-97}}{1070}$  1070 software. Non-hydrogen atoms were anisotropically refined and the hydrogen atoms in the riding mode with isotropic temperature factors fixed at 1.2 times U (equiv) of the parent atoms.

CCDC 794450 and 794451 contain the supplementary crystallographic data for this paper and can be obtained free of charge via [www.ccdc.cam.ac.uk/conts/retrieving.html](http://www.ccdc.cam.ac.uk/conts/retrieving.html) (or from the Cambridge Crystallographic Data Centre, 12, Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033; or [deposit@ccdc.cam.ac.uk\)](http://deposit@ccdc.cam.ac.uk).

#### 5.5. Molecular modeling

Molecular modeling studies were carried out on a Linux work-<br>1080 station. The compounds were built using the SKETCH module implemented in  $SVD$  (version 8.0).<sup>54</sup> Docking was performed using the 3D-coordinates of TDO (pdb code 2D0T) with the help of the

Please cite this article in press as: Dolušić, E.; et al. Bioorg. Med. Chem. (2010), doi:[10.1016/j.bmc.2010.12.032](http://dx.doi.org/10.1016/j.bmc.2010.12.032)

<span id="page-14-0"></span>

12 E. Dolušic´ et al. / Bioorg. Med. Chem. xxx (2010) xxx–xxx

automated  $GOD$  program<sup>35</sup> (active site definition: residues within 7 Å around PIM). In order to take protein flexibility into account, the enzyme-inhibitor complexes were optimized using the MINIMIZE module. The minimization process uses the Powell method with the Tripos force field (dielectric constant 1r) to reach a final convergence of 0.01 kcal mol $^{-1}$ .

#### <sup>1090</sup> Acknowledgments

This work is supported by the FNRS (Belgium) and Biowin (CANTOL: convention  $n°5678$ ). R.F. is greatly indebted to the Belgian 'Fonds de la Recherche Scientifique—FNRS' for the award of a postdoctoral research grant.

#### Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.bmc.2010.12.032](http://dx.doi.org/10.1016/j.bmc.2010.12.032).

#### References and notes

- 1. Munn, D. H.; Mellor, A. L. J. Clin. Invest. 2007, 117, 1147.
- Prendergast, G. C. Oncogene 2008, 27, 3889.
- 3. Rafice, S. A.; Chauhan, N.; Efimov, I.; Basran, J.; Raven, E. L. Biochem. Soc. Trans. 2009, 37, 408.
- Sono, M.; Cady, S. G. Biochemistry 1989, 28, 5392.
- 5. Shimizu, T.; Nomiyama, S.; Hirata, F.; Hayaishi, O. J. Biol. Chem. 1978, 253, 4700. 6. Maghzal, G. J.; Thomas, S. R.; Hunt, N. H.; Stocker, R. J. Biol. Chem. 2008, 283,
- 12014. 7. Vottero, E.; Mitchell, D. A.; Page, M. J.; MacGillivray, R. T.; Sadowski, I. J.;
- Roberge, M.; Mauk, A. G. FEBS Lett. 2006, 580, 2265. 8. Hwu, P.; Du, M. X.; Lapointe, R.; Do, M.; Taylor, M. W.; Young, H. A. J. Immunol. 2000, 164, 3596.
- Dantzer, R.; O'Connor, J. C.; Freund, G. G.; Johnson, R. W.; Kelley, K. W. Nat. Rev. Neurosci. 2008, 9, 46.
- 10. Schwarcz, R. Curr. Opin. Pharmacol. 2004, 4, 12.
- 11. Stone, T. W.; Darlington, L. G. Nat. Rev. Drug Discov. 2002, 1, 609.
- 12. Uyttenhove, C.; Pilotte, L.; Theate, I.; Stroobant, V.; Colau, D.; Parmentier, N.; Boon, T.; Van den Eynde, B. J. Nat. Med. 2003, 9, 1269.
- 13. Muller, A. J.; DuHadaway, J. B.; Donover, P. S.; Sutanto-Ward, E.; Prendergast, G. C. Nat. Med. 2005, 11, 312.
- 14. Okamoto, A.; Nikaido, T.; Ochiai, K.; Takakura, S.; Saito, M.; Aoki, Y.; Ishii, N.; Yanaihara, N.; Yamada, K.; Takikawa, O.; Kawaguchi, R.; Isonishi, S.; Tanaka, T.; Urashima, M. Clin. Cancer Res. 2005, 11, 6030.
- 15. Muller, A. J.; Malachowski, W. P.; Prendergast, G. C. Expert Opin. Ther. Targets 2005, 9, 831.
- 16. Muller, A. J.; Prendergast, G. C. Cancer Res. 2005, 65, 8065.
- 17. Muller, A. J.; Prendergast, G. C. Curr. Cancer Drug Targets 2007, 7, 31.
- 18. Macchiarulo, A.; Camaioni, E.; Nuti, R.; Pellicciari, R. Amino Acids 2009, 37, 219.
- 19. Cady, S. G.; Sono, M. Arch. Biochem. Biophys. 1991, 291, 326. 20. Eguchi, N.; Watanabe, Y.; Kawanishi, K.; Hashimoto, Y.; Hayaishi, O. Arch.
- Biochem. Biophys. 1984, 232, 602.
- <sup>1130</sup> 21. Banerjee, T.; Duhadaway, J. B.; Gaspari, P.; Sutanto-Ward, E.; Munn, D. H.; Mellor, A. L.; Malachowski, W. P.; Prendergast, G. C.; Muller, A. J. Oncogene 2008, 27, 2851.
- 22. Gaspari, P.; Banerjee, T.; Malachowski, W. P.; Muller, A. J.; Prendergast, G. C.; DuHadaway, J.; Bennett, S.; Donovan, A. M. J. Med. Chem. 2006, 49, 684.
- 23. Brastianos, H. C.; Vottero, E.; Patrick, B. O.; Van Soest, R.; Matainaho, T.; Mauk, A. G.; Andersen, R. J. J. Am. Chem. Soc. 2006, 128, 16046.
- 24. Pereira, A.; Vottero, E.; Roberge, M.; Mauk, A. G.; Andersen, R. J. J. Nat. Prod. 2006, 69, 1496.
- 25. Sugimoto, H.; Oda, S.; Otsuki, T.; Hino, T.; Yoshida, T.; Shiro, Y. Proc. Natl. Acad. Sci. U.S.A. 2006, 103, 2611.
- 26. Kumar, S.; Jaller, D.; Patel, B.; LaLonde, J. M.; DuHadaway, J. B.; Malachowski, W. P.; Prendergast, G. C.; Muller, A. J. J. Med. Chem. 2008, 51, 4968.
- 27. Nakashima, H.; Uto, Y.; Nakata, E.; Nagasawa, H.; Ikkyu, K.; Hiraoka, N.; Nakashima, K.; Sasaki, Y.; Sugimoto, H.; Shiro, Y.; Hashimoto, T.; Okamoto, Y.; Asakawa, Y.; Hori, H. Bioorg. Med. Chem. 2008, 16, 8661.
- 28. Carr, G.; Chung, M. K.; Mauk, A. G.; Andersen, R. J. J. Med. Chem. 2008, 51, 2634.
- 29. Yue, E. W.; Douty, B.; Wayland, B.; Bower, M.; Liu, X.; Leffet, L.; Wang, Q.; Bowman, K. J.; Hansbury, M. J.; Liu, C.; Wei, M.; Li, Y.; Wynn, R.; Burn, T. C.; Koblish, H. K.; Fridman, J. S.; Metcalf, B.; Scherle, P. A.; Combs, A. P. J. Med. 1150 Chem. 2009, 52, 7364.
- 30. Terentis, A. C.; Freewan, M.; Sempertegui Plaza, T. S.; Raftery, M. J.; Stocker, R.; Thomas, S. R. Biochemistry 2010, 49, 591.
- 31. Matsuno, K.; Takai, K.; Isaka, Y.; Unno, Y.; **Satoman, M.; Takikawa, O.; Asai, A. Bioorg.** Med. Chem. Lett. 2010, 20, 5126.
- 32. Rohrig, U. F.; Awad, L.; Grosdidier, A.; Larrieu, P.; Stroobant, V.; Colau, D.; Cerundolo, V.; Simpson, A. J.; Vogel, P.; Van den Eynde, B. J.; Zoete, V.; Michielin, O. J. Med. Chem. 2010, 53, 1172.
- 33. Katz, J. B.; Muller, A. J.; Prendergast, G. C. Immunol. Rev. 2008, 222, 206.
- 34. Carr, R. A.; Congreve, M.; Murray, C. W.; Rees, D. C. Drug Discovery Today 2005, 1160 10, 987.
- 35. Jones, G.; Willett, P.; Glen, R. C.; Leach, A. R.; Taylor, R. J. Mol. Biol. 1997, 267, 727.
- 36. Giorgioni, G.; Accorroni, B.; Di Stefano, A.; Marucci, G.; Siniscalchi, A.; Claudi, F. Med. Chem. Res. 2005, 14, 57.
- 37. Posner, G. H.; Whitten, C. E.; McFarland, P. E. J. Am. Chem. Soc. 1972, 94, 5106.
- 38. Sato, T.; Naruse, K.; Enokiya, M.; Fujisawa, T. Chem. Lett. 1981, 1135.
- 39. Shinoda, P.; Tadokoro, M.; Tsukube, H.; Arakawa, R. Chem. Commun. 1998, 181. 40. Colucci, M. C.; Reigan, P.; Siegel, D.; Chilloux, A.; Ross, D.; Moody, C. J. J. Med. **Chem. 2007**, 50, 5780. 1170
- 41. Lehmann, F.; Holm, M.; Laufer, S. Tetrahedron Lett. 2009, 50, 1708.
- 42. Capkova, K.; Yoneda, Y.; Dickerson, T. J.; Janda, K. D. Bioorg. Med. Chem. Lett. 2007, 17, 6463.
- 43. Sundberg, R. G.; Luis, J. G.; Parton, R. L.; Schreiber, S.; Srinivasan, P. C.; Lamb, P.; Forcier, P.; Bryan, R. B. J. Org. Chem. 1978, 23, 4859.
- 44. Lavilla, R.; Gotsen, T.; Gavalda, J.; Josep, M.; Santano, M. C.; Bosch, J. J. Chem.
- Res., Synop. 1996, 8, 380. 45. Troin, Y.; Diez, A.; Bettiol, J.-L.; Rubiralta, M.; David, S.; Husson, H.-P. Heterocycles 1991, 32, 663.
- 46. Prieto, M.; Zurita, E.; Rosa, E.; Muñoz, L.; Lloyd-Williams, P.; Giralt, E. J. Org. 1180 Chem. 2004, 69, 6812.
- 47. Sundberg, R. J.; Smith, F. X. J. Org. Chem. 1975, 40, 2613.
- 48. Facoetti, D.; Abbiati, G.; Rossi, E. Eur. J. Org. Chem. 2009, 2872.
- 49. Salas, M.; Joule, J. A. J. Chem. Res., Synop. 1990, 3, 98.
- 50. Mohan, B.; Nagarathnam, D.; Vedachalam, M.; Srinivasan, P. C. Synthesis 1985, 2, 188.
- 51. Mahboobi, S.; Uecker, A.; Sellmer, A.; Cenac, C.; Hocher, H.; Pongratz, H.; Eichhorn, E.; Hufsky, H.; Trumpler, A.; Sicker, M.; Heidel, F.; Fischer,<br>T.; Stocking, C.; Elz, S.; Bohmer, F. D.; Dove, S. J. *Med. Chem*. **2006**, 49,  $3101.$  1190
- 52. CrysAlisPro. Oxford Diffraction Ltd.
- 53. Sheldrick, G. M. Acta Crystallogr., Sect. A 2008, 64, 112.
- 54. SYBYL 8.0, Tripos Inc. 1699 South Hanley Rd., St. Louis, Missouri 63144, USA.