

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

From contradiction to conciliation

Lambert, Dominique; Hespel, Bertrand

Publication date:
2011

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (HARVARD):
Lambert, D & Hespel, B 2011, *From contradiction to conciliation: a way to "dualize" sheaves*. NAXYS , no. 12-2011, FUNDP. Namur center for complex systems, Namur.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



From contradiction to conciliation:
a way to “dualize” sheaves

by D. Lambert and B. Hespel

Report naXys-12-2011

12 04 2011



Namur Center for Complex Systems

University of Namur
8, rempart de la vierge, B5000 Namur (Belgium)

<http://www.naxys.be>

From contradiction to conciliation: a way to “dualize” sheaves ¹

D. Lambert and B. Hespel

naXys
University of Namur

Abstract

Our aim is to give some insights about how to approach the formal description of situations where one has to conciliate several contradictory statements, rules, laws or ideas. We show that such a conciliation structure can be naturally defined on a topological space endowed with the set of its closed sets and that this specific structure is a kind of “dualization” of the sheaf concept where “global” and “local” levels are reversed.

1 Towards a model of conciliation?

Our aim here is to give some insights about how to approach the formal description of situations where one has to conciliate several statements, rules, laws, ideas that can be contradictory.

In fact it is straightforward to notice that many real situations in human social and political life face us with the need to conciliate contradictory points of view and to find some agreement. Reaching such an agreement does not imply that all contradictions have vanished. The democratic practice shows us that in spite of real contradictions, institutions can survive fruitfully. Of course one realizes easily two things. First, the number of contradictory views or positions has to remain under a certain level above which some dangerous tensions could appear with the risk to destroy completely the relations network. Thus contradictions have to be confined in some restricted and controlled fields in order to avoid complete percolation in the system under consideration. Second, not all contradictory theses can coexist peacefully or without damage for the coherence of institutions, laws or thoughts. But, in some circumstances, contradictions survive coherently in a system, institution or in law corpus because they happen to be interpretations, readings, of a common yet ambiguous statement, idea, rule, law, ... In fact this means that contradictions can merge in a coherent framework if they are rooted in a common soil from which they can be derived as two non-equivalent but possible interpretations.

In this paper we try to formalize this kind of realization. In order to do this, we exhibit a structure called *conciliation* tailored for describing situations in which we impose that two different and even contradictory “things” (ideas, statements, rules, images, ...) defined “locally” (i.e. at the level of a restricted social or political groups, law corpus, ...) can be identified (conciliated) at a higher level, “globally” if you want (for example at the level of a country, or of the international law, ...), if and only if there exists at a “deeper level” (for example at the level of citizens, of articles of laws, ...) a unique element which can be map on both initial “things”. The philosophy of conciliation building is to impose global constraint (forcing the conciliation of objects at high level) and then to exhibit some objects locally that enable the conciliation. The latter are roots providing a deep relation between what can be considered as different or even contradictory at a local level. In fact the existence of such “roots” is not trivial and their existence corresponds precisely to what we mean by a *conciliation*.

¹Both of the authors wish to warmly thank Daniel Dzierzgowski for having greatly improved their manuscript and Professor Chris Mortensen for having quickly shown an interest in their attempt.

We will see that such a conciliation structure happens to be naturally defined on a topological space endowed with the set of its closed sets. This conciliation structure will also appear to be a kind of “dualization” of the sheaf concept where “global” and “local” levels are reversed. In a sheaf, we impose, at a deep level, conditions that enable us to glue coherently objects defined locally in such a way that if the sheaf exists, we can build some new global object whose restrictions give rise to the initial object. So it is possible to use this sheaf-conciliation duality to build some new concepts.

2 The concept of conciliation on a topological space

Let X be a topological space and $\text{Fer}(X)$ the set of its closed sets.

A *pre-conciliation* is defined giving:

- (a) a map G which associates to each closed set W belonging to $\text{Fer}(X)$ a set $G(W)$;
- (b) for every $W_1, W_2 \in \text{Fer}(X)$ such that $W_2 \supseteq W_1$, a map $\delta_{W_1}^{W_2} : G(W_1) \rightarrow G(W_2)$, called a *mediation*, satisfying the following properties:

$$\forall W \in \text{Fer}(X) : \delta_W^W = \text{id}_W \quad (1)$$

where id_W is the identity map on W , and

$$\forall W_1, W_2, W_3 \in \text{Fer}(X) : W_3 \supseteq W_2 \supseteq W_1 \Rightarrow \delta_{W_1}^{W_3} \circ \delta_{W_1}^{W_2} = \delta_{W_1}^{W_3}. \quad (2)$$

A family $(W_i)_{i \in I}$ is a *finite closed co-covering* of $W \in \text{Fer}(X)$ iff I is finite, $\forall i \in I : W_i \in \text{Fer}(X)$ and $W = \bigcap_{i \in I} W_i$.

A pre-conciliation G in X is said to be *unified* iff it satisfies the following property:

- for every $W \in \text{Fer}(X)$, $W \neq \emptyset$,
- and every finite closed co-covering $(W_i)_{i \in I}$ of W ,
- we have: $\forall j \in I \forall t, s \in G(W) : \delta_{W_j}^{W_j}(t) = \delta_{W_j}^{W_j}(s) \Rightarrow t = s$.

A unified pre-conciliation G in X is a *conciliation* in X iff it satisfies the following property:

- for every $W \in \text{Fer}(X)$, $W \neq \emptyset$,
- every finite closed co-covering $(W_i)_{i \in I}$ of W ,
- and every $(t_i)_{i \in I}$ such that $\forall i \in I : t_i \in G(W_i)$,
- if $\forall i, j \in I : \delta_{W_i}^{W_i \cup W_j}(t_i) = \delta_{W_j}^{W_i \cup W_j}(t_j)$
- then $\exists! t \in G(W) \forall i \in I : \delta_{W_i}^{W_i}(t) = t_i$.

Intuitively, the latter property means the following thing. The reason why mediations have succeeded to conciliate, at the global level, ideas which are different or even contradictory, is the fact that there exists, at a deep level, a fundamental idea which enables this conciliation. And it works because this ground for mutual conciliation, rooting divergent ideas in a common soil.

3 Some examples of conciliations

Now we give an example of conciliation that will play a major role to connect this kind of structure to logic. For any $W_i \in \text{Fer}(X)$, let us consider the map:

$$W_i \mapsto \mathfrak{Z}(W_i) = \{Z \in \text{Fer}(X) \mid Z \supseteq W_i\}.$$

Let us define the maps $\delta_{W_i}^{W_j} : \mathfrak{Z}(W_i) \rightarrow \mathfrak{Z}(W_j) : Z_i \mapsto \delta_{W_i}^{W_j}(Z_i) = Z_i \cup W_j$. We check that properties (1), (2) and (3) are trivially satisfied. We want to show that \mathfrak{Z} is a conciliation in X .

Let us suppose that $\forall i, j \in I \forall Z_i \in \mathfrak{Z}(W_i) \forall Z_j \in \mathfrak{Z}(W_j) : \delta_{W_i}^{W_i \cup W_j}(Z_i) = \delta_{W_j}^{W_i \cup W_j}(Z_j)$, we have to prove that $\exists! Z \in \mathfrak{Z}(W_i \cap W_j) : \delta_{W_i \cap W_j}^{W_i}(Z) = Z_i$ and $\delta_{W_i \cap W_j}^{W_j}(Z) = Z_j$.

By hypothesis, we have: $\delta_{W_i}^{W_i \cup W_j}(Z_i) = \delta_{W_j}^{W_i \cup W_j}(Z_j) \Leftrightarrow Z_i \cup W_i \cup W_j = Z_j \cup W_i \cup W_j$.

But $Z_i \in \mathfrak{Z}(W_i)$ and $Z_j \in \mathfrak{Z}(W_j)$, therefore we get: $Z_i \cup W_j = Z_j \cup W_i$. The searched unique closed set Z is nothing but $Z_i \cap Z_j$. Indeed, we have

$$\begin{aligned} \delta_{W_i \cap W_j}^{W_i}(Z_i \cap Z_j) &= (Z_i \cap Z_j) \cup W_i = (Z_i \cup W_i) \cap (Z_j \cup W_i) = Z_i \cap (Z_j \cup W_i) \\ \delta_{W_i \cap W_j}^{W_j}(Z_i \cap Z_j) &= (Z_i \cap Z_j) \cup W_j = (Z_i \cup W_j) \cap (Z_j \cup W_j) = (Z_i \cup W_j) \cap Z_j. \end{aligned}$$

But we have $Z_i \subseteq Z_i \cup W_j = Z_j \cup W_i$, $Z_j \subseteq Z_j \cup W_i = Z_i \cup W_j$, then we can deduce that

$$\delta_{W_i \cap W_j}^{W_i}(Z_i \cap Z_j) = Z_i, \delta_{W_i \cap W_j}^{W_j}(Z_i \cap Z_j) = Z_j$$

and, because $\delta_{W_i \cap W_j}^{W_i}$ and $\delta_{W_i \cap W_j}^{W_j}$ are mediations satisfying property (3), that $Z_i \cap Z_j$ is the unique element of $\mathfrak{Z}(W_i \cap W_j)$ satisfying these two relations.

A second example of conciliation in X is given considering for any $W_i \in \text{Fer}(X)$ the map $1 : W_i \rightarrow \{*\} \upharpoonright_{W_i}$ which associates a singleton to each closed set. It is sufficient to set trivially: $\delta_{W_i}^{W_j}(* \upharpoonright_{W_i}) = * \upharpoonright_{W_j}$ and we check that this defines a conciliation in X .

Conciliation has natural links with the concept of extension. We will show this on a third example using the extension theorem of Tietze (see [18]). This states the following:

Let X be a normal space and let F one of its closed subsets and f a continuous function defined on F with values in a closed interval $[a, b]$ of the set of real numbers.

Then, f has a continuous extension g defined on X and with values in $[a, b]$.

Let us recall that a topological space is normal if it is a T_1 space in which two disjoint closed sets can always be separated by open sets. A T_1 space is a topological space in which each point is a closed set. We are considering now a compact Hausdorff topological space (a space is a Hausdorff space if two distinct points can always be separated by open sets, in the sense that they have open neighbourhood). A compact Hausdorff space is a normal space and furthermore every compact subspace of a Hausdorff space is closed. Let us thus consider two compact subspaces H_1 and H_2 of H such that $H_1 \cup H_2 = H$. Now, let $f_1 : H_1 \rightarrow [a, b]$ and $f_2 : H_2 \rightarrow [a, b]$ be two continuous functions. By Tietze extension theorem, we know that both functions admit two continuous extensions: $g_1 : H \rightarrow [a, b]$ and $g_2 : H \rightarrow [a, b]$. One can write: $f_j = g_j \circ J_j$ ($j = 1, 2$), where J_j denotes the inclusion of H_j in H . Let us call $G(H_1)$ and $G(H_2)$ the sets of continuous functions from the closed sets H_1 and H_2 in $[a, b]$. We check that extensions of these continuous

functions can be viewed as mediations of a conciliation G built on $H_1 \cap H_2 = H^\sharp$ which associates to any closed subset a set of continuous functions on this space and with values in $[a, b]$. We have effectively a conciliation, for if we set the following constraint:

$$g_1 = g_2, \quad g_1 := \delta_{H_1}^H, \quad g_2 := \delta_{H_2}^H,$$

we check that

$$\exists! g : H_1 \cap H_2 \rightarrow H : \delta_{H_1 \cap H_2}^{H_1} g = f_1 \text{ and } \delta_{H_1 \cap H_2}^{H_2} g = f_2.$$

This follows from the fact that $f_1 \circ K_1 = g_1 \circ J_1 \circ K_1 \equiv g_2 \circ J_2 \circ K_2 = f_2 \circ K_2 := g$, where K_j is the inclusion of $H_1 \cap H_2$ in H_j and from the fact that $J_1 \circ K_1 = J_2 \circ K_2 : H_1 \cap H_2 \rightarrow H$. It is interesting to note that in Tietze theorem, the fact that the subspace is closed is crucial. If this subspace is not closed, one cannot ensure the existence of a continuous extension and the constraint to get a conciliation is not satisfied (let us take for example $H = [0, 1]$, $H_1 =]0, 1]$, $f_1 : H_1 \rightarrow H : x \mapsto f_1(x) = \sin(1/x)$, the space H_1 is not closed in H and we easily check that f_1 has no continuous extension on all of H). This shows that the fact that the definition of conciliation starts from closed sets is not at all artificial. On the contrary the use of closed sets seems dictated by the nature of mediations, whose particular case is extensions.

4 Conciliations as cosheaves

Let us note that a pre-conciliation G can be considered as a *covariant functor* from the category of closed sets on X (whose objects are the closed sets and whose arrows are closed sets inclusions) in the category of sets. This functor is described by the following commutative diagram:

$$\begin{array}{ccc} W_j & \xrightarrow{G} & G(W_j) \\ \subseteq \uparrow & & \uparrow \delta_{W_i}^{W_j} \\ W_i & \xrightarrow{G} & G(W_i) \end{array}$$

The above definition of a conciliation in X happens to be a kind of “dualization” of the sheaf concept on X (in that sense a conciliation can be called a “cosheaf”). We know indeed that a sheaf F on X can be defined as follows (see [8]).

Let X be a topological space and $\text{Ouv}(X)$ the set of its open sets.

A *presheaf* F on X is defined giving:

- (a) a map F which associates to each open set U of $\text{Ouv}(X)$ a set $F(U)$;
- (b) for every $U_1, U_2 \in \text{Ouv}(X)$ such that $U_1 \subseteq U_2$, a map $\rho_{U_1}^{U_2} : F(U_2) \rightarrow F(U_1)$, called a *restriction*, satisfying the following properties:

$$\forall U \in \text{Ouv}(X) : \rho_U^U = \text{id}_U \tag{5}$$

where id_U is the identity map on U and

$$\forall U_1, U_2, U_3 \in \text{Ouv}(X) : U_1 \subseteq U_2 \subseteq U_3 \Rightarrow \rho_{U_1}^{U_3} = \rho_{U_1}^{U_2} \circ \rho_{U_2}^{U_3}. \tag{6}$$

A family $(U_i)_{i \in I}$ is a *finite open covering* of $U \in \text{Ouv}(X)$ iff I is finite, $\forall i \in I : U_i \in \text{Ouv}(X)$ and $U = \bigcup_{i \in I} U_i$.

A presheaf on F is said to be *separated* if we assume in addition that:

$$\begin{aligned} & \text{for every } U \in \text{Ouv}(X), \\ & \text{and every finite open covering } (U_i)_{i \in I} \text{ of } U, \\ & \text{we have: } \forall j \in I \forall t, s \in F(U) : \rho_{U_i}^U(t) = \rho_{U_i}^U(s) \Rightarrow t = s. \end{aligned} \quad (7)$$

A separated presheaf F on X is a *sheaf on X* iff it satisfies the following property:

$$\begin{aligned} & \text{for every } U \in \text{Ouv}(X), \\ & \text{every finite open covering } (U_i)_{i \in I} \text{ of } U, \\ & \text{and every } (t_i)_{i \in I} \text{ such that } \forall i \in I : t_i \in F(U_i), \\ & \text{we have: } \exists! t \in F(U) \forall i \in I : \rho_{U_i}^U(t) = t_i. \end{aligned} \quad (8)$$

We know also that a presheaf F on X can be considered as a contravariant functor from the category of open sets (whose objects are open sets and whose arrows are opens sets inclusions) in the category of sets, described by the following commutative diagram:

$$\begin{array}{ccc} U_j & \xrightarrow{F} & F(U_j) \\ \subseteq \uparrow & & \downarrow \rho_{W_i}^{U_j} \\ U_i & \xrightarrow{F} & F(U_i) \end{array}$$

We can push forward this kind of “duality” which relates sheaves and conciliations in order to generate new concepts.

If F is a presheaf on a topological space X , we can define at each point x of X a disjoint union: $\bigcup_{U \ni x} F(U)$. Now let us define an equivalence relation \approx_ρ : for every $U_1, U_2 \in \text{Ouv}(X), t_i \in U_i$,

$$t_1 \approx_\rho t_2 \Leftrightarrow \exists U \subseteq U_1 \cap U_2 : \rho_U^{U_1}(t_1) = \rho_U^{U_2}(t_2).$$

In other words, two “sections” of the sheaf are identified if they have the same restriction on an open set. Then, classically, one can introduce the concept of stalk F_x of F at a point x of X by:

$$F_x := \bigcup_{U_j \ni x} F(U_j) / \approx_\rho = \varinjlim_{U_j \ni x} F(U_j)$$

where the limit is the inductive limit induced by the restrictions:

$$\begin{aligned} F(U_0) & \xrightarrow{\rho_{U_1}^{U_0}} F(U_1) \xrightarrow{\rho_{U_2}^{U_1}} F(U_2) \xrightarrow{\rho_{U_3}^{U_2}} \dots \varinjlim_{U_j \ni x} F(U_j) \\ U_0 & \supseteq U_1 \supseteq U_2 \supseteq \dots \end{aligned}$$

Mimicking this, we can introduce a similar concept for a pre-conciliation G in X , called a *treaty* at x , by

$$G_x := \bigcup_{W_j \ni x} G(W_j) / \approx_\rho = \varinjlim_{W_j \ni x} G(W_j)$$

where the limit is the inductive limit induced by mediations:

$$\begin{aligned} G(W_0) & \xrightarrow{\delta_{U_1}^{W_0}} G(W_1) \xrightarrow{\delta_{U_2}^{W_1}} G(W_2) \xrightarrow{\delta_{U_3}^{W_2}} \dots \varinjlim_{W_j \ni x} G(W_j) \\ W_0 & \subseteq W_1 \subseteq W_2 \subseteq \dots \end{aligned}$$

Here, the equivalence relation \approx_δ is defined by the following equivalence: for every $W_1, W_2 \in \text{Fer}(X), t_i \in W_i$

$$t_1 \approx_\delta t_2 \Leftrightarrow \exists W \supseteq W_1 \cup W_2 : \delta_{W_1}^W(t_1) = \delta_{W_2}^W(t_2).$$

Elements of a *treaty* G_x at x are equivalence classes (the analog of what we call *germs* in sheaf theory). We propose to call such a class an *agreement* at x . Then classes of *agreements* constitute *treaties* and a pre-conciliation can be constructed from such treaties.

Starting from any pre-conciliation, we can build an associated *conciliation*. Let G be a *pre-conciliation*, we can consider the set G^a of the following maps defined on each closed set:

$$a_W : W \rightarrow \bigcup_{x \in X} G_x : x \mapsto a_W(x)$$

such that $a_W(x) \in G_x$. We are showing that G^a is in fact a conciliation. The *agreement* $a(x)$ is an equivalence class of the *treaty* G_x , namely an element $[(s, W)]$, where $s \in G(W), x \in W$ and $[(s, W)] = [(t, V)] \Leftrightarrow s \approx_\delta t$, if $x \in V$. One can then define, in a very natural way, the mediations of G^a by the following formula:

$$G^a \delta_{W_1}^W(a_W(x)) = G^a \delta_{W_1}^{W_2}([(s, W_1)]) := [(G^a \delta_{W_1}^{W_2}(s), W_2)]$$

with $W_1 \subseteq W_2$. One can then prove that G^a is a *conciliation*. This process, generating a conciliation from a preconciliation, could serve as the starting point to define the quotient of two conciliations G and H . Giving the map $W \rightarrow G(W)/H(W)$ whenever it has a meaning, we get a preconciliation Q . One can then define G/H as the associated conciliation Q^a .

5 Conciliations and paraconsistent logic of closed sets

First of all let us introduce the concept of Brouwer algebra (see [11]) which is the dual concept of Heyting algebra (see [1, pp. 129–134]). A Brouwer algebra, or co-Heyting algebra, denoted by $\langle B, \wedge, \vee, 0, 1, \overset{\circ}{\leftarrow} \rangle$, is a bounded distributive lattice with minimal element 0 and maximal element 1, endowed with a binary operation $\overset{\circ}{\leftarrow}$, called “pseudo-difference”, satisfying the following property:

$$\forall a, b, c \in B : (a \overset{\circ}{\leftarrow} b) \leq c \Leftrightarrow a \leq (b \vee c)$$

where \leq is the order relation defined on the lattice by: $a \leq b \Leftrightarrow a = a \wedge b \Leftrightarrow b = a \vee b$.

We know that Boolean algebra is related to classical propositional logic. We also know that Heyting algebra is related to intuitionistic propositional calculus. Brouwer algebra corresponds to paraconsistent propositional logic (see [11], [16], [3]). The correspondence goes as follows. To any proposition p of this paraconsistent logic one can associate an element $a(p)$ belonging to the Brouwer algebra $\langle B, \wedge, \vee, 0, 1, \overset{\circ}{\leftarrow} \rangle$. To the conjunction and to the disjunction, one can associate the binary operations \wedge and \vee , respectively. The contradiction \perp and the tautology \top are associated to the elements 0 and 1 respectively. Finally, the paraconsistent negation \neg corresponds to $1 \overset{\circ}{\leftarrow} a(p)$ in the Brouwer algebra. We recall that paraconsistent logic does not satisfy the non-contradiction principle, i.e., $p \wedge \neg p \leftrightarrow \perp$.

A classical example of Brouwer algebra is built from the set of closed sets of a topological space. We can check that $(\text{Fer}(X), \cap, \cup, \cdot, X, \overset{\circ}{\leftarrow})$ is a Brouwer algebra, where the pseudo-difference is defined as follows:

$$(Z \overset{\circ}{\leftarrow} W) := \overline{\mathbb{C}_X W} \cap Z$$

(where $\mathbb{C}_X W$ denotes the complement of W in X and where the bar denotes the closure of a set, i.e. the least closed set containing this set). In this algebraic interpretation, the paraconsistent negation $\neg p$ corresponds to

$$(X \overset{\circ}{\leftarrow} W(p)) := \overline{\mathbb{C}_X W(p)} \cap X = \overline{\mathbb{C}_X W(p)}.$$

As the space X let us take the real line and let us define $\text{Fer}(X)$ as the set of its closed intervals. We immediately check that the noncontradiction principle $p \wedge \neg p \leftrightarrow \perp$ is not satisfied. Indeed, if we choose $W(p) = [0, 1]$ we get:

$$W(p) \cap \overline{\mathbb{C}_X W(p)} = \{0\} \cup \{1\} \neq \emptyset.$$

Intuitively we can say that contradiction is confined only on the border of the closed set. This confinement prevents contradictions to invade all logical deductions and gives to this paraconsistent logic a non-trivial character.

In fact one can prove that any Brouwer algebra is a subalgebra of a Brouwer algebra of closed sets of a particular topological space (see [11]). This result corresponds to another one that is well-known in the theory of Heyting algebras saying that every Heyting algebra is a subalgebra of a Heyting algebra of open sets of a topological space. We know also this classical result stating that every Boolean algebra can be viewed as a subalgebra of a Boolean algebra of subsets of a set.

Let us consider the set $\mathfrak{Z}(W_i) = \{Z \in \text{Fer}(X) \mid Z \supseteq W_i\}$ defined above. Following James and Mortensen (see [12]), let us show that $\langle \mathfrak{Z}(W_i), \cap, \cup, \emptyset, X, \overset{\circ}{\leftarrow}_{W_i} \rangle$ is a Brouwer algebra, where the pseudo-difference is given by:

$$(Z \overset{\circ}{\leftarrow}_{W_i} W) := (\overline{\mathbb{C}_X W} \cap Z) \cup W_i$$

for all $Z, W \in \mathfrak{Z}(W_i)$.

We have to prove that, for every $Z, W, T \in \mathfrak{Z}(W_i)$,

$$(Z \overset{\circ}{\leftarrow}_{W_i} W) \subseteq T \Leftrightarrow Z \subseteq (W \cup T),$$

i.e.:

$$(\overline{\mathbb{C}_X W} \cap Z) \cup W_i \subseteq T \Leftrightarrow Z \subseteq (W \cup T).$$

But this is easily checked by set considerations. Therefore \mathfrak{Z} is a conciliation in X . Let us emphasize the fact that in this definition we take the union with W_i in order that $(Z \overset{\circ}{\leftarrow}_{W_i} W)$ belongs to $\mathfrak{Z}(W_i)$. In fact, in general, if $Z, W \in \mathfrak{Z}(W_i)$ we do *not* have:

$$\overline{\mathbb{C}_X W} \cap Z = (Z \overset{\circ}{\leftarrow} W) \in \mathfrak{Z}(W_i).$$

The conciliation \mathfrak{Z} in X is a map which associates to any closed set W of X , a particular Brouwer algebra built with all closed sets containing W . The definition of the conciliation \mathfrak{Z} is coherent with the structure of the Brouwer algebra due to the fact that mediations are preserving the pseudo-difference $\overset{\circ}{\leftarrow}_{W_i}$, i.e.

$$\forall W_i \subseteq W_j : \delta_{W_i}^{W_j}(Z \overset{\circ}{\leftarrow}_{W_i} W) = \delta_{W_i}^{W_j}(Z) \overset{\circ}{\leftarrow}_{W_j} \delta_{W_i}^{W_j}(W).$$

The proof of this statement follows from a result previously obtained by James et Mortensen and described in an unpublished book ([5]):

$$\begin{aligned}\delta_{W_i}^{W_j}(Z \xleftarrow[\circ]{W_i} W) &= ((\overline{\mathbb{C}_X W} \cap Z) \cup W_i) \cup W_j = (\overline{\mathbb{C}_X W} \cap Z) \cup W_j = Z \xleftarrow[\circ]{W_i} W \\ \delta_{W_i}^{W_j}(Z) \xleftarrow[\circ]{W_j} \delta_{W_i}^{W_j}(W) &= (\overline{\mathbb{C}_X(W \cup W_j)} \cap (Z \cup W_j)) \cup W_j\end{aligned}$$

Then, we check that:

$$\begin{aligned}(\overline{\mathbb{C}_X(W \cup W_j)} \cap (Z \cup W_j)) \cup W_j &= \overline{(\mathbb{C}_X(W \cup W_j) \cap (Z \cup W_j)) \cup W_j} \\ &= \overline{(\mathbb{C}_X(W \cup W_j) \cup W_j) \cap ((Z \cup W_j) \cup W_j)} \\ &= \overline{(\mathbb{C}_X(W \cup W_j) \cup W_j) \cap (Z \cup W_j)} \\ &= \overline{((\mathbb{C}_X(W) \cap \mathbb{C}_X(W_j)) \cup W_j) \cap (Z \cup W_j)} \\ &= \overline{((\mathbb{C}_X(W) \cup W_j) \cap (\mathbb{C}_X(W_j) \cup W_j)) \cap (Z \cup W_j)} \\ &= \overline{(\mathbb{C}_X(W) \cup W_j) \cap (Z \cup W_j)} \\ &= \overline{(\mathbb{C}_X(W) \cap (Z \cup W_j)) \cup (W_j \cap (Z \cup W_j))} \\ &= \overline{(\mathbb{C}_X(W) \cap Z) \cup (\mathbb{C}_X(W) \cup W_j) \cup W_j} \\ &= \overline{(\mathbb{C}_X(W) \cap Z) \cup W_j} \\ &= \overline{(\mathbb{C}_X(W) \cap Z)} \cup W_j \\ &= Z \xleftarrow[\circ]{W_j} W.\end{aligned}$$

6 The category of conciliations

If we want to compare conciliations in X , it is natural to introduce a notion of morphism of conciliations. Let G and H be conciliations on X . A morphism of conciliations $G \rightarrow H$ can be defined using the following commutative diagram:

$$\begin{array}{ccc} G(Z) & \longrightarrow & H(Z) \\ G\delta_W^Z \uparrow & & \uparrow H\delta_W^Z \\ G(W) & \longrightarrow & H(W) \end{array}$$

This means that the morphism is compatible with mediations of both conciliations (we have distinguished mediations of both conciliations with the help of an index at the left of the mediation symbol).

The set of conciliations in X and the morphisms of conciliations generate objects and arrows of a category: the category of conciliations on X denoted by $\mathbf{Ccs}(X)$. The axioms defining a category can be easily verified. For any conciliation, there exists an identity morphism and the composition of two morphisms is still a morphism. Furthermore, this composition is associative. The conciliation $1 : W_i \rightarrow \{*\}_{|W_i}$, together with the trivial mediations $1_{\delta_{W_i}^{W_j}}(*|_{W_i}) = *|_{W_i}$, happens to be a terminal object of $\mathbf{Ccs}(X)$. For any conciliation G , it exists effectively a unique morphism $G \rightarrow 1$ compatible with all mediations, i.e. rendering the following diagram

commutative:

$$\begin{array}{ccc} G(W_j) & \longrightarrow & 1(W_j) \\ \uparrow \delta_{W_i}^{G, W_j} & & \uparrow 1_{\delta_{W_i}^{W_j}} \\ G(W_i) & \longrightarrow & 1(W_i) \end{array}$$

It is interesting to wonder what are the subobjects of $\mathbf{Ccs}(\mathbf{X})$. What does a subconciliation mean? A subconciliation G' of G is a conciliation satisfying the following commutative diagram, for all $W_i \subseteq W_j$:

$$\begin{array}{ccc} G'(W_j) & \subseteq & G(W_j) \\ \uparrow \delta_{W_i}^{G', W_j} & & \uparrow \delta_{W_i}^{G, W_j} \\ G'(W_i) & \subseteq & G(W_i) \end{array}$$

This diagram means that inclusions remain always compatible with mediations. We are showing that $\mathbf{Ccs}(\mathbf{X})$ can be characterized by the following commutative diagram:

$$\begin{array}{ccc} G'(W_i) & \overset{j}{\subseteq} & G(W_i) \\ \downarrow & & \downarrow \psi_{G'} \\ 1(W_i) & \xrightarrow{\phi} & \mathfrak{Z}(W_i) \end{array}$$

where G' is a subconciliation of G . The maps $\psi_{G'}$ and ϕ are defined as follows:

- ▶ $\psi_{G'}(j(t_i)) = Z_{t_i} = \bigcap \{Z_k \mid Z_k \supseteq W_i \text{ and } t_i \in G'(W_i) \text{ and } \delta_{W_i}^{G, Z_k}(j(t_i)) \in G'(Z_k)\}$.

The image of this map is thus the least closed set containing W_i such that the mediation $\delta_{W_i}^{G, Z_k}$ of G lifts $j(t_i)$ inside $G'(W_i)$.

- ▶ $\phi : 1(W_i) \rightarrow \mathfrak{Z}(W_i) : * \mapsto W_i$.

This maps associates to $*$ the minimal element of $\mathfrak{Z}(W_i)$. When G' is a subconciliation of G , one checks that: $\forall t_i \in G'(W_i) : \psi_{G'}(j(t_i)) = \phi(*) = W_i$.

For two conciliations G_1 et G_2 , one can define their product, associating to each closed set W_i the Cartesian product of $G_1(W_i)$ and $G_2(W_i)$. Given these conciliations, one can define the set of morphisms relating both. The preceding commutative diagram shows also that $\mathbf{Ccs}(\mathbf{X})$ possess an object \mathfrak{Z} classifying subconciliations. This category has thus the properties of a topos. As $\mathfrak{Z}(W_i)$ can be endowed with a structure of Brouwer algebra (and not of Heyting algebra as in an ordinary topos), $\mathbf{Ccs}(\mathbf{X})$ is an example of *complement topos* (see Mortensen and Lavers in [12, p. 105]) (one could call it a co-Heyting or Brouwer topos); namely a topos whose classifying object is a Brouwer algebra. Let us note that in fact $\mathbf{Ccs}(\mathbf{X})$ gives an example of co-Heyting topos which is not a bi-Heyting topos (see [17]).

7 Towards a cohomology of conciliations

The aim of this section is to show that conciliation generates cohomology in the same way that sheaf theory generates sheaf cohomology (see [15], [14], [19]). Let $\{W_j\}_{j \in I}$ be a closed covering of X and let G be a conciliation (of rings) in X , which associates to any closed set W_j a ring $G(W_j) = \langle A, +, \cdot \rangle$. Now we define a *p-cochain with values in this conciliation*, as the map that associates to any union of closed sets $W_{j_0} \cup W_{j_1} \cup \dots \cup W_{j_p}$, $j_\alpha \in I$, $j_0 < j_1 < \dots < j_p$, an

element t_{j_0, j_1, \dots, j_p} of $G(W_{j_0} \cup W_{j_1} \cup \dots \cup W_{j_p})$. A p -cochain is thus a set $\{t_{j_0, j_1, \dots, j_p}\}_{j_0 < j_1 < \dots < j_p}$. The set of p -cochains will be denoted by $C^p(I, G)$. A 0-cochain is a set $\{t_j\}_{j \in I}$, $t_j \in G(W_j)$. A 1-cochain is a set $\{t_{j_0, j_1}\}_{j_0 < j_1}$, $t_{j_0, j_1} \in G(W_{j_0} \cup W_{j_1})$. Etc.

One can define a *coboundary operator* D such that:

$$D : C^p(I, G) \rightarrow C^{p+1}(I, G) : \{t_{j_0, j_1, \dots, j_p}\} \mapsto \{(Dt)_{j_0, j_1, \dots, j_p, j_{p+1}}\}$$

where

$$(Dt)_{j_0, j_1, \dots, j_p, j_{p+1}} = \sum_{k=0}^{p+1} (-1)^k \delta_{W_{j_0} \cup W_{j_1} \cup \dots \cup W_{j_k} \cup \dots \cup W_{j_{p+1}}}^{W_{j_0} \cup W_{j_1} \cup \dots \cup W_{j_{p+1}}} t_{j_0, j_1, \dots, \widehat{j_k}, \dots, j_p, j_{p+1}}$$

($\widehat{}$ means that you have to cancel the symbol). We check that D satisfies the property of a good coboundary operator, namely: $DD = 0$.

If we are computing the action of the coboundary operator on the 0-cochains, we get:

$$(Dt)_{j_0, j_1} = \delta_{W_{j_1}}^{W_{j_0} \cup W_{j_1}} t_{j_1} - \delta_{W_{j_0}}^{W_{j_0} \cup W_{j_1}} t_{j_0}$$

and if we put this to zero, we are lead to the condition (4) satisfied by all elements of $G(W_j)$ if G is a conciliation. Similarly, we can compute the action of the coboundary operator on the 1-cochains:

$$(Dt)_{j_0, j_1, j_2} = \delta_{W_{j_1} \cup W_{j_2}}^{W_{j_0} \cup W_{j_1} \cup W_{j_2}} t_{j_1, j_2} - \delta_{W_{j_0} \cup W_{j_2}}^{W_{j_0} \cup W_{j_1} \cup W_{j_2}} t_{j_0, j_2} + \delta_{W_{j_0} \cup W_{j_1}}^{W_{j_0} \cup W_{j_1} \cup W_{j_2}} t_{j_0, j_1}.$$

Now, let us suppose that the 1-cochain is defined by $t = Ds$. The preceding formula becomes:

$$\begin{aligned} (Dt)_{j_0, j_1, j_2} &= \delta_{W_{j_1} \cup W_{j_2}}^{W_{j_0} \cup W_{j_1} \cup W_{j_2}} \left(\delta_{W_{j_2}}^{W_{j_1} \cup W_{j_2}} s_{j_2} - \delta_{W_{j_1}}^{W_{j_1} \cup W_{j_2}} s_{j_1} \right) \\ &- \delta_{W_{j_0} \cup W_{j_2}}^{W_{j_0} \cup W_{j_1} \cup W_{j_2}} \left(\delta_{W_{j_2}}^{W_{j_0} \cup W_{j_2}} s_{j_2} - \delta_{W_{j_0}}^{W_{j_0} \cup W_{j_2}} s_{j_0} \right) \\ &+ \delta_{W_{j_0} \cup W_{j_1}}^{W_{j_0} \cup W_{j_1} \cup W_{j_2}} \left(\delta_{W_{j_1}}^{W_{j_0} \cup W_{j_1}} s_{j_1} - \delta_{W_{j_0}}^{W_{j_0} \cup W_{j_1}} s_{j_0} \right) \end{aligned}$$

Distributing médiations and applying the property (2) we get:

$$\begin{aligned} (Dt)_{j_0, j_1, j_2} &= \delta_{W_{j_2}}^{W_{j_0} \cup W_{j_1} \cup W_{j_2}} s_{j_2} - \delta_{W_{j_1}}^{W_{j_0} \cup W_{j_1} \cup W_{j_2}} s_{j_1} \\ &- \delta_{W_{j_2}}^{W_{j_0} \cup W_{j_1} \cup W_{j_2}} s_{j_2} + \delta_{W_{j_0}}^{W_{j_0} \cup W_{j_1} \cup W_{j_2}} s_{j_0} \\ &+ \delta_{W_{j_1}}^{W_{j_0} \cup W_{j_1} \cup W_{j_2}} s_{j_1} - \delta_{W_{j_0}}^{W_{j_0} \cup W_{j_1} \cup W_{j_2}} s_{j_0} \end{aligned}$$

this allows us to check that DDs is zero. As in the case of sheaves, we can introduce cohomology groups of conciliation setting: $H^p(I, G) = Z^p(I, G)/B^p(I, G)$ where $Z^p(I, G)$ is the set of p -cocycles, i.e. of p -cochains t such that $Dt = 0$ and where $B^p(I, G)$ is the set of p -coboundaries, i.e. of p -cochains b such that there exists $(p-1)$ -cochain s such that $b = Ds$. The definition of the cohomology group $H^p(I, G)$ is motivated by the fact that the p -cocycle t is only given up to a coboundary $b = Ds$, because $Dt = 0 = D(t + b) = D(t + Ds)$.

8 Globales and universales

We want to suggest now that the classical notions of *locales* and *quantales* have some dual analogs in the framework of conciliations.

A *locale* is a complete lattice L satisfying the following property:

$$x \wedge \bigvee Y = \bigvee (x \wedge Y), \quad x \wedge Y := \{x \wedge y \mid y \in Y \text{ and } Y \subseteq L\}.$$

In other words, L is a “*meet infinitely distributive*” complete distributive lattice.

A *quantale* $\langle Q, \wedge, \vee, \bullet, 0, 1 \rangle$ is a structure defined on a set Q such that

- (a) $\langle Q, \wedge, \vee \rangle$ is a complete lattice with nul element 0 and universal element 1,
- (b) $\langle Q, \bullet \rangle$ is a monoid with identity $1'$ for the operation \bullet
- (c) the operation \bullet is distributive with respect to \vee in the following sense:

$$\forall a \in Q \forall S \subseteq Q : a \bullet \bigvee S = \bigvee \{a \bullet s \mid s \in S\} \text{ and } (\bigvee S) \bullet a = \bigvee \{s \bullet a \mid s \in S\}$$

where $\bigvee S = \{\bigvee_j s_j \mid s_j \in S\}$.

One can immediately check that a locale is a quantale if we take $\forall a \in Q : a \bullet b := a \wedge b$.

One can prove that a *quantale* such that the operation \bullet is idempotent ($\forall a \in Q : a \bullet a = 1'$) and such that the universal element coincide with the monoid identity ($1 = 1'$) is a *locale*.

Let us now introduce the concept of *globale* as a complete lattice L satisfying the following property:

$$w \vee \bigwedge Z = \bigwedge (w \vee Z)$$

where $w \vee Z := \{w \vee z \mid z \in Z \text{ and } Z \subseteq L\}$.

In other words, L is a “*join infinitely distributive*” complete lattice.

An example of *globale* is given by the lattice of closed sets, $\text{Fer}(X)$, of a topological space X . Now, we define $\forall w_1, w_2 \in \text{Fer}(X), w_1 \wedge w_2 := w_1 \cap w_2$. As the union of an arbitrary number of closed sets is not necessarily closed, we have to define $\forall w_1 \in \text{Fer}(X) : \bigvee_j w_j := \overline{\bigcup_j w_j}$ using the closure of the set union. Both operations endowed $\text{Fer}(X)$ with a structure of complete lattice. We give here a proof of the well-known property saying that this lattice is effectively a “*join infinitely distributive*” lattice:

First of all we have:

$$\forall w \in \text{Fer}(X), z \in Z \subseteq \text{Fer}(X) : w \cup z \supseteq w \cup \bigcap Z.$$

We can deduce that:

$$\bigwedge (w \vee Z) = \bigcap (w \cup Z) \supseteq w \cup \bigcap Z = w \vee \bigwedge Z.$$

Let us note that we can write: $w \cup \bigcap Z = w \vee \bigwedge Z$ because $\bigcap Z$ is a closed set and the union of two closed sets is still a closed set.

We have now to prove that: $w \vee \bigwedge Z \supseteq \bigwedge (w \vee Z)$.

We have in fact $\forall z \in Z : w \cup z \supseteq \bigcap (w \cup Z)$. Defining $u = \bigcap (w \cup Z)$ we find $w \cup z \supseteq u$. But we have also:

$$\begin{aligned} z &= (w \cup z) \cap \mathcal{C}_W(w - z) \supseteq u \cap \mathcal{C}_X(w - z) = u \cap \mathcal{C}_X(w \cap \mathcal{C}_X z) = u \cap (\mathcal{C}_X \cup z) \\ &\Rightarrow z \supseteq (u \cap \mathcal{C}_X w) \cup (u \cap z). \end{aligned}$$

Taking the intersection $v = \bigcap Z$ of elements $z \in Z$, we get:

$$\begin{aligned} v &\supseteq (u \cap v) \cup (u \cap \mathcal{C}_X w) = [(u \cap v) \cup u] \cap [(u \cap v) \cup \mathcal{C}_X w] = u \cap (u \cup \mathcal{C}_X w) \cap (v \cup \mathcal{C}_X w) \\ &\Rightarrow v \supseteq u \cap (v \cup \mathcal{C}_X w) \\ &\Rightarrow v \cup w \supseteq [u \cap (v \cup \mathcal{C}_X w)] \cup w = (u \cup w) \cap (v \cup \mathcal{C}_X w \cup w) = u \cup w \supseteq u \\ &\Leftrightarrow w \cup v \supseteq u \Leftrightarrow w \cup \bigcap Z \supseteq \bigcap (w \cup Z). \end{aligned}$$

The lattice of closed sets is thus a globale.

Next we define a kind of “dual” version of the quantale. We will call it a *universale*. As a matter of definition one could propose the following statement: $\langle P, \wedge, \vee, 0, 1, \circ \rangle$ is a *universale* if the set P is such that:

- (a) $\langle P, \wedge, \vee, 0, 1, \circ \rangle$ is a complete lattice with zero element 0 and universal element 1.
- (b) $\langle P, \circ \rangle$ is a monoid with identity element $0'$ with respect to the operation \circ .
- (c) $\forall b \in P, \forall T \subseteq P : b \circ \bigwedge T = \bigwedge (b \circ T) = \bigwedge \{b \circ t \mid t \in T\}; (\bigwedge T) \circ b = \bigwedge (T \circ b)$.

One can show that such a structure has the following properties:

The operation \circ is such that $\forall a, b, c \in P : a \leq b \Rightarrow a \circ c \leq b \circ c$, where the order relation is the relation generated by the lattice: $a \leq b \Leftrightarrow a \wedge b; b = a \vee b$. We have indeed,

$$a \leq b \Leftrightarrow a = a \wedge b \Rightarrow a \circ c = (a \wedge b) \circ c = (a \circ c) \wedge (b \circ c) \Leftrightarrow a \circ c \leq b \circ c$$

If the zero element is identified with the identity element of the monoid operation ($0 = 0'$), we get: $\forall a, b \in P : a \circ b \geq a \vee b$. This is easily proved:

$$\begin{aligned} a \geq 0 = 0' &\Rightarrow a \circ b \geq 0' \circ b = b \Leftrightarrow a \circ b \geq b \Leftrightarrow b = (a \circ b) \wedge b \\ b \geq 0 = 0' &\Rightarrow a \circ b \geq a \circ 0' = a \Leftrightarrow a \circ b \geq a \Leftrightarrow a = (a \circ b) \wedge a \end{aligned}$$

So

$$\begin{aligned} a \vee b &= [(a \circ b) \wedge a] \vee [(a \circ b) \wedge b] \\ &\Leftrightarrow a \vee b = (a \circ b) \wedge (a \vee b) \\ &\Leftrightarrow a \vee b \leq a \circ b. \end{aligned}$$

We can exhibit the link between the *globale* and the *universale* by the following property: if the operation of the *universale* $\langle P, \wedge, \vee, 0, 1, \circ \rangle$ is idempotent and if the zero element of the lattice is identified with the identity element of the monoid, then $\langle P, \wedge, \vee, 0, 1 \rangle$ is a *globale*.

If $\langle P, \wedge, \vee, 0, 1 \rangle$ is a *globale* and if one identifies \circ to \vee , one gets $\forall a \in P : a \circ a = a \vee a = a$ and this shows the idempotency property. Furthermore, one check that

$$\forall a \in P : a \vee 0 = a = a \circ 0' = a \vee 0' \Rightarrow 0 = 0'$$

and that

$$a \circ \bigwedge_j b_j = a \vee \bigwedge_j b_j = \bigwedge_j (a \vee b_j) = \bigwedge_j (a \circ b_j).$$

This shows that $\langle P, \wedge, \vee, 0, 1 \rangle$ is a *universale* with an idempotent monoid operation and with the zero element of the lattice identified with the identity element of the monoid.

If now $\langle P, \wedge, \vee, 0, 1, \circ \rangle$ is a *universale* with idempotent monoid operation and with the universal element of the lattice identified with the identity element of the monoid, then if we identify \circ to \vee we get:

$$\forall a \in P, \forall T \subseteq P : a \circ \bigwedge T = \bigwedge (a \circ T) \Leftrightarrow a \vee \bigwedge T = \bigwedge (a \vee T)$$

and this shows that $\langle P, \wedge, \vee, 0, 1 \rangle$ is a *globale*.

9 Speculations

The main motivation of this paper was to build a formal model of (social, political, legal) situations where one reaches a conciliation of contradictory positions. This leads us to propose axioms defining a structure on a topological space that we called *conciliation*. It was surprising to realize that such a structure was in fact a kind of dualization of a sheaf; the duality here (which has not to be taken in the strict sense of category theory) has to be understood as the reversion of the role of global and local levels combined with the change from open to closed sets. In the case of a sheaf, we glue “locally” some objects defined on open sets, checking a coherence constraint on the intersections of open sets and then, we can reconstruct a well-defined “global” (i.e. at the level of the union of open sets) object of the same type. On the contrary, in the case of a conciliation, we conciliate globally some objects defined on closed sets, checking a coherence constraint on the union of closed sets, and then we can reconstruct well-defined “local” objects (i.e. at the level of intersections of closed sets) of the same type.

In other words, starting from trivial local objects, a sheaf allows us to build non-trivial global object by a gluing procedure. At the opposite, starting from trivial global objects, a conciliation allows us to generate some non-trivial local object.

Our paper may also help to shed some new light on a very important diagram occurring in the book of Shahn Majid, *Foundations of Quantum Group Theory* [9], but also in some of his papers (see for example [10]) and which illustrates the progress towards a search of a quantum gravity theory regulated by the meta-principle of self-duality expressing his representation-theoretic philosophy.

In the upper part of this diagram (see the figure below), we find concepts and theories related to quantum physics. We find, for example, Heyting algebra, which is the starting point for an understanding of quantum logic in which one drops the law of excluded middle. We find also quantales.

In the lower part of the diagram, one can find notions that are connected with general relativity and gravitation. Majid locates, in this part, co-Heyting algebras, i.e. Brouwer algebras. This is because the latter are related to logics in which one can drop the non-contradiction principle. As noted by Lawvere (see [6]), this fact enables us to introduce an operator $\partial p := p \wedge \neg p \neq 0$ which behaves as a derivation operator and which satisfies Leibniz rule: $\partial(p \wedge q) = (\partial p \wedge q) \vee (p \wedge \partial q)$. As derivation is the main ingredient to build vector fields in differential geometry, one can think that Brouwer algebra has something to do with the birth of geometry and step after step with general relativity and gravitation.

The upper and lower part of the diagram are related by duality. For example, if we replace open


sets by closed sets and the intuitionistic implication

$$(U \Rightarrow V) = ((\mathbb{C}_X U) \cup V)$$

($\mathbb{C}_X U$ being the interior of $\mathbb{C}_X U$) by the paraconsistent pseudo-difference ($Z \overset{\circ}{\Leftarrow} T = \overline{(\mathbb{C}_X T \cap Z)}$), we pass from the Heyting algebra in the upper part to Brouwer algebra in the lower part. The horizontal line separating both parts of the diagram corresponds to self-dual structures. An example of such structure is the Boolean algebra which can be considered as an algebra that is at the same time a Heyting and a Brouwer algebra. The set of subsets of a set is a Boolean algebra and its elements are at the same time open and closed sets, what the logicians called clopen sets. This is once again a facet of duality.

It is possible to improve Majid's inspiring diagram adding on the horizontal line "bi-Heyting algebras" (see [1, pp. 249, 300–301]) which are at the same time Heyting and Brouwer algebras without being equivalent to a Boolean algebras. One example of such algebras is the lattice of subgraphs of a given graph (see [17, pp. 29–31]). It is also possible to add in the upper part *locales* which are complete Heyting algebras and which constitute generalizations of the concept of topological space (see [8, pp. 472–475]). One example of *locale* is the lattice of open sets of a topological space (see [4]). These *locales* can pave the way to the *quantales* (see [13], [2]) quoted explicitly by Majid, and which are generalizations of *locales* well suited to formalize quantum theories. An example of such *quantales* is given by the lattice of ideals of a commutative ring with unity. The upper part of Majid's diagram is the natural place for the mathematics of sheaves. We can for instance build sheaves on a locale and the category of such sheaves is called a *localic topos* (see [8, pp. 472–527]), which generalizes the topos of sheaf on a usual topological space.

Using the "duality à la Majid", we can now come back to the lower part of the diagram. The dual of a *locale* is in fact a complete Brouwer algebra and this is what we have called a *globale*. A natural example of such a structure is the lattice of closed sets of a topological space. In fact properties of such lattices have been studied many years ago, in particular in the *Séminaire Dubreil* (see [7]). We can also consider the dual of a *quantale* and we find what we have called above a *universale*. The dual notion of a *sheaf on a locale* is now a *conciliation on a globale* and it would be interesting to study the category of such conciliations: what we might call a *globalic topos*. This topos is maybe a complement topos but this remains to be proved. It is also possible to think about the category of conciliations on a *globale* but up to now we don't know exactly what it is. Using the Majid's duality philosophy we could try to find self-dual structures that are at the same time *locale* and *globale* or at the same time *quantale* and *universale*. But we don't know if this is really relevant. We will try to clarify these problems in a forthcoming publication.



References

- [1] S. Awodey. *Category Theory*. Oxford University Press, 2nd edition, 2010.
- [2] M. Coniglio and F. Miraglia. Non-Commutative Topology and Quantaes. *Studia Logica*, Vol. 65, pp. 223–236, 2000.
- [3] N. Goodman. Logic of Contradiction. *Zeitschr. fur Math. Logik und Grundlagen der Math.*, Vol. 27, pp. 119–126, 1981.
- [4] C. Isham. *Modern Differential Geometry for Physicists*. World Scientific, New Jersey, 2nd edition, 2005.
- [5] W. James and C. Mortensen. Categories, Sheaves and Paraconsistent Logic. Unpublished manuscript.
- [6] F. Lawvere. Intrinsic Co-Heyting Boundaries and the Leibniz Rule in Certain Toposes. In *Category Theory*, Volume 1488 of *Springer Lecture Notes in Mathematics*. Proceeding Como 1990, Springer-Verlag, pp. 279–281, 1991.
- [7] L. Lesieur. Les treillis en topologie I–II. *Séminaire Dubreil. Algèbre et théorie des nombres*, Vol. 7, exp. n° 3, pp. 1–10 and exp. n°4, pp. 1–11, 1953–1954. Available on www.numdam.org.
- [8] S. Mac Lane and I. Moerdijk. *Sheaves in Geometry and Logic. A first Introduction to Topos Theory*. Springer, Berlin, 1992.
- [9] S. Majid. *Foundations of Quantum Group Theory*. Cambridge University Press, 1995.
- [10] S. Majid. Quantum Spacetime and Physical Reality. In S. Majid, editor, *On Space and Time*, pp. 56–140. Cambridge University Press, 2008.
- [11] J. McKinsey and A. Tarski. On Closed Elements in Closure Algebra. *Annals of Mathematics*, Vol. 47, pp. 122–162, 1946.

- [12] C. Mortensen. *Inconsistent Mathematics*. Kluwer Mathematics and Its Applications Series. Kluwer, Dordrecht, 1995.
- [13] C. Mulvey. “&”. *Suppl. Rend. Circ. Mat. Palermo Ser. II*, Vol. 12, pp. 99–104, 1986.
- [14] A. Neeman. *Algebraic and Analytic Geometry*. Cambridge University Press, 2007.
- [15] D. Perrin. *Géométrie analytique*. InterEditions / CNRS Editions, 1995.
- [16] H. Rasiowa. *An Algebraic Approach to Non-Classical Logics*. North-Holland/PWN-Polish Scientific Publishers, Amsterdam/Warszawa, 1974.
- [17] G. Reyes and H. Zolfaghari. Bi-Heyting Algebras, Toposes and Modalities. *Journal of Philosophical Logic*, Vol. 25, pp. 25–43, 1996.
- [18] G. Simmons. *Introduction to Topology and Modern Analysis*. McGraw-Hill, New York, 1963.
- [19] R. Ward and R. Wells Jr. *Twistor Geometry and Field Theory*. Cambridge University Press, 1990.

Dominique Lambert (d.lambert@fundp.ac.be)
Bertrand Hespel (bertrand.hespel@fundp.ac.be)

Département Sciences-Philosophies-Sociétés
Faculté des sciences
Centre ESPHIN et NAXYS
Université de Namur - FUNDP
61, rue de Bruxelles – B-5000 Namur (Belgium)