

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

Characterization of CYP26B1-selective inhibitor, DX314, as a potential therapeutic for keratinization disorders

Veit, Joachim; DE GLAS, VALERIE; Balau, Benoit; Liu, Haoming; Bourlond, Florence; Paller, Amy; Poumay, Yves; Diaz, Philippe

Published in:

Journal of Investigative Dermatology

DOI:

[10.1016/j.jid.2020.05.090](https://doi.org/10.1016/j.jid.2020.05.090)

Publication date:

2021

Document Version

Peer reviewed version

[Link to publication](#)

Citation for published version (HARVARD):

Veit, J, DE GLAS, VALERIE, Balau, B, Liu, H, Bourlond, F, Paller, A, Poumay, Y & Diaz, P 2021, 'Characterization of CYP26B1-selective inhibitor, DX314, as a potential therapeutic for keratinization disorders', *Journal of Investigative Dermatology*, vol. 141, pp. 72-83. <https://doi.org/10.1016/j.jid.2020.05.090>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



THE JOURNAL OF
INVESTIGATIVE
DERMATOLOGY

Characterization of CYP26B1-selective Inhibitor, DX314, as a Potential Therapeutic for Keratinization Disorders

Journal:	<i>Journal of Investigative Dermatology</i>
Manuscript ID	JID-2020-0105.R2
Article Type:	Original Article
Date Submitted by the Author:	29-Apr-2020
Complete List of Authors:	Veit, Joachim; University of Montana Missoula, Biomedical and Pharmaceutical Sciences De Glas, Valerie; University of Namur, URPHYM-NARLIS Balau, Benoît; University of Namur, URPHYM-NARLIS Liu, Haoming; Northwestern University Feinberg School of Medicine, Departments of Dermatology and Pediatrics Bourlond, Florence; Hôpital Erasme, Université Libre de Bruxelles, Service de Dermatologie Paller, Amy; Northwestern University Feinberg School of Medicine, Departments of Dermatology and Pediatrics Poumay, Yves; University of Namur, URPHYM-NARLIS Diaz, Philippe; University of Montana Missoula, Biomedical and Pharmaceutical Sciences; DermaXon LLC, R&D
Keywords:	Drug Development, Keratinization Disorders, Barrier Function, Pharmacology, Ichthyosis

SCHOLARONE™
Manuscripts

Editor comments:

Section/Deputy Editor: 1

Comments to the Author:

Revision is responsive to reviewer comments and improved by revision.

- We would like to thank the editors for their time and feedback regarding our submission.
- The only change we are submitting in this revision is to a minor figure reference issue we found in the Figure 4 legend. The reference to parts "(a,c,d)" in the last sentence was changed to correctly read "(a,b,d)". This correction was highlighted and underlined in yellow in the manuscript text.

Reviewer comments:**Reviewer: 1**

Comments to the Author


This is a very careful Revision of an excellent paper. My concern on the experimental design used, namely a simultaneous co-treatment with all trans retinoic acid in the keratinocyte cultures is well and fully addressed and authors Point out that in the rhino mice model no cotreatment was used, but similar effects were achieved. Moreover, they clarify that in their in vitro conditions no RA precursors and no appreciable levels of all trans retinoic acid will be present which justifies their Approach and they provide several prominent references like that of Giltaire et al 2009 on this culture model. All specific points have been fully and carefully addressed. No further comment.

Reviewer: 2

Comments to the Author

I believe that the authors have responded adequately to my questions.

- We would like to sincerely thank reviewer #1 and reviewer #2 for their time, effort and comments regarding our submission.



527x355mm (96 x 96 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Informative Title:

Characterization of CYP26B1-selective Inhibitor, DX314, as a Potential Therapeutic for Keratinization Disorders

Short Title:

DX314: Keratinization Disorder Therapeutic

Joachim G.S. Veit¹, Valérie De Glas², Benoît Balau², Haoming Liu⁴, Florence Bourlond⁵, Amy S. Paller⁴, Yves Poumay², Philippe Diaz^{1,3}

¹Department of Biomedical and Pharmaceutical Sciences, University of Montana, Missoula, MT

²URPHYM-NARLIS, University of Namur, Namur, Belgium

³DermaXon LLC, Missoula, MT

⁴Departments of Dermatology and Pediatrics, Northwestern University Feinberg School of Medicine, Chicago, IL

⁵Service de Dermatologie, Hôpital Erasme, Université Libre de Bruxelles, Belgique

Correspondence: Philippe Diaz, Department of Biomedical and Pharmaceutical Sciences, University of Montana, 32 Campus Dr., Missoula, MT, 59812. Email: philippe.diaz@umontana.edu

ORCIDs:

Joachim G.S. Veit: <https://orcid.org/0000-0001-5908-0648>

Valérie De Glas: <https://orcid.org/0000-0001-7366-2601>

Benoît Balau: <https://orcid.org/0000-0002-7314-860X>

Haoming Liu: <https://orcid.org/0000-0002-3190-5422>

Florence Bourlond: <https://orcid.org/0000-0003-1071-3306>

Amy S. Paller: <https://orcid.org/0000-0001-6187-6549>

Yves Poumay: <http://orcid.org/0000-0001-5200-3367>

Philippe Diaz: <https://orcid.org/0000-0002-2339-8803>

1
2
3 Abbreviations: RAMBA, retinoic acid metabolism blocking agents; *atRA*, all-*trans*-retinoic acid;
4 RHE, reconstructed human epidermis; TEER, transepithelial electrical resistance; TEWL,
5 transepidermal water loss; CYP26, cytochrome p450 family 26; DD, Darier disease; LI, lamellar
6 ichthyosis; RXLI, Recessive x-linked ichthyosis. See supplementary materials for additional
7 abbreviations (**Table S4**)
8
9
10
11
12

13 Funding: This work was supported by the by the National Institute of Arthritis and
14 Musculoskeletal and Skin Disease of the National Institutes of Health: R44AR069416 (PD),
15 P20GM103546 (PD, JV). The content of this paper is solely the responsibility of the authors and
16 does not necessarily reflect the official views of the National Institutes of Health.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Inhibition of cytochrome P450 (CYP)-mediated retinoic acid (RA) metabolism by RA metabolism blocking agents (RAMBAs) increases endogenous retinoids and is an alternative to retinoid therapy. Currently available RAMBAs (i.e. liarozole and talarozole) tend to have fewer adverse effects than traditional retinoids but lack target specificity. Substrate-based inhibitor DX314 has enhanced selectivity for RA-metabolizing enzyme CYP26B1 and may offer an improved treatment option for keratinization disorders such as congenital ichthyosis and Darier disease. In this study we use RT-qPCR, RNA sequencing, pathway, upstream regulator, and histological analyses to demonstrate that DX314 can potentiate the effects of all-*trans*-RA (*a*tRA) in healthy and diseased reconstructed human epidermis (RHE). We unexpectedly discovered that DX314, but not *a*tRA or previous RAMBAs, appears to protect epidermal barrier integrity. Additionally, DX314-induced keratinization and epidermal proliferation effects are observed in a rhino mice model. Altogether, results indicate that DX314 inhibits *a*tRA metabolism with minimal off-target activity and shows therapeutic similarity to topical retinoids *in vitro* and *in vivo*. Findings of a unique barrier-protecting effect require further mechanistic study but may lead to a novel strategy in barrier-reinforcing therapies. DX314 is a unique and promising candidate compound for further study and development in the context of keratinization disorders.

INTRODUCTION

Therapeutics targeting retinoid biopathways have been implemented in the clinical treatment of keratinization disorders such as the congenital ichthyoses (Vahlquist et al. 2008), Darier disease (DD) (Casals et al. 2009; Cooper and Burge 2003; Dicken et al. 1982; Steijlen et al. 1993), and other skin disorders (e.g. acne, psoriasis) (Dawson and Dellavalle 2013; Fisher and Voorhees 1996) to alleviate patient symptoms. Such therapies leverage the role of endogenous retinoids in regulating keratinocyte proliferation and differentiation. Retinoid bioactivity is primarily, although not solely, mediated by transcription factors such as retinoic acid receptors (RAR) and retinoid X receptors (RXR) (Fisher et al. 1994).

However, retinoid treatments also result in adverse effects including dry skin, irritation, redness, photosensitivity, teratogenicity and barrier impairment (Orfanos et al. 1997). Endogenously-occurring retinoids (e.g. tretinoin) autoinduce their own metabolism (Van Der Leede et al. 1997; Marikar et al. 1998), which in turn demands higher exogenous doses for effective treatment, resulting in increased systemic exposure. Synthetic retinoids, particularly tazarotenic acid, will both activate RARs and inhibit RA-metabolism (Foti et al. 2016), which may overstimulate retinoid biopathways and increase associated adverse effects. For these reasons, RA metabolism blocking agents (RAMBAs) (Verfaille et al. 2008), including liarozole and talarozole, were developed to target the primary RA-specific metabolizing enzymes of the cytochrome p450 family 26 (CYP26) (Ray et al. 1997).

RAMBAs, particularly when used topically, achieve therapeutic effects without high exposure to systemic levels of RA by utilizing endogenously available RA rather than high doses of exogenous retinoids, which theoretically reduces overexposure and adverse effects. Topical RAMBAs could be implemented either as standalone treatments or as adjunct therapies to reduce

1
2
3 oral retinoid dosing without loss of therapeutic efficacy. Reports illustrate successful inhibition
4 of RA metabolism using liarozole (Van Wauwe et al. 1992) and proven efficacy in treating
5 several skin disorders (Berth-Jones et al. 2000; Bhushan et al. 2001; Kang et al. 1996; Kuijpers
6 et al. 1998; Lucker et al. 2005; Lucker et al. 1997; Vahlquist et al. 2014), including in a
7 comparative trial vs acitretin (an oral retinoid) for congenital ichthyosis, which showed a trend
8 towards a better safety profile (Verfaillie et al. 2007b). Problematically, liarozole also inhibits
9 off-target CYPs such as aromatase (CYP19), an important enzyme in estradiol biosynthesis
10 (Nelson et al. 2013). Previous RAMBAs have not progressed past clinical trials, suggesting the
11 need for improved RAMBA candidates.
12
13
14
15
16
17
18
19
20
21
22
23

24 The promising results of early-generation RAMBAs led Diaz et al. to develop a series of
25 compounds targeting RA metabolism *via* specific inhibition of two CYP26 isoforms (A1 and B1)
26 (Diaz et al. 2016). Removal of the heme-interacting azole moiety, thought to contribute to the
27 non-specific effects of previous azole-containing RAMBAs, may preserve the desired effects
28 while minimizing off-target activity. One of these compounds, a CYP26B1-specific inhibitor,
29 DX314 (IC₅₀: CYP26A1=1752nM; CYP26B1=108nM), was described in US patent
30 US009963439B2 as example 39 (Diaz et al. 2018).
31
32
33
34
35
36
37
38
39

40 Endogenous all-*trans*-RA (*at*RA) is a well-known regulator of epidermal proliferation
41 and differentiation (Fisher and Voorhees 1996), in part by inducing heparin-binding EGF-like
42 growth factor (HBEGF), which stimulates keratinocyte proliferation (Rittié et al. 2006; Stoll and
43 Elder 1998; Xiao et al. 1999; Yoshimura et al. 2003), and involucrin (IVL), a late marker of
44 epidermal differentiation (Eckert et al. 2004; Monzon et al. 1996; Poumay et al. 1999).
45
46
47
48
49

50 Expression of both CYP26A1 and CYP26B1 is induced by *at*RA, but only the CYP26A1
51
52
53
54
55
56
57
58
59
60

1
2
3 promotor contains a RA response element (RARE) which directly binds RARs (Pavez Loriè et
4 al. 2009a).

5
6
7
8 In this study, we investigate RAMBAs *in vitro* under tightly-controlled growth conditions
9
10 that specifically do not contain RA or RA precursors (Giltaire et al. 2009; Minner et al. 2010;
11 Pavez Loriè et al. 2009a; Poumay et al. 1999). In these conditions, a highly specific RAMBA
12
13 will have a negligible effect in the absence of *at*RA, however when co-treated with a nanomolar
14
15 dose of *at*RA, facilitate a relative increase in *at*RA concentration by inhibiting its metabolism,
16
17 and therefore potentiate the expression of RA-responsive genes. Since *in vivo* concentrations of
18
19 *at*RA in healthy human skin are typically 2-4nM (Mihály et al. 2011), we co-treated RAMBAs *in*
20
21 *vitro* with 1nM *at*RA to provide a near-physiological basal level, without saturating nanomolar-
22
23 sensitive *at*RA effects, which allowed us to observe if RAMBAs can potentiate those effects.
24
25
26
27

28
29 Skin acts as an effective barrier to the environment. Changes in epidermal barrier
30
31 integrity can be investigated using transepithelial electrical resistance (TEER) and
32
33 transepidermal water loss (TEWL). TEER is an *in vitro* assay assessing electrical properties to
34
35 evaluate possible changes in trans- and paracellular (regulated by tight junctions) ion
36
37 permeability across the epidermis. TEWL measures passive diffusion of water across the
38
39 epidermis and can be performed *in vivo* or *in vitro*. Decreased TEER or increased TEWL
40
41 typically indicates barrier integrity disruption.
42
43

44
45 Our data show that DX314 potentiates the effects of *at*RA on gene expression in healthy
46
47 and diseased epidermis by inhibiting CYP26B1-mediated RA metabolism. Unexpectedly,
48
49 DX314 mitigated the epidermal barrier dysregulation and irregular morphology displayed by
50
51 other RAMBAs and high dose *at*RA. In addition, topical DX314 induced comedolytic/anti-
52
53
54
55
56
57
58
59
60

1
2
3 keratinizing effects in the rhino mouse model, reflecting those observed in previous retinoid
4 studies (Ashton et al. 1984; Fort-Lacoste et al. 1999; Kligman and Kligman 1979).
5
6

7
8 Together, this study suggests that DX314 exhibits potential as a new keratinization
9 disorder therapeutic that may address some shortcomings of previous retinoid-based treatments.
10
11

12 13 14 **RESULTS**

15 **DX314 potentiates the effects of *at*RA in healthy and keratinization disorder keratinocytes**

16
17 Based on RT-qPCR assays, a 4-day *at*RA treatment of RHE caused a dose-dependent
18 increase in HBEGF, CYP26A1 and IVL gene expression (**Figure 1a**). DX314, together with
19 near-physiological dosing of *at*RA (1nM; to provide basal RA without saturating sensitive RA
20 pathways) mimics the consequences of high dose *at*RA on the expression of every gene
21 analyzed, indicating potentiation of *at*RA. Liarozole with *at*RA significantly increases HBEGF
22 and CYP26A1 expression relative to control, but only CYP26A1 expression is potentiated
23 compared to 1nM *at*RA alone.
24
25
26
27
28
29
30
31
32
33
34

35 Immunostaining shows IVL (**Figure 1b**) primarily localized in the upper epidermis of
36 control and DX314-alone RHE. Induction of early IVL expression is observed in basal and
37 suprabasal layers of the epidermis with *at*RA-alone, and even more so, DX314 with 1nM *at*RA.
38
39
40
41

42 RNAseq confirmed that DX314 alone (*at*RA-free conditions) had no effect on HBEGF,
43 IVL, CYP26A1 or CYP26B1 mRNA expression (**Figure 1c**), and that DX314 potentiated the
44 effects of 1nM *at*RA, not only on these genes, but on numerous other known retinoid-responsive
45 genes, among them several keratins (KRT) (Radoja et al. 1997), lecithin:retinol acyl-transferase
46 (LRAT) (Kurlandsky et al. 1996), retinol binding protein 1 (RBP1) (Kang et al. 1995), and
47 cellular retinoic acid binding protein 2 (CRABP2) (Aström et al. 1994). Changes in expression of
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 other genes that involve both direct (RARE-containing promotor, indicated by adjacent green
4 box) (Aström et al. 1992; Fisher et al. 1995; Lalevée et al. 2011; Laursen et al. 2015; Loudig et
5 al. 2000; Radoja et al. 1997; de Thé et al. 1990; Tomic-Canic et al. 1992; Vasios et al. 1989) and
6 indirect or unknown RA pathways supported the potentiation effect.
7
8
9
10

11
12 RNAseq data was further analyzed with Ingenuity Pathway Analysis (IPA) (Krämer et al.
13 2014) software for canonical pathway and upstream regulator prediction. The applied
14 significance cutoffs (see **Methods**) resulted in 1360, 5480, 169, and 3015 differentially
15 expressed genes in RHE treated with 1nM *atRA*, 100nM *atRA*, 1000nM DX314, and 1000nM
16 DX314 with 1nM *atRA*, respectively. Analysis results were sorted by overall |z-score|, indicating
17 the strength and direction of each prediction (positive score = activation, negative score =
18 inhibition). As expected, the upstream regulator with largest activation score was *atRA*
19 (tretinoin), which displayed an activation pattern consistent with *atRA* potentiation (**Figure 1d**).
20 Of the top 20 scoring regulators, only three weakly displayed any predicted activity by DX314
21 alone. Canonical pathway analysis (**Figure 1e**) found the overall most activated (“Integrin
22 Signaling”) and inhibited (“RhoGDI Signaling”) pathways both display activation patterns
23 suggesting a potentiation of *atRA* by DX314.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40 To test the potential of DX314 in certain keratinization disorders, we investigated
41 CYP26A1 gene expression in keratinocytes from patients with Darier disease (DD), recessive x-
42 linked ichthyosis (RXLI), and lamellar ichthyosis (LI). DX314 potentiates the effects of *atRA* in
43 DD RHE (**Figure 2a**), RXLI full-thickness RHE (**Figure 2b**), LI RHE (**Figure 2c**), and RXLI
44 monolayer cultures (**Figure 2d**). Talarozole potentiated *atRA* in DD RHE and liarozole
45 potentiated *atRA* in RXLI full-thickness RHE.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Histological analysis shows that *at*RA induces robust morphological changes in DD RHE
4 (**Figure 3a**) including a dramatic loss of SG (and their filaggrin-containing keratohyalin granules
5 (KG)), denucleation and flattening of stratum spinosum (SS) keratinocytes, and an overall
6 unhealthy appearance. When treated alone, DX314 and talarozole caused no major
7 morphological changes. Co-treatment with talarozole and 1nM *at*RA shifted the cell
8 morphology, most notably with loss of SG and flattening of epidermal keratinocytes, to more
9 closely resemble the appearance of high dose *at*RA. DX314 with *at*RA also affected morphology
10 relative to DX314 alone, but to a lesser extent, with the appearance not significantly different
11 than control or only 1nM *at*RA-treated RHE.
12
13
14
15
16
17
18
19
20
21
22
23

24 KRT10, a commonly used marker of epidermal differentiation that localizes to the
25 suprabasal epidermis, showed reduced expression in *at*RA-treated RHE (**Figure 3b**). Treatment
26 with DX314 or talarozole alone led to no change in KRT10 localization, but co-treatment with
27 near-physiological levels of *at*RA reduced staining in the lower SS. KRT10 gene expression in
28 DD RHE (**Figure 3c**) was decreased by *at*RA in a dose-dependent manner and was potentiated
29 by DX314 and talarozole. This effect on KRT10 was also observed in healthy RHE (**Figure 1c**
30 and **S1**).

31 **DX314 induces barrier effects in healthy and diseased RHE**

32
33
34
35
36
37
38
39
40
41
42 As a measure of barrier integrity, TEER of healthy RHE was assessed. Independent runs
43 with RHE from several donors were pooled and normalized to their respective controls (**Figure**
44 **4a**). High dose *at*RA significantly decreased TEER. Liarozole and talarozole alone showed no
45 effect on TEER, but co-treatment with *at*RA resulted in significant TEER decrease. Surprisingly,
46 in both healthy and LI (**Figure 4b**) RHE, DX314 alone increased TEER and caused no decrease
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 relative to control with *at*RA co-treatment. TEWL was similarly affected by *at*RA, but no
4
5 significant change was seen with DX314 alone.
6

7
8 Morphologically, *at*RA disrupted LI (**Figure 4c**) and healthy RHE (**Figure S2 and S3**)
9
10 structure as described above. DX314 led to no major changes in morphology when dosed alone.
11
12 However, unlike other RAMBAs (**Figure S2**), DX314 with *at*RA co-treatment reduced
13
14 disruption in normal morphology (improved SG/KG, more columnar basal keratinocytes, and
15
16 less disorganized upper epidermis). Semi-quantitative analysis of SG surface area (**Figure 4d**),
17
18 measured in healthy RHE (**Figure S3**), confirmed a dramatic reduction of the SG by *at*RA, but
19
20 no significant loss in DX314 treated groups.
21
22

23
24 The epidermal differentiation complex (EDC) is a cluster of genes on human locus 1q21
25
26 that are essential for epidermal differentiation (Kypriotou et al. 2012). As noted using RNAseq,
27
28 expression of these genes (**Figure 4e**) is generally consistent with other retinoid-responsive
29
30 genes. However, many cornified envelope (CE) precursor family genes, such as late cornified
31
32 envelope (LCE) and small proline rich (SPRR) proteins, were dramatically downregulated by
33
34 high dose *at*RA, but not affected by DX314 with *at*RA, which may play a role in the observed
35
36 barrier effects. In addition, FLG expression was increased 2-fold with the DX314-*at*RA
37
38 combination, but not with high dose *at*RA.
39
40
41

42
43 Nuclear receptor profiling revealed that DX314 acts as an inverse agonist for RAR-
44
45 related orphan receptors (ROR) α and γ (**Figure S4**), while showing no activity on any other
46
47 nuclear receptors studied.
48

49 **DX314 reduces epidermal abnormalities in rhino mice**

50
51 Rhino mice are commonly used as an *in vivo* model for screening comedolytic and anti-
52
53 keratinizing compounds such as retinoids (Ashton et al. 1984; Fort-Lacoste et al. 1999; Griffiths
54
55
56
57
58
59
60

1
2
3 et al. 1993; Seiberg et al. 1997). Overall, DX314 treatment improved skin morphology (**Figure**
4 **5**). DX314 decreased comedo density (**Figure 6a** and **6b**), increased the mean comedo profile
5
6 (ratio of comedo opening size to internal diameter) (**Figure 6c**), suggesting comedolysis, and
7
8 induced epidermal thickening (**Figure 6d**), consistent with previous studies of topical retinoids.
9
10 Again, DX314 treatment did not change TEWL relative to vehicle (**Figure 6e**). No abnormal
11
12 behavior, adverse skin changes, changes in body weight or DRAIZE scoring (**Table S5** and **S6**)
13
14 were observed throughout the study.
15
16
17
18
19
20
21

22 **DISCUSSION**

23
24 Retinoid-based drugs are well-accepted therapeutics for the treatment of many skin
25
26 diseases (Dawson and Dellavalle 2013; Fisher and Voorhees 1996; Vahlquist et al. 2008).
27
28 Despite their efficacy, use often leads to adverse reactions from their wide spectrum of non-
29
30 therapeutically relevant endogenous roles, which are exacerbated by metabolic autoinduction and
31
32 tolerance (Digiovanna et al. 2013; Orfanos et al. 1997). A strategy involving RAMBAs showed
33
34 potential in preclinical and clinical studies (Berth-Jones et al. 2000; Bhushan et al. 2001;
35
36 Bovenschen et al. 2007; Giltaire et al. 2009; Kang et al. 1996; Kuijpers et al. 1998; Lucker et al.
37
38 2005; Lucker et al. 1997; Pavez Loriè et al. 2009b; Stoppie et al. 2000; Vahlquist et al. 2014;
39
40 Verfaille et al. 2007a; Van Wauwe et al. 1992), however, first-generation RAMBAs have not
41
42 progressed to approved for clinical use. A highly selective RAMBA, with low risk of adverse
43
44 events, could address the downsides of current treatment options. This study investigates the
45
46 CYP26B1-selective compound, DX314, as a potential next-generation RAMBA.
47
48
49
50

51 Potentiation of the effects of a low, physiologically relevant dose of *α*RA by DX314 in
52
53 healthy and keratinization disorder keratinocytes, but not DX314 in an *α*RA-free environment,
54
55
56
57
58
59
60

1
2
3 confirms that DX314 acts by inhibiting *α*RA metabolism. These gene expression patterns were
4 reproduced in keratinocyte cultures from individuals with DD and congenital ichthyosis, in
5 addition to healthy skin, suggesting that the bioactivity of DX314 can be therapeutically relevant
6 in skin disorders.
7
8
9
10
11

12 A broader investigation of gene expression changes using RNAseq also showed a strong
13 pattern indicating potentiation of *α*RA by DX314 on both RARE-promoted, and indirectly
14 regulated genes. Pathway analysis found compelling supporting evidence in predicted upstream
15 regulator and canonical pathway activation patterns.
16
17
18
19
20

21 Immunostaining confirmed the *α*RA potentiating effects of DX314 on IVL localization
22 in healthy RHE and KRT10 localization in DD RHE.
23
24
25

26 These experiments showed that DX314 alone had minimal effect on gene expression and
27 therefore, minimal potential for off-target adverse effects, despite therapeutically relevant effects
28 when paired with endogenous levels of *α*RA.
29
30
31
32

33 Keratinization disorders are associated with intrinsic epidermal barrier disruption and a
34 therapy that improves barrier function would be highly desirable. Surprisingly, this study found a
35 significant increase in TEER compared to controls in RHE treated with DX314 alone, and unlike
36 with liarozole and talarozole, no decrease from control when co-treated with *α*RA. Although
37 higher doses of *α*RA significantly decreased TEER and DX314 otherwise appeared to potentiate
38 the effects of *α*RA, DX314 with *α*RA did not impair barrier function below that of the control.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TEWL in LI RHE, which is expected to increase with retinoid treatment or ablation of CYP26B1 (Okano et al. 2012), was not increased by DX314 alone or beyond that of 1nM *α*RA when added as co-treatment. DX314 did not decrease TEWL in RHE, however, a lack of correlation between *in vitro* TEWL and barrier function has been previously documented (Chilcott et al. 2002).

1
2
3 Despite previous studies in rhino mice showing retinoids inducing substantial increases in TEWL
4 (Elias et al. 1981; Gendimenico et al. 1994), our study found a slight, albeit a non-significant,
5 decrease in TEWL, which additionally suggests an *in vivo* barrier-protecting effect.
6
7

8
9
10 Morphologically, DX314-treated RHE show dramatically less disruption of the SG and
11 KG, and display an overall healthy appearance compared to *at*RA, liarozole or talarozole-treated
12 healthy, DD, and LI RHE. Semi-quantitative analysis of SG surface area also showed a
13 significant decrease in *at*RA-treated RHE, but no significant change in RHE treated with DX314,
14 with or without *at*RA. Nevertheless, DX314-treated RHE tissue sections had a more continuous
15 SG/KG layer relative to controls, which may translate to an increase in TEER.
16
17

18
19 We observed minimal changes in the expression of many cornified envelope precursor
20 proteins within the EDC, despite a large decrease in expression by high dose *at*RA and DX314
21 potentiating of the effects of *at*RA on other retinoid-responsive genes. We speculate that this
22 preservation of CE protein expression may contribute to the ameliorative effect of DX314 on
23 barrier integrity. In addition, DX314 displays unique inverse agonist activity on ROR α and γ .
24 Affinity for ROR γ may be explained by DX314's structural similarity to *at*RA, which has been
25 previously shown to bind to, and inhibit, ROR γ (Stehlin-Gaon et al. 2003). Conversely, *at*RA
26 was not found to act on ROR α , so the DX314 inverse agonism represents another unique
27 property. Furthermore, a previously studied topical ROR α/γ inverse agonist was found to inhibit
28 inflammation in mouse models of atopic dermatitis (Dai et al. 2017), a skin disorder displaying
29 pathological barrier disruption. Future investigations should explore the potential link between
30 DX314's ROR α/γ activity and its barrier effects, as well as potential contribution to therapeutic
31 effects of skin barrier protection.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Use of rhino mice to study the *in vivo* effects of dermatologically active compounds such
4 as retinoids is well-established, and unlike *in vitro*, does not require co-treatment with *atRA*
5 since adequate RA is generated *in vivo* through dietary sources. Reduction of the acne-like cysts
6 (comedones), as well as the associated epidermal thickening (hyperplasia), are sensitive to
7 retinoid treatment. In this preliminary study, DX314 led to significant improvement overall in
8 comedo number (comedones per cm of skin), profile, and induced epidermal thickening.
9
10 Optimization of the DX314 formulation (to eliminate the harsh acetone vehicle), dosing to
11 improve bioavailability, and extending the treatment duration are likely to amplify DX314
12 efficacy in this model.
13
14
15
16
17
18
19
20
21
22
23

24 We conclude that our results provide strong evidence that DX314, which is known to
25 specifically inhibit the RA-metabolizing enzyme CYP26B1, potentiates the effects of
26 physiological levels of *atRA* in keratinocytes from healthy skin and keratinization disorders *in*
27 *vitro*; may protect from epidermal skin barrier disruption by retinoids; and has a restorative effect
28 on changes *in vivo* rhino mouse skin consistent with previous retinoid treatments. These
29 observations merit further investigation as a unique keratinization disorder treatment with the
30 ability to simultaneously correct abnormal keratinization while protecting critical skin barrier
31 function. Together these findings present an exciting new therapeutic candidate aimed at
32 providing improved patient outcomes with minimal adverse effects, in contrast to currently
33 available treatments.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

MATERIALS AND METHODS

Primary keratinocytes

Healthy and DD primary keratinocytes, provided by Dr. Poumay's lab (Namur, Belgium), were isolated as previously described (Poumay et al. 2004) from skin samples provided by Drs. B. Bienfait and J.S. Blairvacq (Clinique St Luc, Namur-Bouge, Belgium). Additional healthy keratinocytes were purchased from ThermoFisher (Cascade Biologics, Portland, OR). RXLI and LI keratinocytes were provided by Dr. Paller (Northwestern University, IL). Details in **Table S1**.

Monolayer culture and RHE

Monolayer cultures were prepared as previously described (Minner et al. 2010). Upon reaching confluence, keratinocytes were treated for 20hr to compounds solubilized in media (0.1% DMSO vehicle for all *in vitro* studies).

RHE were produced as previously described (Poumay et al. 2004; De Vuyst et al. 2014) in Epilife media with 1.5mM Ca²⁺ (Cascade Biologics, Portland, OR), 10ng/mL keratinocyte growth factor (Sigma, Saint Louis, MO) and 50μg/mL vitamin C (Sigma, Saint Louis, MO). Treatments were started day 7 of growth, refreshed day 9, and halted day 11.

Full-thickness RHE were prepared as previously described (Zheng et al. 2012). Briefly, keratinocytes were seeded atop a simulated dermis (collagen matrix containing J2-3T3 fibroblasts) and allowed to develop into stratified epidermis before receiving a 4-day treatment (refreshed day 2).

RHE histological analysis and immunostaining

RHE were processed as previously described (Frankart et al. 2012; De Vuyst et al. 2014) and stained with hematoxylin-eosin (HE) or prepared for immunostaining. Further described in

Supplementary Methods.

Measures of epidermal barrier function

TEER was measured by a previously described method (Frankart et al. 2012) with a ERS-2 voltohmmeter (Millipore, Burlington, MA). TEWL measurement used an AquaFlux AF200 evaporimeter (Biox Systems, London, England). For RHE, a sterilized gasket was placed between the cell culture insert and TEWL probe to form airtight seal. TEWL was measured over 60-90s until reaching a steady state. Analysis of SG surface area was performed using Fiji/ImageJ (Schindelin et al. 2012). SG surface area, defined by the presence of KG, was manually outlined and the area divided by each tissue's total area.

RNA isolation, RT-qPCR, RNAseq, and bioinformatics

Details on RT-qPCR, RNA-seq, and bioinformatics (Andrews 2010; Bolger et al. 2014; Durinck et al. 2009; Kim et al. 2017; Kim et al. 2016; Kim et al. 2015; Li et al. 2009; Love et al. 2013; Pertea et al. 2015; RCoreTeam 2018; Zhu et al. 2018) are provided in **Supplementary Methods**. Primer sequences (Giltaire et al. 2009) provided in **Table S2**.

When applicable, a method described by (Willems et al. 2008) was used to standardize the qPCR data to correct for interindividual variability before analysis.

Nuclear receptor profiling

Refer to **Supplementary Methods**.

Rhino mice

Eleven RHJ/LeJ rhino mice (2-3 males, 3 females per group) received daily topical application of 50 μ L vehicle (acetone), or 1% DX314, on a 2x2cm area of back skin for 11 days.

All animal studies were approved by IACUC under NIH guidelines. Details in **Supplementary**

Methods.

Statistical analysis

Statistics, apart from separately described RNAseq portion, were performed as described in respective figure legends using Prism 6 (Graphpad Software, La Jolla, CA).

Data availability statement

Data available upon request.

CONFLICTS OF INTEREST

PD is cofounder of DermaXon™ and inventor of the technology, he and The University of Montana are entitled to future royalty payments. JV was employed at DermaXon™ during a portion of this study.

ACKNOWLEDGMENTS

We would like to thank Drs. B. Bienfait and JS. Blairvacq (Clinique St. Luc, Namur-Bouge, Belgium) for providing skin samples used in this study. We would also like to thank Joanna Kreitinger and Larissa Walker of DermaXon™ and the Northwestern University Skin Biology and Diseases Resource-based Center (P30AR075049) for their assistance with various lab tasks.

AUTHOR CONTRIBUTIONS

Conceptualization: *lead-* PD; *equal-* JV, YP, AP
Data Curation: *lead-* JV
Formal Analysis: *lead-* JV
Funding Acquisition: *lead-* PD; *equal-* AP; *supportive-* JV
Investigation: *lead-* JV; *supportive-* VDG, BB, HL
Methodology: *equal-* JV, YP, PD; *supportive-* AP
Project Administration: *lead-* PD; *supportive-* JV, AP, YP
Resources: *equal-* JV, YP, PD; *supportive-* AP, VDG, BB, HL
Software: *lead-* JV
Supervision: *lead-* PD; *supportive-* AP, YP
Validation: *equal-* JV, YP, PD; *supportive-* FB
Visualization: *lead-* JV
Writing – Original Draft Preparation: *lead-* JV
Writing – Review and Editing: *equal-* JV, AP, YP, PD

REFERENCES

- Andrews S. FastQC: A quality control tool for high throughput sequence data. [Internet]. 2010. Available from: <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>
- Ashton RE, Connor MJ, Lowe NJ. Histologic changes in the skin of the rhino mouse (hrrhrrh) induced by retinoids. *J. Invest. Dermatol.* Elsevier Masson SAS; 1984;82(6):632–5
- Aström A, Pettersson U, Chambon P, Voorhees JJ. Retinoic acid induction of human cellular retinoic acid-binding protein-II gene transcription is mediated by retinoic acid receptor-retinoid X receptor heterodimers bound to one far upstream retinoic acid-responsive element with 5-base pair spacing. *J. Biol. Chem.* 1994;269(35):22334–9
- Aström A, Pettersson U, Voorhees JJ. Structure of the human cellular retinoic acid-binding protein II gene. Early transcriptional regulation by retinoic acid. *J. Biol. Chem.* 1992;267(35):25251–5
- Berth-Jones J, Todd G, Hutchinson PE, Thestrup-Pedersen K, Vanhoutte FP. Treatment of psoriasis with oral liarozole: a dose-ranging study. *Br. J. Dermatol.* 2000;143(6):1170–6
- Bhushan M, Burden AD, McElhone K, James R, Vanhoutte FP, Griffiths CEM. Oral liarozole in the treatment of palmoplantar pustular psoriasis: A randomized, double-blind, placebo-controlled study. *Br. J. Dermatol.* 2001;145(4):546–53
- Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics.* 2014;30(15):2114–20
- Bovenschen HJ, Otero ME, Langewouters AMG, van Vlijmen-Willems IMJJ, van Rens DWA, Seyger MMB, et al. Oral retinoic acid metabolism blocking agent Rambazole for plaque psoriasis: an immunohistochemical study. *Br. J. Dermatol.* 2007;156(2):263–70
- Casals M, Campoy A, Aspiólea F, Carrasco MA, Camps A. Successful treatment of linear Darier's disease with topical adapalene. *J. Eur. Acad. Dermatol. Venereol.* 2009;23(2):237–8
- Chilcott RP, Dalton CH, Emmanuel AJ, Allen CE, Bradley ST. Transepidermal water loss does not correlate with skin barrier function in vitro. *J. Invest. Dermatol.* Elsevier Masson SAS; 2002;118(5):871–5 Available from: <http://dx.doi.org/10.1046/j.1523-1747.2002.01760.x>
- Cooper SM, Burge SM. Darier's disease: epidemiology, pathophysiology, and management. *Am. J. Clin. Dermatol.* 2003;4(2):97–105
- Dai J, Choo M-K, Park JM, Fisher DE. Topical ROR Inverse Agonists Suppress Inflammation in Mouse Models of Atopic Dermatitis and Acute Irritant Dermatitis. *J. Invest. Dermatol.* 2017;137(12):2523–31
- Dawson AL, Dellavalle RP. Acne vulgaris. *BMJ.* 2013;346(6):f2634
- Diaz P, Huang W, Keyari CM, Buttrick B, Price L, Guilloteau N, et al. Development and Characterization of Novel and Selective Inhibitors of Cytochrome P450 CYP26A1, the Human Liver Retinoic Acid Hydroxylase. *J. Med. Chem.* 2016;59(6):2579–95
- Diaz P, Isoherranen N, Buttrick B, Guilloteau N. Specific Inhibitors of Cytochrome P450 26 Retinoic Acid Hydroxylase. United States: United States Patent Office; 2018.
- Dicken CH, Bauer EA, Hazen PG, Krueger GG, Marks JG, McGuire JS, et al. Isotretinoin

- 1
2
3 treatment of Darier's disease. *J. Am. Acad. Dermatol.* 1982;6(4):721–6
4
5 Digiovanna JJ, Mauro T, Milstone LM, Schmuth M, Toro JR. Systemic retinoids in the
6 management of ichthyoses and related skin types. *Dermatol. Ther.* 2013;26(1):26–38
7
8 Durinck S, Spellman PT, Birney E, Huber W. Mapping identifiers for the integration of genomic
9 datasets with the R/Bioconductor package biomaRt. *Nat. Protoc.* 2009;4(8):1184–91
10
11 Eckert RL, Crish JF, Efimova T, Dashti SR, Deucher A, Bone F, et al. Regulation of involucrin
12 gene expression. *J. Invest. Dermatol.* Elsevier Masson SAS; 2004;123(1):13–22 Available from:
13 <http://dx.doi.org/10.1111/j.0022-202X.2004.22723.x>
14
15 Elias PM, Fritsch PO, Lampe M, Williams ML, Brown BE, Nemanic M, et al. Retinoid effects
16 on epidermal structure, differentiation, and permeability. *Lab. Investig.* 1981;44(6):531–40
17
18 Fisher GJ, Reddy AP, Datta SC, Kang S, Yi JY, Chambon P, et al. All-trans retinoic acid induces
19 cellular retinol-binding protein in human skin in vivo. *J. Invest. Dermatol.* 1995;105(1):80–6
20
21 Fisher GJ, Talwar HS, Xiao JH, Datta SC, Reddy AP, Gaub MP, et al. Immunological
22 identification and functional quantitation of retinoic acid and retinoid X receptor proteins in
23 human skin. *J. Biol. Chem.* 1994;269(32):20629–35
24
25 Fisher GJ, Voorhees JJ. Molecular mechanisms of retinoid actions in skin. *Faseb J.*
26 1996;10(9):1002–13
27
28 Fort-Lacoste L, Verscheure Y, Tisne-Versailles J, Navarro R. Comedolytic effect of topical
29 retinaldehyde in the rhino mouse model. *Dermatology.* 1999;199(SUPPL. 1):33–5
30
31 Foti RS, Isoherranen N, Zelter A, Dickmann LJ, Buttrick BR, Diaz P, et al. Identification of
32 Tazarotenic Acid as the First Xenobiotic Substrate of Human Retinoic Acid Hydroxylase
33 CYP26A1 and CYP26B1. *J. Pharmacol. Exp. Ther.* 2016;357(2):281–92
34
35 Frankart A, Malaisse J, De Vuyst E, Minner F, de Rouvroit CL, Poumay Y. Epidermal
36 morphogenesis during progressive in vitro 3D reconstruction at the air-liquid interface. *Exp.*
37 *Dermatol.* 2012;21(11):871–5
38
39 Gendimenico GJ, Stim TB, Corbo M, Mezick JA, Janssen B. A Pleiotropic Response Is Induced
40 in F9 Embryonal Carcinoma Cells and Rhino Mouse Skin by All-trans-Retinoic Acid, a RAR
41 Agonist but Not by SR11237, a RXR-Selective Agonist. *J. Invest. Dermatol.* Elsevier Masson
42 SAS; 1994;102(5):676–80
43
44 Giltaire S, Herphelin F, Frankart A, Hérin M, Stoppie P, Poumay Y. The CYP26 inhibitor
45 R115866 potentiates the effects of all-trans retinoic acid on cultured human epidermal
46 keratinocytes. *Br. J. Dermatol.* 2009;160(3):505–13
47
48 Griffiths CE, Elder JT, Bernard BA, Rossio P, Cromie MA, Finkel LJ, et al. Comparison of
49 CD271 (adapalene) and all-trans retinoic acid in human skin: dissociation of epidermal effects
50 and CRABP-II mRNA expression. *J. Invest. Dermatol.* 1993. p. 325–8
51
52 Kang S, Duell EA, Fisher GJ, Datta SC, Wang ZQ, Reddy AP, et al. Application of retinol to
53 human skin in vivo induces epidermal hyperplasia and cellular retinoid binding proteins
54 characteristic of retinoic acid but without measurable retinoic acid levels or irritation. *J. Invest.*
55 *Dermatol.* Elsevier Masson SAS; 1995;105(4):549–56
56
57 Kang S, Duell EA, Kim KJ, Voorhees JJ. Liarozole inhibits human epidermal retinoic acid 4-

- 1
2
3 hydroxylase activity and differentially augments human skin responses to retinoic acid and
4 retinol in vivo. *J. Invest. Dermatol.* 1996;107(2):183–7
5
6 Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory
7 requirements. *Nat. Methods.* 2015;12(4):357–60
8
9 Kim B, Langmead B, Salzberg S. genome_tran [Internet]. 2017 [cited 2018 Jun 25]. Available
10 from: ftp://ftp.ccb.jhu.edu/pub/infphilo/hisat2/data/grch38_tran.tar.gz
11
12 Kim D, Pertea M, Kim D, Pertea GM, Leek JT, Salzberg SL. Transcript-level expression
13 analysis of RNA-seq experiments with HISAT , StringTie and Transcript-level expression
14 analysis of RNA-seq experiments with HISAT , StringTie and Ballgown. *Nat. Protoc. Nature*
15 *Publishing Group;* 2016;11(9):1650–67
16
17 Kligman LH, Kligman AM. The effect on rhino mouse skin of agents which influence
18 keratinization and exfoliation. *J. Invest. Dermatol. Elsevier Masson SAS;* 1979;73(5 I):354–8
19
20 Krämer A, Green J, Pollard J, Tugendreich S. Causal analysis approaches in ingenuity pathway
21 analysis. *Bioinformatics.* 2014;30(4):523–30
22
23 Kuijpers ALA, Van Pelt JPA, Bergers M, Boegheim PJ, Den Bakker JEN, Siegenthaler G, et al.
24 The effects of oral liarozole on epidermal proliferation and differentiation in severe plaque
25 psoriasis are comparable with those of acitretin. *Br. J. Dermatol.* 1998;139(3):380–9
26
27 Kurlandsky SB, Duell EA, Kang S, Voorhees JJ, Fisher GJ. Auto-regulation of retinoic acid
28 biosynthesis through regulation of retinol esterification in human keratinocytes. *J. Biol. Chem.*
29 1996;271(26):15346–52
30
31 Kypriotou M, Huber M, Hohl D. The human epidermal differentiation complex: Cornified
32 envelope precursors, S100 proteins and the “fused genes” family. *Exp. Dermatol.*
33 2012;21(9):643–9
34
35 Lalevée S, Anno YN, Chatagnon A, Samarut E, Poch O, Laudet V, et al. Genome-wide in silico
36 identification of new conserved and functional retinoic acid receptor response elements (direct
37 repeats separated by 5 bp). *J. Biol. Chem.* 2011;286(38):33322–34
38
39 Laursen KB, Kashyap V, Scandura J, Gudas LJ. An alternative retinoic acid-responsive Stra6
40 promoter regulated in response to retinol deficiency. *J. Biol. Chem.* 2015;290(7):4356–66
41
42 Van Der Leede BJM, Van Den Brink CE, Pijnappel WWM, Sonneveld E, Van Der Saag PT,
43 Van Der Burg B. Autoinduction of retinoic acid metabolism to polar derivatives with decreased
44 biological activity in retinoic acid-sensitive, but not in retinoic acid-resistant human breast
45 cancer cells. *J. Biol. Chem.* 1997;272(29):17921–8
46
47 Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
48 Alignment/Map format and SAMtools. *Bioinformatics.* 2009;25(16):2078–9
49
50 Loudig O, Babichuk C, White J, Abu-Abed S, Mueller C, Petkovich M. Cytochrome
51 P450RAI(CYP26) promoter: a distinct composite retinoic acid response element underlies the
52 complex regulation of retinoic acid metabolism. *Mol. Endocrinol.* 2000;14(9):1483–97
53
54 Love M, Anders S, Huber W. Differential analysis of RNA-Seq data at the gene level using the
55 DESeq package. 2013;1–32
56
57 Lucker GPH, Heremans AMC, Boegheim PJ, Van De Kerkhof PCM, Steijlen PM. Oral

- 1
2
3 treatment of ichthyosis by the cytochrome P-450 inhibitor liarozole. *Br. J. Dermatol.*
4 1997;136(1):71–5
5
- 6 Lucker GPH, Verfaillie CJ, Heremans AMC, Vanhoutte FP, Boegheim JPJ, Steijlen PPM.
7 Topical liarozole in ichthyosis: A double-blind, left-right comparative study followed by a long-
8 term open maintenance study [2]. *Br. J. Dermatol.* 2005;152(3):566–9
9
- 10 Marikar Y, Wang Z, Petkovich M, Voorhees JJ, Fisher GJ, Duell EA. Retinoic Acid Receptors
11 Regulate Expression of Retinoic Acid 4-Hydroxylase that Specifically Inactivates All-Trans
12 Retinoic Acid in Human Keratinocyte HaCaT Cells. *J. Invest. Dermatol. Elsevier Masson SAS;*
13 1998;111(3):434–9
14
- 15 Mihály J, Gamlieli A, Worm M, Rühl R. Decreased retinoid concentration and retinoid
16 signalling pathways in human atopic dermatitis. *Exp. Dermatol.* 2011;20(4):326–30
17
- 18 Minner F, Herphelin F, Poumay Y. Study of Epidermal Differentiation in Human Keratinocytes
19 Cultured in Autocrine Conditions. *Biol. Integument.* 2010. p. 71–82
20
- 21 Monzon RI, LaPres JJ, Hudson LG. Regulation of involucrin gene expression by retinoic acid
22 and glucocorticoids. *Cell Growth Differ.* 1996;7(12):1751–9
23
- 24 Nelson CH, Buttrick BR, Isoherranen N. Therapeutic potential of the inhibition of the retinoic
25 acid hydroxylases CYP26A1 and CYP26B1 by xenobiotics. *Curr. Top. Med. Chem. NIH Public*
26 *Access;* 2013;13(12):1402–28
27
- 28 Okano J, Lichti U, Mamiya S, Aronova M, Zhang G, Yuspa S, et al. Increased retinoic acid
29 levels through ablation of Cyp26b1 determine the processes of embryonic skin barrier formation
30 and peridermal development. *J. Cell Sci.* 2012;125(Pt 7):1827–36
31
- 32 Orfanos CE, Zouboulis CC, Almond-Roesler B, Geilen CC. Current use and future potential role
33 of retinoids in dermatology. *Drugs.* 1997;53(3):358–88
34
- 35 Pavez Loriè E, Chamcheu JC, Vahlquist A, Törmä H. Both all-trans retinoic acid and
36 cytochrome P450 (CYP26) inhibitors affect the expression of vitamin A metabolizing enzymes
37 and retinoid biomarkers in organotypic epidermis. *Arch. Dermatol. Res. Springer-Verlag;*
38 2009a;301(7):475–85
39
- 40 Pavez Loriè E, Cools M, Borgers M, Wouters L, Shroot B, Hagforsen E, et al. Topical treatment
41 with CYP26 inhibitor talarozole (R115866) dose dependently alters the expression of retinoid-
42 regulated genes in normal human epidermis. *Br. J. Dermatol. Blackwell Publishing Ltd;*
43 2009b;160(1):26–36
44
- 45 Perteau M, Perteau GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL. StringTie enables
46 improved reconstruction of a transcriptome from RNA-seq reads. *Nat. Biotechnol.*
47 2015;33(3):290–5
48
- 49 Poumay Y, Dupont F, Marcoux S, Leclercq-Smekens M, Hérim M, Coquette A. A simple
50 reconstructed human epidermis: Preparation of the culture model and utilization in in vitro
51 studies. *Arch. Dermatol. Res.* 2004;296(5):203–11
52
- 53 Poumay Y, Herphelin F, Smits P, De Potter IY, Pittelkow MR. High-cell-density phorbol Ester
54 and retinoic acid upregulate involucrin and downregulate Suprabasal Keratin 10 in autocrine
55 cultures of human epidermal keratinocytes. *Mol. Cell Biol. Res. Commun.* 1999;2(2):138–44
56
57
58
59
60

- 1
2
3 Radoja N, Diaz D V., Minars TJ, Freedberg IM, Blumenberg M, Tomic- Canic M. Specific
4 organization of the negative response elements for retinoic acid and thyroid hormone receptors in
5 keratin gene family. *J. Invest. Dermatol.* 1997;109(4):566–72
6
- 7 Ray WJ, Bain G, Yao M, Gottlieb DI. CYP26, a novel mammalian cytochrome P450, is induced
8 by retinoic acid and defines a new family. *J. Biol. Chem.* 1997;272(30):18702–8
9
- 10 RCoreTeam. R: A Language and Environment for Statistical Computing [Internet]. Vienna,
11 Austria; 2018. Available from: <https://www.r-project.org>
12
- 13 Rittié L, Varani J, Kang S, Voorhees JJ, Fisher GJ. Retinoid-induced epidermal hyperplasia is
14 mediated by epidermal growth factor receptor activation via specific induction of its ligands
15 heparin-binding EGF and amphiregulin in human skin in vivo. *J. Invest. Dermatol.*
16 2006;126(4):732–9
17
- 18 Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-
19 source platform for biological-image analysis. *Nat. Methods.* 2012;9(7):676–82
20
- 21 Seiberg M, Siock P, Wisniewski S, Cauwenbergh G, Shapiro SS. The effects of trypsin on
22 apoptosis, utriculi size, and skin elasticity in the Rhino mouse. *J. Invest. Dermatol.*
23 1997;109(3):370–6
24
- 25 Stehlin-Gaon C, Willmann D, Zeyer D, Sanglier S, Van Dorsselaer A, Renaud JP, et al. All-trans
26 retinoic acid is a ligand for the orphan nuclear receptor ROR β . *Nat. Struct. Biol.*
27 2003;10(10):820–5
28
- 29 Steijlen PM, Reifenschweiler DOH, Ramaekers FCS, van Muijen GNP, Happle R, Link M, et al.
30 Topical treatment of ichthyoses and Darier's disease with 13-cis-retinoic acid - A clinical and
31 immunohistochemical study. *Arch. Dermatol. Res.* 1993;285(4):221–6
32
- 33 Stoll SW, Elder JT. Retinoid regulation of heparin-binding EGF-like growth factor gene
34 expression in human keratinocytes and skin. *Exp. Dermatol.* 1998;7(6):391–7
35
- 36 Stoppie P, Borgers M, Borghgraef P, Dillen L, Goossens J, Sanz G, et al. R115866 inhibits all-
37 trans-retinoic acid metabolism and exerts retinoidal effects in rodents. *J. Pharmacol. Exp. Ther.*
38 2000;293(1):304–12
39
- 40 de Thé H, Vivanco-Ruiz MM, Tiollais P, Stunnenberg H, Dejean A. Identification of a retinoic
41 acid responsive element in the retinoic acid receptor beta gene. *Nature.* 1990;343(6254):177–80
42
- 43 Tomic-Canic M, Sunjevaric I, Freedberg IM, Blumenberg M. Identification of the retinoic acid
44 and thyroid hormone receptor-responsive element in the human K14 keratin gene. *J. Invest.*
45 *Dermatol.* 1992;99(6):842–7
46
- 47 Vahlquist A, Blockhuys S, Steijlen P, Van Rossem K, Didona B, Blanco D, et al. Oral liarozole
48 in the treatment of patients with moderate/severe lamellar ichthyosis: Results of a randomized,
49 double-blind, multinational, placebo-controlled phase II/III trial. *Br. J. Dermatol.*
50 2014;170(1):173–81
51
- 52 Vahlquist A, Gånemo A, Virtanen M. Congenital ichthyosis: An overview of current and
53 emerging therapies. *Acta Derm. Venereol.* 2008. p. 4–14
54
- 55 Vasios GW, Gold JD, Petkovich M, Chambon P, Gudas LJ. A retinoic acid-responsive element
56 is present in the 5' flanking region of the laminin B1 gene. *Proc Natl Acad Sci U S A.*
57
58
59
60

1
2
3 1989;86(23):9099–103
4

5 Verfaille CJ, Borgers M, van Steensel MAM. Retinoic acid metabolism blocking agents
6 (RAMBAs): a new paradigm in the treatment of hyperkeratotic disorders. *J. der Dtsch.*
7 *Dermatologischen Gesellschaft = J. Ger. Soc. Dermatology JDDG.* 2008;6(5):355–64
8

9 Verfaille CJ, Thissen CACB, Bovenschen HJ, Mertens J, Steijlen PM, Van De Kerkhof PCM.
10 Oral R115866 in the treatment of moderate to severe plaque-type psoriasis. *J. Eur. Acad.*
11 *Dermatology Venereol.* 2007a;21(8):1038–46
12

13 Verfaille CJ, Vanhoutte FP, Blanchet-Bardon C, Van Steensel MA, Steijlen PM. Oral liarozole
14 vs. acitretin in the treatment of ichthyosis: A phase II/III multicentre, double-blind, randomized,
15 active-controlled study. *Br. J. Dermatol.* 2007b;156(5):965–73
16

17 De Vuyst E, Charlier C, Giltaire S, De Glas V, de Rouvroit CL, Poumay Y. Reconstruction of
18 normal and pathological human epidermis on polycarbonate filter. *Methods Mol. Biol.*
19 2014;1195(1202):191–201
20

21 Van Wauwe J, Van Nyen G, Coene MC, Stoppie P, Cools W, Goossens J, et al. Liarozole, an
22 inhibitor of retinoic acid metabolism, exerts retinoid-mimetic effects in vivo. *J. Pharmacol. Exp.*
23 *Ther.* 1992;261(2):773–9
24

25 Willems E, Leyns L, Vandesompele J. Standardization of real-time PCR gene expression data
26 from independent biological replicates. *Anal. Biochem.* 2008;379(1):127–9
27

28 Xiao JH, Feng X, Di W, Peng ZH, Li L a, Chambon P, et al. Identification of heparin-binding
29 EGF-like growth factor as a target in intercellular regulation of epidermal basal cell growth by
30 suprabasal retinoic acid receptors. *EMBO J.* 1999;18(6):1539–48
31

32 Yoshimura K, Uchida G, Okazaki M, Kitano Y, Harii K. Differential expression of heparin-
33 binding EGF-like growth factor (HB-EGF) mRNA in normal human keratinocytes induced by a
34 variety of natural and synthetic retinoids. *Exp. Dermatol.* Munksgaard International Publishers;
35 2003;12 Suppl 2(s2):28–34
36

37 Zheng D, Giljohann DA, Chen DL, Massich MD, Wang X-Q, Iordanov H, et al. Topical delivery
38 of siRNA-based spherical nucleic acid nanoparticle conjugates for gene regulation. *Proc. Natl.*
39 *Acad. Sci.* 2012;109(30):11975–80
40

41 Zhu A, Ibrahim JG, Love MI. Heavy-tailed prior distributions for sequence count data: removing
42 the noise and preserving large differences. *Bioinformatics.* 2018;(November):1–9
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FIGURE LEGENDS

Figure 1: DX314 potentiates *at*RA gene expression effects in healthy RHE. (a) Relative expression of HBEGF, IVL, and CYP26A1 mRNA by RT-qPCR. Symbol underneath indicates comparison group (n=3-4 times in duplicate; mean±95% CI; *p≤0.05, **p≤0.01, ***p≤0.001; one-way ANOVA with Tukey's correction; *vs Control, †vs 1nM *at*RA, ‡vs 10nM *at*RA, ⊗vs DX314-alone, ⊕vs Liarozole-alone). **(b)** IVL localization in healthy RHE. **(c)** RNAseq: Relative mRNA expression of retinoid-responsive genes in healthy RHE. Non-grey cells differ from controls (FDR≤0.05; n=3-5). Adjacent green cell indicates likely RAR-mediated effect based on presence of RARE-promotor for respective gene. Predicted activation z-score of **(d)** upstream regulators or **(e)** canonical pathways determined by IPA software utilizing RNAseq data. Black dots indicate statistical insignificance (p≤0.05 and z-score ≥2 or ≤-2). Additional abbreviations: **Table S4.**

Figure 2: DX314 potentiates the effects of *at*RA on CYP26A1 mRNA expression in keratinocytes from individuals with keratinization disorders. Relative CYP26A1 mRNA expression by RT-qPCR in; **(a)** Darier disease RHE, **(b)** recessive x-linked ichthyosis (RXLI) full-thickness RHE, **(c)** lamellar ichthyosis RHE, and **(d)** RXLI monolayer keratinocyte cultures. RHE were treated for 4 days and monolayer keratinocytes for 20hrs. Statistical significance was computed with **(a)** autoscaled or **(b-d)** raw dCt values. Symbol below each treatment indicates comparison group (n=3 independent replicates with technical duplicates; mean±95% CI; *p≤0.05; **p≤0.01; ***p≤0.001; one-way ANOVA with Tukey's correction; *vs Control, †vs 1nM *at*RA, ‡vs 10nM *at*RA, §vs 100nM *at*RA, ◆vs 1000nM *at*RA, ⊗vs DX314-alone, ⊕vs Liarozole-alone, ⊖vs Talarozole-alone).

1
2
3
4
5
6 **Figure 3: DX314 potentiates the effects of *at*RA on the expression and localization of**
7
8 **keratin 10 (KRT10) in Darier disease (DD) RHE. *At*RA, but not DX314, induces a loss of**
9
10 **stratum granulosum. (a) HE staining and (b) immunofluorescent staining of KRT10 (green)**
11
12 localization with nuclear stain (blue), in DD RHE treated for 4 days. Scale bars: black = 20 μ m,
13
14 white =50 μ m. (c) Relative KRT10 mRNA expression by qPCR. Symbol below each treatment
15
16 indicates comparison group. (n=3 independent replicates with technical duplicates; mean \pm 95%
17
18 CI; *p \leq 0.05; **p \leq 0.01; ***p \leq 0.001; one-way ANOVA with Tukey's correction on autoscaled
19
20 values; *vs Control, †vs 1nM *at*RA, ‡vs 10nM *at*RA, §vs 100nM *at*RA, ¶vs DX314-alone, ¯vs
21
22 Talarozole-alone).

23
24
25
26
27
28 **Figure 4: DX314 protects barrier function in RHE. (a) Transepithelial electrical resistance**
29
30 (TEER) in healthy RHE. TEER was normalized to control RHEs for each run, then pooled for
31
32 analysis. Graph shows Tukey's boxplot with outliers. Sample sizes (n) are shown above x-axis.
33
34 (b) LI RHE TEER (top), transepidermal water loss (middle), and the linear correlation between
35
36 the two measures (bottom). (c) HE staining of lamellar ichthyosis (LI) RHE. Scale bar = 50 μ m.
37
38 (d) Semi-quantitative analysis of relative stratum granulosum (SG) surface area in healthy RHE.
39
40 (e) Relative expression of epidermal differentiation complex (EDC) genes and regulators by
41
42 RNAseq. Colored (non-grey) cells indicate statistical significance from control (FDR \leq 0.05; n=3-
43
44 5). All RHE received a 4-day treatment. (a,b,d) (*p \leq 0.05; **p \leq 0.01; ***p \leq 0.001; one-way
45
46 ANOVA with Dunnett's correction vs control).
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Figure 5: DX314 reduces rhino mouse skin abnormalities.** Representative HE staining of skin
4
5 biopsies from rhino mice topically treated for 11 days with vehicle (acetone) or 1% DX314.
6
7 ImageJ software was used to quantify comedonal number, profile (d/D, ratio of opening to inner
8
9 diameter), and epidermal thickness. Epidermal thickness was measured at multiple points across
10
11 each sample by measuring the sum of epidermal areas (yellow), excluding the corneal layer, and
12
13 dividing by the sum of the length of the basal layers (dotted blue line). Scale bar =200µm.
14
15
16
17
18
19

20 **Figure 6: DX314 treatment reduces comedonal number, induces epidermal thickening, and**
21 **increases comedonal profile, while having no effect on transepidermal water loss (TEWL)**
22 **in treated rhino mice.** Semi-quantitative analysis of changes in (a) total (open + closed) and (b)
23
24 open comedonal number, (c) comedonal profile, and (d) epidermal thickness in rhino mice
25
26 topically treated with vehicle (acetone) or 1% DX314 over 11 days. (e) Daily TEWL
27
28 measurements did not reveal any statistically significant differences between treatment groups.
29
30 (n=5-6 mice per treatment; mean±SD; *p≤0.05; **p≤0.01; Student's t-test vs vehicle control).
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1: DX314 potentiates atRA gene expression effects in healthy RHE. (a) Relative expression of HBEGF, IVL, and CYP26A1 mRNA by RT-qPCR. Symbol underneath indicates comparison group (n=3-4 times in duplicate; mean±95% CI; *p≤0.05, **p≤0.01, ***p≤0.001; one-way ANOVA with Tukey's correction; *vs Control, †vs 1nM atRA, ‡vs 10nM atRA, ⊗vs DX314-alone, ⊕vs Liarozole-alone). **(b)** IVL localization in healthy RHE. **(c)** RNAseq: Relative mRNA expression of retinoid genes in healthy RHE. Non-grey cells differ from controls (FDR≤0.05; n=3-5). Adjacent green cell indicates likely RAR-mediated effect based on presence of RARE-promotor for respective gene. Predicted activation z-score of **(d)** upstream regulators or **(e)** canonical pathways determined by IPA software utilizing RNAseq data. Black dots indicate statistical insignificance (p≤0.05 and z-score ≥2 or ≤-2). Additional abbreviations: **Table S4**.




Figure 2: DX314 potentiates the effects of atRA on CYP26A1 mRNA expression in keratinocytes from individuals with keratinization disorders. Relative CYP26A1 mRNA expression by RT-qPCR in; **(a)** Darier disease RHE, **(b)** recessive x-linked ichthyosis (RXLI) full-thickness RHE, **(c)** lamellar ichthyosis RHE, and **(d)** RXLI monolayer keratinocyte cultures. RHE were treated for 4 days and monolayer keratinocytes for 20hrs. Statistical significance was computed with **(a)** autoscaled or **(b-d)** raw dCt values. Symbol below each treatment indicates comparison group (n=3 independent replicates with technical duplicates; mean±95% CI; *p≤0.05; **p≤0.01; ***p≤0.001; one-way ANOVA with Tukey's correction; *vs Control, †vs 1nM atRA, ‡vs 10nM atRA, &vs 100nM atRA, ◆vs 1000nM atRA, ⊗vs DX314-alone, ⊕vs Liarozole-alone, ∅vs Talarozole-alone).

150x135mm (600 x 600 DPI)




Figure 3: DX314 potentiates the effects of atRA on the expression and localization of keratin 10 (KRT10) in Darier disease (DD) RHE. AtRA, but not DX314, induces a loss of stratum granulosum. (a) HE staining and (b) immunofluorescent staining of KRT10 (green) localization with nuclear stain (blue), in DD RHE treated for 4 days. Scale bars: black = 20 μ m, white = 50 μ m. (c) Relative KRT10 mRNA expression by qPCR. Symbol below each treatment indicates comparison group. (n=3 independent replicates with technical duplicates; mean \pm 95% CI; *p \leq 0.05; **p \leq 0.01; ***p \leq 0.001; one-way ANOVA with Tukey's correction on autoscaled values; *vs Control, †vs 1nM atRA, ‡vs 10nM atRA, §vs 100nM atRA, ⊗vs DX314-alone, ∅vs Talarozole-alone).

182x171mm (600 x 600 DPI)




Figure 4: DX314 protects barrier function in RHE. (a) Transepithelial electrical resistance (TEER) in healthy RHE. TEER was normalized to control RHEs for each run, then pooled for analysis. Graph shows Tukey's boxplot with outliers. Sample sizes (n) are shown above x-axis. **(b)** LI RHE TEER (top), transepidermal water loss (middle), and the linear correlation between the two measures (bottom). **(c)** HE staining of lamellar ichthyosis (LI) RHE. Scale bar = 50μm. **(d)** Semi-quantitative analysis of relative stratum granulosum (SG) surface area in healthy RHE. **(e)** Relative expression of epidermal differentiation complex (EDC) genes and regulators by RNAseq. Colored (non-grey) cells indicate statistical significance from control (FDR≤0.05; n=3-5). All RHE received a 4-day treatment. **(a,b,d)** (*p≤0.05; **p≤0.01; ***p≤0.001; one-way ANOVA with Dunnett's correction vs control).

86x119mm (600 x 600 DPI)




Figure 5: DX314 reduces rhino mouse skin abnormalities. Representative HE staining of skin biopsies from rhino mice topically treated for 11 days with vehicle (acetone) or 1% DX314. ImageJ software was used to quantify comedonal number, profile (d/D , ratio of opening to inner diameter), and epidermal thickness. Epidermal thickness was measured at multiple points across each sample by measuring the sum of epidermal areas (yellow), excluding the corneal layer, and dividing by the sum of the length of the basal layers (dotted blue line). Scale bar = $200\mu\text{m}$.

221x256mm (600 x 600 DPI)




Figure 6: DX314 treatment reduces comedonal number, induces epidermal thickening, and increases comedonal profile, while having no effect on transepidermal water loss (TEWL) in treated rhino mice. Semi-quantitative analysis of changes in **(a)** total (open + closed) and **(b)** open comedonal number, **(c)** comedonal profile, and **(d)** epidermal thickness in rhino mice topically treated with vehicle (acetone) or 1% DX314 over 11 days. **(e)** Daily TEWL measurements did not reveal any statistically significant differences between treatment groups. (n=5-6 mice per treatment; mean±SD; *p≤0.05; **p≤0.01; Student's t-test vs vehicle control).

113x110mm (600 x 600 DPI)

SUPPLEMENTARY METHODS

Immunostaining

Slides were sequentially exposed to the following: PBS rinse, 2x 3min 0.1M glycine in dH₂O, PBS rinse, then 1hr in PBS/BSA/Triton (PBS with 0.2% BSA and 0.02% Triton X-100). A hydrophobic marker was used to encircle the tissue and 50 μ L of primary antibody (diluted in PBS/BSA/Triton) was applied. The slides were placed in a humidity chamber and incubated for 1hr at room temperature. The slides were then rinsed three times in PBS/BSA/Triton before 1hr humidified incubation with 50 μ L of the respective secondary antibody. The slides were again rinsed three times in PBS/BSA/Triton before a 15min incubation with 50 μ L of Hoechst nuclear stain (diluted in PBS/BSA/Triton) followed by 3x 5min PBS rinses. Coverslips were mounted with Mowiol 40-88 and the slides were stored at 4°C until imaged on an Olympus DX63 microscope with Olympus XM10 camera. Antibodies and dilutions can be found below (**Table S3**).

RNA isolation and quantitative PCR

Following treatment, RHEs intended for RNA extraction were flash frozen at -80°C until use. RNA was isolated using the NucleoSpin RNA (Macherey-Nagel, Bethlehem, PA) kit, as recommended by the manufacturer. Variations from the standard protocol include homogenization with 600 μ L (rather than 350 μ L) of RA1 lysis buffer, addition of 6 μ L of β -mercaptoethanol to RA1 to aid tissue lysis, and addition 600 μ L of 70% ethanol (rather than 350 μ L) during nucleic acid precipitation. RNA obtained from monolayer and full-thickness RHE cultures were isolated using TRIzol (Life Technologies, Burlington, Canada) phenol-chloroform extraction as described by the manufacturer. Variations from the standard protocol include chilling the sample following the addition of isopropanol to encourage nucleic acid precipitation,

1
2
3 and the addition of a second chilled 75% ethanol wash prior to drying the pellet, which greatly
4
5 improved consistency in RNA purity.
6

7
8 Isolated RNA purity and concentration were measured by NanoDrop 2000c (Thermo
9
10 Scientific, Rockford, IL) and integrity was confirmed by gel electrophoresis. 100-200ng of
11
12 template RNA was reverse-transcribed to cDNA with the Superscript III reverse transcriptase kit
13
14 (Invitrogen, Aalst, Belgium). The cDNA was then diluted 1:10 with water. 2 μ L of cDNA was
15
16 added to a 10 μ L real-time qPCR reaction, which used Takyon No ROX SYBR 2X MasterMix
17
18 (Eurogentec, Seraing, Belgium) on a Roche Lightcycler 96 (activation: 3min - 95°C; 40 cycles:
19
20 10sec - 95°C, 20sec - 60°C, 30sec - 72°C). 500nM of each primer pair (**Table S2**), optimized for
21
22 an annealing temperature of 60°C, was used for each reaction. RPL13a and 36B4 (RPLP0) were
23
24 used as reference genes.
25
26
27

28 **RNA sequencing and bioinformatics**

29
30
31 RNA samples were sent to the University of Colorado's Genomics and Sequencing Core
32
33 Facility (Denver, CO) for library preparation and sequencing. Purity and concentration were
34
35 measured with an Agilent Bioanalyzer (Agilent Technologies, Santa Clara, CA). 200-500ng of
36
37 RNA was used to prepare the Illumina HiSeq libraries according to manufacturer's instructions
38
39 for the TruSeq Stranded RNA kit (Illumina, San Diego, CA). Sequencing was done as 2x151bp
40
41 paired end reads on the Illumina HiSeq4000. The bioinformatics pipeline used was an adaptation
42
43 of several described methods (Kim et al. 2016; Love et al. 2013): FastQC (Andrews 2010)
44
45 (v0.11.3) → Trimmomatic (Bolger et al. 2014) (v0.38) → FastQC → Hisat2 (Kim et al. 2015)
46
47 (v2.1.0) → SamTools (Li et al. 2009) (v1.8) → Stringtie (Pertea et al. 2015) (v1.3.4d). Reads
48
49 were mapped to the H. sapien GRCh38 *genome_tran* index provided by HiSat2 developers (Kim
50
51 et al. 2017). Differential expression analysis was performed in R (RCoreTeam 2018) (v3.5.2)
52
53
54
55
56
57
58
59
60

1
2
3 using DESeq2 (Love et al. 2014) (v1.22.2). False-discovery rate (FDR) was determined with the
4
5 Benjamini-Hochberg correction, logarithmic-fold change shrinkage to correction was performed
6
7 using the apeglm (Zhu et al. 2018) (v1.4.2) algorithm, HGNC names were attached using
8
9 BiomaRt (Durinck et al. 2009) (v2.38.0). Treatment groups: Control (n=5), 1nM *atRA* (n=4),
10
11 100nM *atRA* (n=4), 1000nM DX314 (n=3), and 1000nM DX314 + 1nM *atRA* (n=4).
12
13

14
15 Fold-change (FC) and FDR values mapped to Ensembl IDs were imported into Ingenuity
16
17 Pathway Analysis (IPA) (Qiagen, Germantown, MD) software for analysis. Analysis was limited
18
19 to genes with $FDR \leq 0.05$, and $FC \geq 1.5$ or ≤ -1.5 . IPA uses Fisher's exact test ($p \leq 0.05$) to
20
21 predict the activation or inhibition of canonical pathways and upstream regulators. A z-score of \geq
22
23 2.0 is considered activated, a $z \leq -2.0$ is considered inhibited.
24
25

26 **Nuclear receptor profiling**

27
28 Nuclear receptor activity profiling was performed by luciferase gene reporter assay using
29
30 the Steady-Glo luciferase assay system (Promega, E2550) and 16 recombinant HeLa cell lines in
31
32 which the respective human nuclear receptors were overexpressed. Compounds were solubilized
33
34 at 10mM in DMSO and all assays were performed at a final concentration of 0.1% DMSO.
35
36 Dose-response curves were generated with 8 compound concentrations (n=2) in 4-fold serial
37
38 dilutions from a maximum test dose of 10 μ M. Cells in growth media (DMEM with 1g/L glucose
39
40 (Invitrogen, 31885-023), 2mM L-glutamine (Invitrogen, 25030-024), 1x MEM non-essential
41
42 amino acids (Invitrogen, 11140-035), 0.1% antibiotic-antimycotic (Invitrogen, 15240-062),
43
44 0.5mg/mL G418 (Invitrogen, 10131-027), 0.5 μ g/mL puromycin (Sigma, P8833), 10% FBS
45
46 (Biowest, S800)) were seeded in 96-well plates at 2e4 cells/well and incubated at 37°C with 5%
47
48 CO₂ for 4hrs. Compounds in test media (growth media with 0.1% Pluronic F-127 (Interchim, FP-
49
50 379951)) were then added to each well and the plates were returned to the incubator for 18-
51
52
53
54
55
56
57
58
59
60

1
2
3 24hrs. 50 μ L/well of Steady-Glo was added to each well and plates were shaken for 5min at room
4
5 temperature before measuring luciferase activity (RLU) using a BMG Clariostar plate reader.
6

7
8 Data was normalized to negative (Min) and positive (Max) controls: $\frac{(x - Min) * 100}{(Max - Min)}$. 4-
9
10 parameter logistic model fitting was performed using XLfit software (IDBS). Where n is the hill
11
12 coefficient: $Normalized\ Activation\ (\%) = Min + \frac{(Max - Min)}{1 + (\frac{EC_{50}}{x})^n}$.
13
14
15

16 17 **Rhino mice**

18
19 Rhino mice (2-3 males, 3 females per group), aged 5-7 weeks were acquired from
20
21 Jackson Laboratory (Sacramento, CA) and acclimated to standard housing and provisions for one
22
23 week prior to start. All animal studies were approved by Institutional Animal Care and Use
24
25 Committee under NIH guidelines. Mice received daily topical application of 50 μ L vehicle
26
27 (acetone), or freshly made 1% DX314, on a 2x2cm area of back skin for 11 days. Daily clinical
28
29 observations, body weights, TEWL, and DRAIZE scoring were recorded (**Table S5** and **S6**).
30
31
32 Mice underwent necropsy on day 12 and skin tissue was collected and processed.
33
34

35
36 Quantitative analysis of the tissue was performed using Fiji/ImageJ software following a
37
38 previously described standardized method (Bouclier et al. 1991). Each sample was analyzed at
39
40 least two different section depths and totaling 6.2-12.8mm of skin per subject. Open comedone
41
42 profile was quantified as the ratio of the opening size (d) and the internal diameter (D).
43
44 Epidermal thickness was measured (17+ unique points per subject) by measuring the epidermal
45
46 area, excluding the SC, and dividing by the length of the underlying basal layer (**Figure 5**).
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Keratinocytes	Source	Pathology	Age/Sex	Figure(s)
NAK209	Poumay/Bienfait	Healthy	48/F	1a-b, 4a
NAK219			37/M	1a, 4a
NAK214			32/F	1a, 4a, S1, S2
HEKa	Gibco, C0055C		40/F	1c-e, 4d-e, S3
DARK1	Poumay/Bienfait	Darier disease	Child/M	2a, 3a-c
RXLI1653	Paller	Recessive x-linked ichthyosis	13/M	2b
RXLI1658			29/M	2d
LI173			Lamellar ichthyosis	9/M

For Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Target	Forward	Reverse
36B4	ATCAACGGGTACAAACGAGTC	CAGATGGATCAGCCAAGAAGG
RPL13a	CTCAAGGTCGTGCGTCTGAA	TGGCTGTCACTGCCTGGTACT
HBEGF	TGGCCCTCCACTCCTCATC	GGGTCACAGAACCATCCTAGCT
IVL	TGAAACAGCCAACCTCCAC	TTCCTCTTGCTTTGATGGG
KRT10	ATCGATGACCTTAAAAATCAGATTCTC	GCAGAGCTACCTCATTCTCATAC
CYP26A1	GGGAGAGCGGCTGGACAT	TCCAAAGAGGAGTTCGGTTGA
CYP26B1	CCGCTTCCATTACCTCCCGTTC	CCACCGCCAGCACCTTCAG

For Review Only

Target	Species	Dilution	Source
IVL	Mouse	1:200	Invitrogen, I9018
KRT10	Mouse	1:100	DAKO, M7002
2°, HRP-conjugated (anti-mouse)	Horse	1:100	Vectastain, PK-4002
2°, Alexa Fluor 488-conjugated (anti-mouse)	Goat	1:1000	Life Technologies, A11001
Hoechst 33258, nuclear stain	-	1:100	Life Technologies, H3569

For Review Only


AD	atopic dermatitis
ANOVA	analysis of variance
AQP	aquaporin
ARNT	aryl hydrocarbon receptor nuclear translocator
ARP	actin-related protein
atRA	all-trans retinoic acid
BHLH	basic helix-loop-helix
Clorf	chromosome 1 open reading frame
CASP	caspase
CE	cornified envelope
CLDN	claudin
CRABP	cellular retinoic acid binding protein
CRBP	cellular retinol binding protein
CRCT	cysteine rich C-terminal
CRNN	cornulin
CYP	cytochrome P450
DD	Darier disease
DMSO	dimethyl sulfoxide
DNA	deoxyribonucleic acid
DSC	desmocollin
ECM	extracellular matrix
EDC	epidermal differentiation complex
EDTA	ethylenediaminetetraacetic acid
EGF	epidermal growth factor
EHF	ETS homologous factor
ELF	E74 like ETS transcription factor
ERBB	Erb-B2 Receptor Tyrosine Kinase
Erk	extracellular signal-regulated kinases
FBS	fetal bovine serum
FDR	false discovery rate
FGF	fibroblast growth factor
FLG	filaggrin
GATA	GATA binding protein
GRHL	grainyhead like transcription factor
GTF	gene transfer format
HBEGF	heparin-binding EGF-like growth factor
HE	hematoxylin and eosin
HGNC	HUGO gene nomenclature committee
HIC	HIC ZBTB transcriptional repressor
HKGS	human keratinocyte growth supplement
HRNR	hornerin
IF	immunofluorescence
IFNG	interferon gamma
IHC	immunohistochemistry
IL	interleukin
ILK	integrin-linked kinase
IPA	ingenuity pathway analysis
IVL	involucrin
JAK/STAT	janus kinases/signal transducer and activator of transcription

JAM	junctional adhesion molecules
KG	keratohyalin granules
KGF	keratinocyte growth factor
KLF	kruppel-like factor
KLK	kallikrein related peptidase
KRT	keratin
LAMB	laminin subunit beta
LCE	late cornified envelope
LI	lamellar ichthyosis
LOR	loricrin
LRAT	lecithin:retinol acetyltransferase
LXR	liver X receptor
MAPK	mitogen-activated protein kinases
MEIS	Meis homeobox
ML	Monolayer
NFE	nuclear factor, erythroid
NR	nuclear receptor
NRIP	nuclear receptor interacting protein
OCLN	occludin
OR	organotypic raft
P/S	penicillin/streptomycin
PBS	phosphate buffered saline
PKC	protein kinase C
PPAR	peroxisome-proliferator-activated receptor
RA	retinoic acid
RABL	rab-like protein
RAL	retinal
RalDH	retinal dehydrogenase
RAMBA	retinoic acid metabolism blocking agents
RAR	retinoic acid receptor
RARE	retinoic acid response element
RARRES	retinoic acid receptor responder
RB1	retinoblastoma-associated protein
RBP	retinol binding protein
RE	retinyl esters
RHE	reconstructed human epidermis
RHOGDI	RHO protein GDP dissociation inhibitor
RNA	ribonucleic acid
ROL	retinol/vitamin A
ROR	RAR-related orphan receptor
RPTN	repetin
RT qPCR	real-time quantitative polymerase chain reaction
RXLI	recessive x-linked ichthyosis
RXR	retinoid X receptor
S100	S100 calcium binding protein
SAM	sequence alignment map
SB	stratum basale
SC	stratum corneum
SDR	short-chain dehydrogenases/reductases

SG	stratum granulosum
SMARC	SWI/SNF-related matrix-associated actin-dependent regulator of chromatin
SPINK	serine protease inhibitor kazal-type
SPR	small proline-rich protein
SS	stratum spinosum
SSWL	subsurface water loss
STRA	stimulated by retinoic acid
TCHH/THH	trichohyalin
TEER	transepithelial electrical resistance
TEWL	trans-epidermal water loss
TGF	transforming growth factor
TGM	transglutaminases
THOP	thimet oligopeptidase
TJ	tight junction
TLR	toll-like receptor
TNF	tumor necrosis factor
TP53	tumor protein p53
TR	thyroid receptor
TREM	triggering receptor expressed on myeloid cells
TTR	transthyretine
UV	ultraviolet
VDR	vitamin D receptor
WASP	Wiskott–Aldrich Syndrome protein
Wnt	wingless/integrated
ZO	zona occludens

Group	Animal Number	Clinical Observation (Days Observed)
Vehicle	101*	NSO (Day 0-5), slight dry skin on upper back (Day 6), dry skin on upper back (Day 7-8), slight dry skin on upper back (Day 9-11) NSO Day 12)
	102*	NSO (Day 0-8), very slight dry skin on upper back (Day 9), NSO (Day 10-12)
	103*	NSO (Day 0-5), slight dry skin on upper back (Day 6-7), NSO (Day 8), two small bite marks at base of tail (Day 9-11), NSO (Day 12)
	152^Λ	NSO (Day 0-6), Slight dry skin on upper back (Day 7-8), NSO (Day 9), Slight dry skin on upper back (Day 10-11), NSO (Day 12)
	153^Λ	NSO (Day 0-7), Slight dry skin on upper back (Day 8), NSO (Day 9-12)
1% DX314	401*	NSO (Day 0-5), Slight dry skin on upper back (Day 6-8), brown/tan patch across mid back (Day 9-12)
	402*	NSO (Day 0-6), Dry skin on upper back (Day 7), Slight dry skin on upper back (Day 8), Brown/tan patch across mid back (Day 9-12)
	403*	NSO (Day 0-6), Dry skin on upper back (Day 7), Slight dry skin on upper back (Day 8), Darker brown/tan patch across mid back (Day 9-12)
	451^Λ	NSO (Day 0-6), slight dry skin on upper back (Day 7-8), NSO (Day 9-11), slight tan patch on mid back (Day 12)
	452^Λ	NSO (Day 0-6), Slight dry skin on upper back (Day 7-8), NSO (Day 9-12)
	453^Λ	NSO (Day 0-6), Slight dry skin on upper back (Day 7-8), NSO (Day 9-11), OD: corneal edema, swollen and opaque; tan patch on back (Day 12)

Group	Animal	Weight						DRAIZE Scoring (Erythema : Edema)		
		Day 0		Day 7		Day 12		Day 0	Day 7	Day 12
		g	%	g	%	g	%			
Vehicle	101*	24.5	100.0	20.8	84.9	21.9	89.4	0 : 0	0 : 0	0 : 0
	102*	24.4	100.0	25.9	106.1	25.2	103.3	0 : 0	0 : 0	0 : 0
	103*	18.5	100.0	19.1	103.2	19.8	107.0	0 : 0	0 : 0	0 : 0
	152^	19.3	100.0	19.6	101.6	19.7	102.1	0 : 0	0 : 0	0 : 0
	153^	21.9	100.0	21.8	99.5	23.2	105.9	0 : 0	0 : 0	0 : 0
1% DX314	401*	23.4	100.0	23.1	98.7	23.8	101.7	0 : 0	0 : 0	0 : 0
	404*	22.6	100.0	23.2	102.7	23.4	103.5	0 : 0	0 : 0	0 : 0
	403*	21.9	100.0	22.8	104.1	23.2	105.9	0 : 0	0 : 0	0 : 0
	451^	19.1	100.0	19.7	103.1	20.5	107.3	0 : 0	0 : 0	0 : 0
	452^	21.4	100.0	22.4	104.7	22.3	104.2	0 : 0	0 : 0	0 : 0
	453^	20.6	100.0	20.7	100.5	21.4	103.9	0 : 0	0 : 0	0 : 0



16 **Figure S1: DX314 potentiates the effects of atRA on KRT10 localization in healthy RHE.**
17 Immunofluorescent staining of KRT10 (green) localization with nuclear stain (blue) in healthy RHE treated
18 for 4 days. Scale bar = 50μm.

19 148x39mm (300 x 300 DPI)

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure S2: AtRA causes abnormal morphological effects in healthy RHE, which are potentiated by liarozole and talarozole, but not DX314. HE staining of healthy RHE treated for 4 days by atRA or RAMBAs with, and without, atRA. Scale bar = 20 μ m. (SC, stratum corneum; SG, stratum granulosum; SS, stratum spinosum; SB, stratum basale).

110x92mm (300 x 300 DPI)




Figure S3: Tissues used in semi-quantitative analysis of relative stratum granulosum (SG) surface area. Healthy RHE were treated for 4 days before processing, HE staining, and imaging. Shown are cropped images of the samples used to determine SG surface area as a percent of total tissue surface area (**Figure 4d**). Scale bar = 50 μ m.

169x100mm (300 x 300 DPI)




Figure S4: DX314 acts as an inverse agonist on ROR α and ROR γ , displays no activity on other nuclear receptors. (a) Table of EC₅₀ values for DX314 and respective reference ligand. Non-linear regression of normalized **(b)** ROR α and **(c)** ROR γ luciferase reporter assay activity. IA = Inactive.

233x103mm (300 x 300 DPI)