

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

Neisseria cinerea Expresses a Functional Factor H Binding Protein Which Is Recognized by Immune Responses Elicited by Meningococcal Vaccines

Lavender, Hayley; Poncin, Katy; Tang, Christoph M.

Published in:
Infection and Immunity

DOI:
[10.1128/iai.00305-17](https://doi.org/10.1128/iai.00305-17)

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (HARVARD):
Lavender, H, Poncin, K & Tang, CM 2017, 'Neisseria cinerea Expresses a Functional Factor H Binding Protein Which Is Recognized by Immune Responses Elicited by Meningococcal Vaccines', *Infection and Immunity*, vol. 85, no. 10, e00305-17. <https://doi.org/10.1128/iai.00305-17>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Neisseria cinerea Expresses a Functional Factor H Binding Protein Which Is Recognized by Immune Responses Elicited by Meningococcal Vaccines

Hayley Lavender, Katy Poncin,* Christoph M. Tang

Sir William Dunn School of Pathology, University of Oxford, Oxford, United Kingdom

ABSTRACT *Neisseria meningitidis* is a major cause of bacterial meningitis and sepsis worldwide. Capsular polysaccharide vaccines are available against meningococcal serogroups A, C, W, and Y. More recently two protein-based vaccines, Bexsero and Trumenba, against meningococcal serogroup B strains have been licensed; both vaccines contain meningococcal factor H binding protein (fHbp). fHbp is a surface-exposed lipoprotein that binds the negative complement regulator complement factor H (CFH), thereby inhibiting the alternative pathway of complement activation. Recent analysis of available genomes has indicated that some commensal *Neisseria* species also contain genes that potentially encode fHbp, although the functions of these genes and how immunization with fHbp-containing vaccines could affect the commensal flora have yet to be established. Here, we show that the commensal species *Neisseria cinerea* expresses functional fHbp on its surface and that it is responsible for recruitment of CFH by the bacterium. *N. cinerea* fHbp binds CFH with affinity similar to that of meningococcal fHbp and promotes survival of *N. cinerea* in human serum. We examined the potential impact of fHbp-containing vaccines on *N. cinerea*. We found that immunization with Bexsero elicits serum bactericidal activity against *N. cinerea*, which is primarily directed against fHbp. The shared function of fHbp in *N. cinerea* and *N. meningitidis* and cross-reactive responses elicited by Bexsero suggest that the introduction of fHbp-containing vaccines has the potential to affect carriage of *N. cinerea* and other commensal species.

KEYWORDS *Neisseria cinerea*, *Neisseria meningitidis*, complement, fHbp, vaccine

The genus *Neisseria* includes two major human pathogens, *Neisseria meningitidis* and *Neisseria gonorrhoeae* (1, 2). *N. meningitidis* is a commensal of the human nasopharynx in approximately 10 to 40% of the general population but can traverse the epithelial layer and cause systemic disease (3), while the gonococcus is a leading cause of sexually transmitted disease (4). In contrast, nonpathogenic *Neisseria* species are ubiquitous commensals of the human oral cavity and nasopharynx in healthy individuals and constitute approximately 10% of the human oral bacterial flora (1, 5, 6). Even though *Neisseria polysaccharea*, *Neisseria lactamica*, and *Neisseria cinerea* are closely related to the meningococcus and the gonococcus (7, 8), there are only rare case reports of invasive disease caused by these commensal species, and they are usually limited to immunocompromised hosts (9).

N. meningitidis can be categorized into 13 serogroups based on the structure of its polysaccharide capsule, with 6 serogroups (A, B, C, W, X, and Y) responsible for the majority of disease worldwide (3, 10–13). Capsular polysaccharide conjugate vaccines have been developed to protect individuals against disease caused by some serogroups of *N. meningitidis* (i.e., A, C, W, and Y), while outer membrane vesicle (OMV) vaccines have been successfully employed to combat epidemic disease caused by single strains (14, 15). However, these approaches cannot prevent endemic serogroup

Received 2 May 2017 Returned for modification 2 June 2017 Accepted 7 July 2017

Accepted manuscript posted online 24 July 2017

Citation Lavender H, Poncin K, Tang CM. 2017. *Neisseria cinerea* expresses a functional factor H binding protein which is recognized by immune responses elicited by meningococcal vaccines. Infect Immun 85:e00305-17. <https://doi.org/10.1128/IAI.00305-17>.

Editor Andreas J. Bäumlér, University of California, Davis

Copyright © 2017 Lavender et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Christoph M. Tang, christoph.tang@path.ox.ac.uk.

* Present address: Katy Poncin, Microorganisms Biology Research Unit (URBM), Narilis, University of Namur, Namur, Belgium.

B. N. meningitidis disease, which is still a significant cause of morbidity and mortality in Europe and the United States (3). This is because immunization with the serogroup B capsule, which is identical to a modification on NCAM-1 (16), can induce autoimmunity, while OMV vaccines offer only limited cross protection against diverse strains (17–20).

Two recently licensed serogroup B vaccines, Bexsero and Trumemba, contain factor H binding protein (fHbp) as a key antigen (21–23). fHbp is a 27-kDa surface lipoprotein that is expressed by almost all disease-causing isolates of *N. meningitidis* (24, 25). The protein elicits serum bactericidal activity (SBA), an accepted correlate of protection against meningococcal disease (21, 22, 24). fHbp can be classified into one of three variant groups (V1, V2, or V3) or two subfamilies (A and B) based on its amino acid sequence (22, 26). Immunization with fHbp generally induces cross protection against strains expressing an fHbp within the same variant group, with limited cross protective immunity between variant groups (21).

fHbp promotes survival of *N. meningitidis* in serum by binding the negative complement regulator complement factor H (CFH) at high affinity (25). CFH consists of 20 complement control protein (CCP) domains joined by short linker sequences, with each domain consisting of approximately 60 amino acids (27–29). CFH inhibits the alternative pathway (AP) of complement by acting as a cofactor for factor I-mediated cleavage of C3b, a key molecule in the complement cascade, and by accelerating the decay of the AP C3 convertase (30–32). These functions are mediated by the first four CCP domains (CFH_{1–4}), while other CCP domains have distinct functions, such as binding to heparin and glycosaminoglycans (33, 34). *N. meningitidis* fHbp binds CCP domains 6 and 7 of CFH (CFH_{6–7}) (25, 35), while a further complement protein, CFHR3, also binds to fHbp and acts as a competitive antagonist of CFH (36).

Recent genome analysis of 80 pathogenic and commensal *Neisseria* strains identified *fhbp* in all *N. meningitidis* strains examined and homologues of *fhbp* in some commensal species (37). A homologue of *fhbp* was also present in isolates of *N. gonorrhoeae*, although the protein encoded by this gene, Ghfp, does not bind CFH to any significant degree (38, 39). *fhbp* was also found in all isolates of *N. cinerea* and *N. polysaccharea* examined (37–39). Of note, *N. cinerea fhbp* genes have significant nucleotide identity with V1 *fhbp*, whereas *N. gonorrhoeae* and *N. polysaccharea* harbor homologues of V3 *fhbp* (37, 39).

Several lines of evidence indicate that commensal *Neisseria* spp. can induce immunity against *N. meningitidis*. For example, colonization with *N. lactamica* during early childhood is associated with the development of adaptive immunity against the meningococcus (5, 40, 41). Moreover, individuals colonized with *N. lactamica* following live challenge develop cross-reactive salivary IgA and serum IgG antibodies and appear to be protected against subsequent acquisition of *N. meningitidis* (42). Therefore, the presence of antigens, such as fHbp, in commensal *Neisseria* spp. raises the possibility that implementation of protein-based vaccines containing fHbp could impact the colonization of commensal species in the upper airway.

Here, we investigated whether *fhbp* in the *N. cinerea* genome encodes a functional protein and examined its roles in binding CFH and complement evasion. We found that *N. cinerea* expresses fHbp on its surface, where it binds CFH at high affinity and mediates resistance against complement-mediated lysis. Furthermore, we evaluated whether immune responses elicited by Bexsero have significant activity against *N. cinerea*. We demonstrate that immune responses elicited by Bexsero have SBA against *N. cinerea*, with fHbp the main target of SBA responses elicited by Bexsero against *N. cinerea*, indicating that introduction of fHbp-containing vaccines could affect carriage of commensal *Neisseria* spp.

RESULTS

***N. cinerea* fHbp is predicted to be structurally similar to meningococcal fHbp.**

The genome sequences of five *N. cinerea* isolates, CCUG 346T, CCUG 5746, CCUG 25879, CCUG 27178A, and CCUG 53043, from the PubMLST *Neisseria* BIGSdb database (<http://pubmlst.org/neisseria/>) were interrogated for the presence of *fhbp*. Previously anno-

A

N. meningitidis V1.1 *fhbp* VNRTAFCCSLT**T**ALILITACSSGGGGVAADIGAGLADALTAPLDHKDKGLQSL
N. cinerea V1.110 *fhbp* VNRTAFCCSLT**A**ALILITACSSGGGGVAADIGAGLADALTAPLDHKDKGLQSL

N. meningitidis V1.1 *fhbp* TLDQSVRKNEKLKLAQAQGAEKTYGNGDSLNTGKLNKDKVSRFDFIRQIEVDGQ
N. cinerea V1.110 *fhbp* TLDQSVRKNEKLKLAQAQGAEKTYGNGDSLNTGKLNKDKVSRFDFIRQIEVDGQ

N. meningitidis V1.1 *fhbp* LITLESGEFQVYKQSHSALTAFQTEQIQDSEHSGKMVAKR**Q**FRIGDIAGEHTS
N. cinerea V1.110 *fhbp* LITLESGEFQVYKQSHSALTAFQTEQIQDSEHSGKMVAKR**R**FRIGDIAGEHTS

N. meningitidis V1.1 *fhbp* FDKLPEGGRATYRGTAFGSDDAGGKLTYTIDFAAKQ**NG**KIEHLKSPELNVDL
N. cinerea V1.110 *fhbp* FDKLPEGGRATYRGTAFGSDDAGGKLTYTIDFAAKQ**HG**KIEHLKSPELNVDL

N. meningitidis V1.1 *fhbp* AAADIKP**DGKR**HAVISGSVLYN**Q**AEKGSYSLGIFGGKAQEVAGSAEVEKTVNGI
N. cinerea V1.110 *fhbp* AAADIKP**DEKH**HAVISGSVLYN**Q**DEKGSYSLGIFGGKAQEVAGSAEVEKTVNGI

N. meningitidis V1.1 *fhbp* RHIGLAAKQ
N. cinerea V1.110 *fhbp* RHIGLAAKQ

B




FIG 1 *N. cinerea* fhbp is predicted to be structurally similar to V1.1 fhbp. (A) Sequence alignment of *N. meningitidis* V1.1 *fhbp* and *N. cinerea* V1.110 *fhbp*. The blue residues show sequence differences. (B) Predicted structure of *N. cinerea* V1.110 fhbp (gray ribbon) generated with the I-TASSER server and V1.1 fhbp (blue ribbon [25]; PDB accession no. 2W80). The figure was drawn using Pymol. Residues in red (shown as ball-and-stick representations) are different in *N. cinerea* fhbp and V1.1 fhbp. Residues R193 and E267 (red) are in close proximity to the CFH (green ribbon) binding site.

tated *fhbp* sequences were verified by BLAST using the V1.1 *fhbp* nucleotide sequence as the search query. All the isolates contain *fhbp* with a predicted signal peptide, a conserved lipobox, and a variable linker sequence (see Fig. S1 in the supplemental material), with a high degree of nucleotide sequence identity (~95%) with V1.1 *fhbp* from *N. meningitidis*, consistent with previous findings (37). Furthermore, *N. cinerea* CCUG 346T *fhbp* is identical to the meningococcal *fhbp* peptide variant V1.110 (<http://pubmlst.org/neisseria/>). We further characterized *fhbp* from *N. cinerea* using CCUG 346T, as the strain has previously been shown to be genetically tractable (43). The predicted amino acid sequence of fhbp from *N. cinerea* CCUG 346T has six amino acid differences from V1.1 fhbp (Fig. 1A). A structural model of *N. cinerea* fhbp was generated using V1.1 fhbp as the threading template (25; Protein Data Bank [PDB] accession no. 2W80) (Fig. 1B). One different amino acid, A58, is predicted to reside in the leader peptide, while three of the residues unique to *N. cinerea* fhbp (H243, H269, and D282) are in the C-terminal barrel of fhbp, distant from the CFH-fhbp interface (44). Residues R193 and E267 in *N. cinerea* fhbp reside at the CFH-fhbp interface in close proximity (i.e., within 4 Å) to S366 and N338 of CFH, respectively, which could potentially lead to the formation of additional salt bridges between fhbp and CFH. Alanine substitution for Q193 in meningococcal V1.1 fhbp leads to a 2-fold decrease in CFH binding compared to wild-type V1.1 fhbp, while alanine substitution at position 267 in V2.21 fhbp (E267) showed significantly reduced CFH binding, with no binding of CFH₆₋₇ detected (44).




FIG 2 *N. cinerea* V1.110 fHbp binds CFH at high affinity. (A) Recombinant V1.110 fHbp from *N. cinerea* (*N.c*) CCUG 346T, with V1.1 fHbp, and V1.1^{I331A} fHbp from *N. meningitidis* (*N.m*) were recognized by Western blotting using polyclonal anti-V1.1 fHbp sera. *N. cinerea* V1.110 fHbp and V1.1 fHbp bound CFH by far-Western analysis with NHS as the source of CFH. Molecular masses are shown, and Coomassie blue staining of proteins is shown as the loading control. (B) SPR analysis of *N. cinerea* fHbp binding to recombinant CFH₆₋₇ (concentrations are indicated) with 1:1 Langmuir fit (black lines). (C) Calculated K_D values for CFH₆₋₇ binding to *N. cinerea* V1.110 fHbp and V1.1 fHbp; NBD, no binding detected. (D) ELISA of *N. cinerea* fHbp, V1.1 fHbp, and V1.1^{I311A} fHbp binding to full-length CFH. (E) Significance of CFH binding by ELISA analyzed at $5 \mu\text{g}/\text{ml}$ of CFH. The error bars indicate the standard error of the mean (SEM) from three independent experiments, and *P* values (***, $P < 0.001$; ns, $P > 0.05$) were calculated using a two-tailed unpaired *t* test.

***N. cinerea* fHbp binds CFH at high affinity.** As amino acids in *N. cinerea* and *N. meningitidis* fHbp proteins differ at the CFH interface, we next evaluated whether *N. cinerea* fHbp binds CFH (37). *N. cinerea* fHbp was expressed as a recombinant protein and purified as previously described (44). The protein was recognized by anti-V1.1 fHbp serum by Western blot analysis, as expected (Fig. 2A). Initially, binding of fHbp from *N. cinerea* and V1.1 fHbp from *N. meningitidis* to CFH was determined by far-Western analysis using normal human serum (NHS) as the source of CFH. V1.1^{I311A} fHbp was used as a negative control, as it has significantly impaired binding to CFH (44). The results of far-Western analysis demonstrated that *N. cinerea* fHbp and wild-type V1.1 fHbp bind CFH, while V1.1^{I311A} fHbp has no detectable CFH binding, as previously described (44) (Fig. 2A).

To further characterize this interaction, we determined the affinity of *N. cinerea* fHbp for CFH₆₋₇ by surface plasmon resonance (SPR). *N. cinerea* fHbp binds to CFH₆₋₇ with an affinity similar to that of V1.1 fHbp, with equilibrium binding constants (K_D s) of 3.5 ± 0.5 nM and 3.0 ± 0.5 nM, respectively (Fig. 2B and C), indicating that the amino acid substitutions in *N. cinerea* V1.110 fHbp do not affect its ability to bind CFH. We also




FIG 3 Functional fHbp was expressed by all *N. cinerea* strains examined. (A) Western blot analysis of fHbp expression using anti-V1.1 fHbp sera against *N. cinerea* strains and *N. meningitidis* H44/76 and H44/76ΔfHbp. Far-Western analysis demonstrated that fHbps from different *N. cinerea* isolates bound CFH. RecA was used as a loading control. (B) Western and far-Western analyses demonstrating that binding of CFH to *N. cinerea* is dependent on fHbp expression. (C) Temperature-dependent regulation of fHbp expression by *N. cinerea*, examined by Western blot analysis using anti-V1.1 fHbp sera. RecA, loading control, with relative fHbp expression levels (rel. values) shown.

determined the levels of CFH binding to *N. cinerea* fHbp and V1.1 fHbp by enzyme-linked immunosorbent assay (ELISA) using full-length CFH (Fig. 2D and E). The ELISA data were consistent with results from both far-Western and SPR analyses, with no significant difference in CFH binding to *N. cinerea* fHbp and V1.1 fHbp over a range of concentrations ($P > 0.05$ by two-tailed t test) (Fig. 2E). Taken together, these results demonstrate that *N. cinerea* fHbp binds CFH and that this interaction is mediated by CFH₆₋₇ of CFH.

***N. cinerea* isolates express fHbp on the bacterial surface.** Next, the expression of fHbp by other *N. cinerea* isolates was analyzed using polyclonal mouse serum raised against V1.1 fHbp, which cross-reacts with *N. cinerea* fHbp from CCUG 346T (Fig. 2A). A band of approximately 27 kDa corresponding to the molecular mass of fHbp was detected in lysates of all five *N. cinerea* isolates by Western blotting (Fig. 3A). Moreover, five *N. cinerea* isolates were analyzed for the ability to bind full-length CFH present in NHS; *N. meningitidis* H44/76 and H44/76ΔfHbp were used as positive and negative controls, respectively. Far-Western analysis detected a band corresponding to fHbp in whole-cell lysates of *N. cinerea* isolates and the positive control, H44/76; no CFH binding to *N. meningitidis* H44/76ΔfHbp was detected, as previously described (39) (Fig. 3A). We also sought to establish whether CFH binding to *N. cinerea* is dependent on fHbp.

Therefore, we constructed an *fhbp* mutant of *N. cinerea* CCUG 346T by insertional inactivation of the gene with a kanamycin resistance cassette. Analysis of the wild-type *N. cinerea* and the $\Delta fhbp$ strain demonstrated that the $\Delta fhbp$ mutant did not express fHbp, as expected, while far-Western blot analysis demonstrated that CFH binding to *N. cinerea* is dependent on fHbp (Fig. 3B).

Our previous work demonstrated that *N. meningitidis* fHbp is subject to thermal regulation, with higher levels of fHbp produced by strains grown at higher temperatures (i.e., 37°C and 42°C compared with 30°C) (45, 46). Therefore, we investigated whether *N. cinerea* fHbp is also thermoregulated. *N. cinerea* CCUG 346T was harvested following growth of the bacteria in brain heart infusion (BHI) broth to mid-log phase at 30°C, 37°C, or 42°C; the relative fHbp expression levels were quantified by Western blot analysis (Fig. 3C). The results demonstrated that increasing temperature is associated with elevated fHbp expression in *N. cinerea*, similar to the meningococcus ($P = 0.0382$; one-way analysis of variance [ANOVA] for 30°C versus 42°C) (Fig. 3C). Inspection of sequences corresponding to the 5' untranslated region (UTR) of *fhbp* from *N. cinerea* CCUG 346T indicated that there is a single base change (C-U) in the fumarate-nitrate reduction regulator (FNR) box, but the temperature-regulatory domains identified in *N. meningitidis* MC58 (46) are conserved in *N. cinerea* CCUG 346T (see Fig. S2 in the supplemental material). Two putative α -ribosome binding sites (α -RBS) in the open reading frame (ORF) of fHbp of *N. meningitidis* MC58 have been shown to determine temperature regulation of fHbp (46), and *N. cinerea* CCUG 346T retains an identical sequence for α -RBS1 (CUGCCU) (see Fig. S2 in the supplemental material) but has a C-U substitution in the second α -RBS.

To analyze surface expression of fHbp in *N. cinerea*, we performed flow cytometry analysis using anti-V1.1 fHbp serum to detect fHbp on the surfaces of *N. cinerea* CCUG 346T and CCUG 346T $\Delta fhbp$ (Fig. 4A and B); *N. meningitidis* H44/76 and H44/76 $\Delta fhbp$ were included as positive and negative controls, respectively (Fig. 4C and D). The results demonstrated that fHbp is detected on the surface of *N. cinerea* CCUG 346T, as well as *N. meningitidis* H44/76; fHbp was not detected on either of the *fhbp* mutants (two-tailed *t* test; $P < 0.0001$ and $P = 0.0183$ compared with the respective wild-type strains).

***N. cinerea* fHbp promotes complement resistance.** Next, we investigated whether fHbp can recruit CFH to the surface of *N. cinerea*. *N. cinerea* CCUG 346T and CCUG 346T $\Delta fhbp$ were incubated in heat-inactivated (HI) NHS as a source of CFH, and CFH was detected by flow cytometry with an α -CFH monoclonal antibody (MAb), OX24 (Fig. 5A and B) (47); binding was quantified as the geometric mean fluorescence intensity. Bacteria incubated in phosphate-buffered saline (PBS) or in CFH-depleted serum (HI-CFH depleted) were used as controls. Following incubation with HI-NHS, CFH was detected on the surface of *N. cinerea* (Fig. 5A and B), while no CFH binding was found either after incubating wild-type bacteria with HI-CFH-depleted serum or PBS or after incubating the *fhbp* mutant in HI-NHS (two-tailed *t* test; $P = 0.0002$) (Fig. 5A to D), demonstrating that CFH binds to the surface of *N. cinerea* in an fHbp-dependent manner.

Next, we examined the functional consequences of CFH binding to *N. cinerea* by performing serum survival assays. *N. cinerea* CCUG 346T and CCUG 346T $\Delta fhbp$ were incubated in NHS (concentrations ranging from 0 to 20%) for 30 min, and the percent survival was calculated by comparing the number of CFU recovered after incubation in serum with the number of CFU recovered after incubation in PBS. *N. cinerea* CCUG 346T was significantly more resistant to complement-mediated lysis than CCUG 346T $\Delta fhbp$ over a range of serum concentrations (Fig. 6A). For example, there was 54% and 31% survival for CCUG 346T and CCUG 346T $\Delta fhbp$, respectively, in 5% NHS (two-tailed *t* test; $P = 0.0161$). Furthermore, we examined whether these differences in survival were observed in the presence of only the AP of complement by inhibiting the activities of the classical and lectin pathways by the addition of 10 mM EGTA and 5 mM MgCl₂ to sera (48). *N. cinerea* CCUG 346T still demonstrated a trend toward enhanced serum resistance compared to CCUG 346T $\Delta fhbp$ when only




FIG 4 *N. cinerea* fHbp is present on the bacterial surface. Flow cytometry analysis with anti-V1.1 fHbp serum demonstrated detection of fHbp on the surfaces of *N. cinerea* (solid black line) (A and B) and *N. meningitidis* (solid blue line) (C and D). The error bars indicate the standard error of the mean (SEM) from three independent experiments. Significance was calculated using a two-tailed unpaired *t* test; ****, $P < 0.0001$; *, $P < 0.05$.

the AP was active in a 15% serum concentration, although this did not reach statistical significance (two-tailed *t* test; $P = 0.127$) (52.28% and 38.94% survival for CCUG 346T and CCUG 346TΔfHbp, respectively) (Fig. 6B). Furthermore, we analyzed whether an *N. meningitidis* fHbp mutant, H44/76ΔfHbp, could be complemented with *N. cinerea* fHbp (V1.110). *N. meningitidis* H44/76, H44/76ΔfHbp, and H44/76ΔfHbp complemented with either meningococcal V1.1 fHbp or *N. cinerea* fHbp were incubated in NHS (concentrations ranging from 0 to 75%) for 30 min, and the percent survival was calculated. Both complemented strains, H44/76ΔfHbp::V1.1fHbp and H44/76ΔfHbp::V1.110fHbp, were significantly more resistant to NHS than H44/76ΔfHbp (one-way ANOVA; $P = 0.0023$ and $P = 0.0046$, respectively), and no significant difference in survival was observed between the complemented strains and H44/76 at 75% NHS (Fig. 6C). Taken together, these results demonstrate that *N. cinerea* fHbp contributes to the evasion of killing by the complement system.

fHbp-containing vaccines elicit SBA against *N. cinerea*. Finally we examined whether the licensed meningococcal vaccine, Bexsero, which contains meningococcal fHbp, elicits immune responses against *N. cinerea*. Mice were immunized with Bexsero or an equivalent amount (20 μg) of V1.1 fHbp; sera from mice immunized with adjuvant alone were used as a negative control. Initially, we used ELISA to measure the ability of antibodies in pooled immune sera to recognize V1.1 fHbp and *N. cinerea* fHbp. Interestingly, there was no significant difference in the recognition of V1.1 fHbp and *N. cinerea* fHbp by antibodies in sera from mice immunized with Bexsero or V1.1 fHbp (Fig. 7A; data from mice immunized with fHbp V1.1 not shown) ($P > 0.05$ by two-tailed *t* test). Next, the reactivity of sera was examined by Western blotting of whole-cell lysates




FIG 5 Surface-bound CFH contributes to complement resistance of *N. cinerea* in an fHbp-dependent manner. Shown is flow cytometry analysis of CFH binding to *N. cinerea* CCUG 346T (A and B) and *N. cinerea* CCUG 346TΔfHbp (C and D). The shaded areas indicate results from bacteria incubated with PBS, the solid green lines show bacteria incubated in HI-NHS, and the dashed green lines show bacteria incubated in HI-CFH-depleted sera. The data are presented as the mean fluorescence intensity. The error bars indicate the standard error of the mean (SEM) from three independent experiments, and the *P* values were calculated using a two-tailed unpaired *t* test; ***, *P* < 0.001; ns, *P* > 0.05.

from *N. cinerea* CCUG 346T and *N. meningitidis* H44/76 (Fig. 7B). Sera from mice immunized with V1.1 fHbp detected a band of the size expected for fHbp by Western blot analysis of lysates of wild-type strains, while no protein was detected in the ΔfHbp mutants, as previously observed for polyclonal sera against V1.1 fHbp (see Fig. S3A in the supplemental material) (44). As expected, sera from mice immunized with Bexsero detected multiple protein bands in whole-cell lysates (see Fig. S3B) due to the complex mixture of the vaccine (23), although a band corresponding to the molecular mass of fHbp was detected in lysates of *N. cinerea* CCUG 346T, but not from the ΔfHbp mutant. Sera from mice immunized with either recombinant fHbp or Bexsero also recognized recombinant V1.1 and *N. cinerea* V1.110 fHbp (Fig. 7B).

Next, the potential impact of immune responses raised against Bexsero on *N. cinerea* was examined by measuring SBA. Immunization with V1.1 fHbp or Bexsero elicited SBA titers of 128 against *N. cinerea* CCUG 346T; SBA titers of ≥8 are correlated with protection against serogroup B meningococcal disease (Fig. 7C) (49). Of note, sera from mice immunized with Bexsero had a higher SBA titer (256) against *N. cinerea* CCUG 346T complemented with *N. cinerea* fHbp under an IPTG (isopropyl-β-D-thiogalactopyranoside)-inducible promoter than wild-type bacteria, consistent with higher expression of fHbp by the complemented strain (not shown). Furthermore, no SBA was detected against *N. cinerea* CCUG 346TΔfHbp with either V1.1 fHbp or Bexsero, demonstrating that SBA elicited by Bexsero against *N. cinerea*




FIG 6 *N. cinerea* fHbp contributes to complement resistance. (A and B) Sensitivity of *N. cinerea* and *N. cinerea* CCUG 346TΔ*fHbp* to complement-mediated lysis in the presence of NHS (A) and in the presence of the AP of complement only (B). (C) Survival in human serum of *N. meningitidis* H44/76, H44/76Δ*fHbp*, and H44/76Δ*fHbp* complemented with either V1.1 fHbp or *N. cinerea* V1.110 fHbp. The error bars indicate the standard error of the mean (SEM) from three independent experiments.

is mediated by fHbp. In conclusion, immune responses elicited by Bexsero recognize *N. cinerea* fHbp and have significant SBA.

DISCUSSION

fHbp is a key virulence factor for the meningococcus and binds the negative complement regulator CFH, promoting bacterial survival in human serum (25, 50, 51). The importance of fHbp as a vaccine antigen is evident from its inclusion in two recently licensed serogroup B meningococcal vaccines (21, 22). Here, we show that fHbp expression is not restricted to pathogenic *N. meningitidis* but that the antigen is also expressed by the commensal species *N. cinerea*, in which fHbp is surface located and binds CFH at high affinity. Recruitment of CFH by *N. cinerea* promotes its survival




FIG 7 Immunization with Bexsero elicits SBA against *N. cinerea*. (A) ELISA analysis of anti-Bexsero mouse sera (3-fold dilutions from 1/300) against recombinant fHbp; BSA was used as a control. The error bars indicate the standard error of the mean (SEM) from three independent experiments. (B) fHbp (~27 kDa) is recognized by sera from mice immunized with V1.1 fHbp or Bexsero as determined by Western blot analysis. The higher-molecular-mass band observed when whole-cell extracts were probed with sera from mice immunized with Bexsero was likely due to responses against other proteins in the vaccine. (C) Serum bactericidal titers of pooled sera from mice immunized with recombinant V1.1 fHbp or Bexsero against *N. cinerea* strains; NBT, no bactericidal titer.

in human serum complement, while serum raised against Bexsero has SBA against *N. cinerea* expressing fHbp.

Previous work has analyzed the expression and function of fHbp in pathogenic *Neisseria* spp. (39, 50, 51). fHbp is a highly variable protein that can be grouped into three distinct variants based on its amino acid sequence (26). Using SPR, we found that *N. cinerea* fHbp binds CFH in the nanomolar range, similar to the affinity of CFH for V1.1 fHbp from *N. meningitidis*, indicating that high-affinity binding of CFH to the surface of the bacterium is not only involved during survival in the bloodstream, but likely aids colonization and persistence in the human nasopharynx. In contrast, *N. gonorrhoeae*, which resides in the genitourinary tract, expresses a nonfunctional fHbp called Ghfp (39) and instead recruits CFH via an abundant surface porin, PorB.1 (52, 53). A frameshift mutation in the nucleotide sequence at position 40 of *ghfp* disrupts the lipobox, resulting in the loss of surface localization of Ghfp (39), and sequence changes in *ghfp* result in a protein that is unable to bind CFH (39). The distinct mechanisms by which *N. meningitidis* and *N. cinerea* recruit CFH compared with *N. gonorrhoeae* are likely to reflect differences in concentrations of CFH and the levels of other complement proteins at the sites they occupy in the body. Complement proteins, including C3b, may be derived from exudates on the nasopharyngeal mucosa from the subepithelial microvasculature (54, 55). Levels of C3 in nasopharyngeal secretions are estimated to be approximately 6.6% of serum C3 levels (range, 4.2 to 20.2%), indicating that there is a significant level of complement in the nasopharynx (56). Although CFH could not be detected on respiratory epithelial cells by immunohistochemistry (57), levels of the complement regulator in saliva range from 0.007 to 0.059 $\mu\text{g/ml}$ (0.04 to 0.38 nM) (58, 59), which is significantly lower than in serum. There are few studies on CFH in the genital tract, although low levels of CFH have been detected in urothelial cells, which also can produce C3 (57). High-affinity interactions with CFH might be required at mucosal surfaces due to low availability of this important complement regulator (58, 59). Furthermore, levels of CFH and other complement proteins may change during

inflammation, when CFH recruitment might provide an advantage for commensal and pathogenic bacteria in avoiding exudates of serum components and putatively could aid *N. cinerea* colonization of the nasopharynx (56, 60, 61).

Binding of CFH is important for nasopharyngeal colonization by other bacteria, including *Streptococcus pneumoniae* (62), by mediating interactions with respiratory epithelial cells. Binding of CFH to PspC on the pneumococcus significantly increases its attachment to human epithelial and endothelial cells by acting as a molecular bridge between the bacteria and host cells (63). Moreover, the RGD motif in CCP₄ of CFH can recognize integrins on cells to induce uptake of the pneumococcus (63, 64). A similar role for CFH acting as a molecular bridge has not been described for *N. meningitidis* or *N. cinerea* as yet and merits further investigation, as similar mechanisms might influence *N. meningitidis* and *N. cinerea* colonization of the human nasopharynx.

Although *N. cinerea* expresses *fHbp*, other, closely related commensal species, such as *N. lactamica* and *N. polysaccharea*, do not express fHbp. The majority of *N. polysaccharea* isolates have a frameshift in the ORF, potentially producing a truncated protein, while *fHbp* has not been identified in the *N. lactamica* genome, indicating that fHbp is not essential for successful colonization of the human nasopharynx by *Neisseria* spp.

Previous work has suggested that NspA and PorB2 of *N. meningitidis* can bind CFH, although these proteins have little effect in wild-type strains and specific binding affinities are not known (65, 66). Although *N. cinerea* CCUG 346T carries a gene encoding PorB2, we found no evidence by Western blotting or flow cytometric analysis that molecules other than fHbp are responsible for *N. cinerea* binding to CFH. Of note, *N. cinerea* CCUG 346T is not predicted to harbor a functional *nspA* gene. Other functional roles that could enhance bacterial survival have also been attributed to meningococcal fHbp, including protecting against the cationic antimicrobial peptide LL-37 (67). LL-37 is a cathelicidin that is expressed by epithelial cells in the nasopharynx (68, 69). Furthermore, both fHbp and Ghfp have been postulated to act as siderophore receptors and to recruit enterobactin and so could contribute to iron scavenging on mucosal surfaces (70).

Previously, we demonstrated that meningococcal fHbp levels are elevated with increasing temperatures, i.e., from 30°C to 37 and 42°C (45, 46). Thermoregulation of meningococcal fHbp is mediated by two putative α -RBS within the ORF of *fHbp*; at lower temperatures, the transcript forms a stem-loop structure including the RBS, thus reducing fHbp expression (46, 71). Sequence analysis of *N. cinerea* CCUG 346T indicated that the 5' UTR upstream of *fHbp*, including the monocistronic promoter, is conserved compared with *N. meningitidis* MC58 and includes the RBS and one repeat of the α -RBS within the *fHbp* ORF (see Fig. S2 in the supplemental material). The second α -RBS has a single residue change, C-U, although *N. cinerea* fHbp is still subject to temperature regulation (Fig. 3C). Temperature is also an indicator of local inflammation, which could expose the bacterium to host immune effectors. Therefore, upregulation of fHbp could provide *N. cinerea* with a selective advantage over other commensal *Neisseria* spp. that do not express the antigen (72, 73).

Conjugate vaccines against *N. meningitidis* have been highly successful at reducing carriage of the pathogen but do not affect commensal *Neisseria* spp., as they target specific capsular polysaccharides that are not expressed by commensal species (74, 75). Limited data are currently available on the impact of Bexsero and other protein-based vaccines on meningococcal carriage (76, 77). As IgG and complement are present on mucosal surfaces (54, 78, 79), it is possible that immune responses against fHbp could affect the nasopharyngeal carriage of bacteria expressing fHbp. We found that immunization of mice with Bexsero elicited SBA against *N. cinerea*, with titers similar to those found in mice immunized with fHbp alone. Furthermore, there was no SBA against *N. cinerea* lacking fHbp, indicating that fHbp is largely responsible for the SBA seen with Bexsero. With regard to other protein antigens found in Bexsero, genes encoding GNA2091 and GNA1030 are present in all *N. cinerea* isolates, whereas only fragments of *nadA* are found in some isolates (37). NHBA and PorA have not been identified in any *N. cinerea* isolate analyzed to date. However, as *N. cinerea* lacks a capsule, it may be more susceptible than *N. meningitidis* to immune responses elicited by Bexsero against

TABLE 1 *N. cinerea* and *N. meningitidis* strains used in this study

Strain	fHbp variant	Yr identified	Reference
<i>N. cinerea</i>			
CCUG 346T	V1.110	Unknown	7
CCUG 346TΔfHbp			This study
CCUG 346TΔfHbp::fHbp	V1.110		This study
CCUG 25879	V1.564	1989	7
CCUG 27178 A	V1.110	1983	7
CCUG 53043	V1.534	2006	85
CCUG 5746	V1.563	1977	7
<i>N. meningitidis</i>			
H44/76	V1.1	1976	86
H44/76ΔfHbp			39
H44/76ΔfHbp::V1.1 fHbp	V1.1		39
H44/76ΔfHbp::V1.110 fHbp	V1.110		This study

surface proteins. Therefore, it will be important to evaluate the impact of Bexsero on the carriage of commensal *Neisseria*, as carriage of nonpathogenic species can induce cross protective immunity (5, 42, 80).

In summary, we have shown that *N. cinerea* expresses functional fHbp on its surface, where it contributes to complement evasion. Moreover, Bexsero elicits bactericidal antibodies against *N. cinerea*, with SBA responses similar to immunization with fHbp alone. Therefore, implementation of vaccines containing fHbp has the potential to affect the commensal nasopharyngeal flora, and the impact of multicomponent vaccines should be assessed in future challenge studies.

MATERIALS AND METHODS

Sequence analysis and model of *N. cinerea* fHbp. *N. cinerea* fHbp sequences were extracted from previously annotated whole-genome sequences available in the PubMLST BIGSdb database (<http://pubmlst.org/neisseria/>). Alignments of the predicted *N. cinerea* fHbp sequence against the V1.1 fHbp were generated by Clustal OMEGA (<http://www.ebi.ac.uk/Tools/msa/clustalo/>); residues were numbered according to V1.1 fHbp. A model of *N. cinerea* fHbp was generated using the iTASSER server (81) with V1.1 fHbp (25; PDB accession no. 2W80). The resulting model is illustrated in PyMol (<https://www.pymol.org/>) and has high confidence, with a C score of 1.68.

Bacterial strains and growth. The strains used in this study are listed in Table 1. *Neisseria* species were grown on BHI agar (1.5% [wt/vol]; Oxoid) supplemented with 5% defibrinated horse blood. Bacteria grown on solid media were incubated overnight at 37°C in the presence of 5% CO₂. Liquid cultures in BHI broth were inoculated with 10⁹ CFU per 10 ml and grown with shaking at 180 rpm at the specified temperatures to an A₆₀₀ of ~0.5 unless otherwise indicated. Antibiotics were added to media at the following concentrations: kanamycin, 100 μg ml⁻¹, and erythromycin, 10 μg ml⁻¹. *Escherichia coli* was grown in Luria-Bertani (LB) broth or on LB agar. Overnight liquid cultures were grown at 37°C with shaking at 180 rpm in 5 ml of medium inoculated from a single colony. The bacteria were diluted 1 in 100 the next morning into larger volumes for protein expression. Antibiotics were added at the following concentrations for *E. coli*: kanamycin, 50 μg ml⁻¹, and carbenicillin, 100 μg ml⁻¹.

Generation of plasmids and protein purification. Genomic DNA was isolated using a Wizard Genomic DNA purification kit (Promega) according to the manufacturer's instructions. fHbp was amplified from *N. cinerea* genomic DNA using primers F_{fHbp} (5'-GCCATATGATGGCCGCCGACAT-3'; NdeI site underlined) and R_{fHbp} (5'-GCCTCGAGTTGCTTGGCGCAAGGCCGATAT-3'; XhoI site underlined). The product was ligated into pET-21b (Table 2), resulting in a construct for expression of fHbp with a C-terminal His tag under an inducible promoter. fHbp expression was performed as described previously (44).

In brief, for protein expression, *E. coli* BL21(D3) (Agilent) containing fHbp expression constructs was grown in liquid medium to an A₆₀₀ of 0.4 to 0.8; then, IPTG was added to a final concentration of 1 mM. After 4 h, the bacteria were harvested by centrifugation at 5,000 × g for 30 min at 4°C prior to cell lysis in an EmulsiFlex-C5 homogenizer (Avestin) at 15,000 lb/in². The lysates were centrifuged at 50,000 × g for 30 min at 4°C, and recombinant fHbp was purified by affinity chromatography using a HisTrap column (GE Healthcare) and eluted with 200 mM imidazole. Further purification was performed with an AKTA

TABLE 2 *E. coli* BL21(DE3)Lys plasmids

Plasmid	Insert	Modification	Source
pET21b	V1.1 fHbp		44
pET21b	V1.1 fHbp	I311A	44
pET21b	V1.110 fHbp		This study

purifier (GE Healthcare) by anion-exchange chromatography (HiTrapQ HP column; GE Healthcare). Protein concentrations were estimated using a Nanodrop 2000c spectrophotometer (Thermo Scientific).

SDS-PAGE and Western blot analysis. Recombinant protein (10 μ g) and whole-cell extracts were separated on 12% or 14% polyacrylamide gels prior to transfer to Immobilon P polyvinylidene difluoride (PVDF) membranes (Millipore, USA) using the Trans-Blot SD semidry transfer system (Bio-Rad, USA) or staining for 10 min with Coomassie blue. For whole-cell extracts, *N. meningitidis* and *N. cinerea* were grown on solid medium overnight and resuspended in PBS. The number of bacteria was determined by measuring the DNA concentration by the A_{260} of 20 μ l of the bacterial suspension in 980 μ l of lysis buffer (0.1 M NaOH, 1% SDS); 10^9 CFU was mixed with an equal volume of 2 \times SDS-PAGE loading buffer (100 mM Tris-HCl, pH 6.8, 20 μ M β -mercaptoethanol, 4% SDS, 0.2% bromophenol blue, 20% glycerol) and boiled for 10 min. For Western blot analysis, membranes were blocked in 0.05% (wt/vol) dry milk-PBS with 0.05% (vol/vol) Tween 20 overnight and then incubated with primary antibodies or with NHS as a source of CFH. Primary antibodies were used at the following concentrations: anti-V1.1 fHbp polyclonal sera, 1:1,000 (44); anti-RecA, 1:5,000 (Abcam, Cambridge); anti-CFH, 1:10,000 (Millipore). Horseradish peroxidase (HRP)-conjugated secondary antibodies, goat anti-mouse HRP (Dako, United Kingdom), rabbit anti-goat HRP (Dako, United Kingdom), and goat anti-rabbit HRP (Santa Cruz Biotechnology, Germany), were used at a final concentration of 1:10,000. Membranes were washed three times in PBS with 0.05% (vol/vol) Tween 20, and binding was detected using an ECL Western blotting detection kit (Amersham, USA).

To examine the effect of temperature on fHbp expression, *N. cinerea* was grown on solid medium overnight at 37°C prior to inoculating BHI broth (10 ml) with 10^9 bacteria. Cultures were grown to mid-log phase (A_{600} ~0.5) at 30°C, 37°C, or 42°C. Bacteria were collected by centrifugation (14,000 \times g for 5 min at room temperature) and then resuspended in 100 μ l sample buffer prior to analysis by Western blotting. Relative expression values were calculated by measuring band intensities with ImageJ software; normalized to the signal for RecA, the loading control; and shown as a ratio relative to bacteria grown at 30°C.

Surface plasmon resonance and ELISA. SPR was performed using a BiaCore 3000 instrument (GE Healthcare). The purified proteins were immobilized on a CM5 sensor chip to approximately 800 reaction units (RU) by amine coupling (GE Healthcare). Unreacted groups on the chip were inactivated with ethanolamine. HSB-EB (HSB buffer; 0.01 M HEPES, 150 mM NaCl, 3 mM EDTA, pH 7.4) was used throughout. Increasing concentrations of CFH₆₋₇, prepared as described previously (82) (0.5 nM to 32 nM), were injected over the chip at a flow rate of 40 μ l/min using the KINJECT command with a dissociation time of 300 s. K_D values were calculated and analyzed with BIAevaluation software. Kinetic data were referenced against both a blank cell and subtraction of a blank injection (HSB-EB).

To determine CFH binding by ELISA, 96-well plates (F96 MaxiSorp; Nunc) were coated with recombinant fHbp (3 μ g/ml; 50 μ l per well) overnight at room temperature prior to blocking with 3% bovine serum albumin (BSA) in PBS with 0.05% Tween 20. The plates were incubated with full-length human CFH (Sigma) in 2-fold dilutions from 10 μ g/ml, and binding was detected with anti-CFH antibody (Millipore) and HRP-conjugated goat anti-mouse polyclonal antibody (Dako). CFH binding was visualized with 3,3',5,5'-tetramethylbenzidine (TMB) substrate reagent (Roche) and 2 N sulfuric acid stop solution (Roche) according to the manufacturer's instructions, and the A_{450} was measured (SpectraMax M5; Molecular Devices). Statistical significance was tested using an unpaired Student *t* test (GraphPad Prism v.6.0) to compare means and standard deviations (SD), with a *P* value of <0.05 as the cutoff for significance. To analyze the reactivity of sera from mice immunized with Bexsero, ELISA plates were coated with V1.1 fHbp, V1.110 fHbp, or BSA (2.5 μ g/ml; 50 μ l per well) prior to incubation with sera; antibody binding was detected as described above. Pooled mouse sera were used as a primary antibody with 3-fold dilutions from a starting dilution of 1:300.

Generation of *N. cinerea* fHbp mutants and complemented strains. *N. cinerea* strain CCUG 346T Δ fHbp was constructed by insertional inactivation of the *fHbp* gene with a cassette encoding kanamycin resistance. *Nc* Δ fHbp (forward primer; 5'-GCCATATGGCTGAAAGTCATCAACGAAT-3' [NdeI site underlined]) and *Nc* Δ fHbp (reverse primer; 5'-GCCTCGAGGTGCAATACAAAATGCCGTCCGAACGGTAA-3' [XhoI site underlined]) were used to amplify *fHbp* and flanking sequences from *N. meningitidis* strain H44/76 Δ fHbp. The flanking sequences between *N. meningitidis* H44/76 and *N. cinerea* CCUG 346T shared 99.28% nucleic acid sequence identity. *N. cinerea* CCUG 346 T was then transformed with a linear PCR fragment as described previously (83). The complemented *N. cinerea* and *N. meningitidis* strains CCUG 346T Δ fHbp::fHbp and H44/76 Δ fHbp::V1.110 were constructed by amplifying *fHbp* from *N. cinerea* genomic DNA using primers *Nc* fHbp forward (5'-CGGTTAATTAAGGAGTAATTTTGTGAACC-3'; PacI site underlined) and *Nc* fHbp reverse (5'-CGGTTAATTAATTTGCTTGGCGGC-3'; PacI site underlined). The PCR products were ligated into pNCC1 for transformation into *N. cinerea* (43) or PGCC4 for transformation into *N. meningitidis* (84), which were linearized by ClaI digestion before transforming into *N. cinerea* or *N. meningitidis* as described previously (83).

Flow cytometry. *N. meningitidis* and *N. cinerea* were grown on solid medium overnight and quantified as described above. The bacteria (10^9 CFU) were fixed in 3% paraformaldehyde for 2 h at room temperature and then washed with PBS. fHbp was detected using anti-V1.1 fHbp serum and goat anti-mouse IgG Alexa Fluor 647-conjugated polyclonal antibody (Molecular Probes, Life Technologies). To evaluate CFH binding, 5×10^7 CFU was resuspended in 20 μ l of HI-NHS or CFH-depleted serum (CompTech, USA) for 30 min at room temperature. Binding of CFH was detected with an anti-CFH MAb (OX24) and goat anti-mouse IgG Alexa Fluor 647-conjugated polyclonal antibody (Molecular Probes, Life Technologies) (47). Samples were analyzed using a FACSCalibur (BD Biosciences), and at least 10^4 events were recorded; the results were analyzed by calculating the geometric mean fluorescence intensity in

FlowJo vX software (Tree Star). Statistical significance was tested using an unpaired Student *t* test (GraphPad Prism v.6.0).

Serum sensitivity assays. NHS was obtained by collecting venous blood and allowing it to coagulate at room temperature for 60 min before centrifuging it at $3,000 \times g$ for 20 min at 4°C prior to storage at -80°C. The NHS was heat inactivated (HI-NHS) at 56°C for 30 min to inactivate complement. *N. cinerea* and *N. meningitidis* were grown overnight on BHI agar, and then 10^4 CFU was incubated in dilutions of NHS in PBS (0 to 20% [vol/vol] for *N. cinerea*; 0 to 75% [vol/vol] for *N. meningitidis*) for 30 min at 37°C in the presence of CO₂. To analyze the effect of the AP on bacterial survival, NHS was preincubated with 5 mM MgCl₂-10 mM EGTA to inhibit the classical and lectin pathways (48). Bacterial survival was determined by plating onto BHI agar in triplicate. The percent survival was calculated by comparing bacterial recovery in serum with recovery from samples containing no serum. Statistical significance was tested using an unpaired Student *t* test or one-way ANOVA (GraphPad Prism v.6.0).

Generation of immune sera and SBA. Eight female BALB/c mice (6 to 8 weeks old; Charles River, Margate, UK) were immunized with recombinant fHbp (20 μg) adsorbed to aluminum hydroxide [final composition, 0.5 mg/ml Al(OH)₃, 10 mM histidine-HCl] by mixing overnight at 4°C or with Bexsero (total protein, 20 μg). Antigens were given by the intraperitoneal route on days 0, 21, and 35, and sera were collected on day 49 by terminal anesthesia and cardiac puncture. All animal experiments were carried out under protocols reviewed and approved by the Home Office, United Kingdom, under license number PPL 30/3194.

To measure SBA, *N. cinerea* was grown overnight on BHI agar prior to replating onto solid medium and was grown for a further 5 h at 37°C in 5% CO₂. A total of 5×10^4 CFU/ml of *N. cinerea* was mixed with an equal volume of baby rabbit complement (Cedarlane) diluted 1:10 in SBA buffer (Dulbecco's PBS containing 0.1% [wt/vol] glucose, 1 mM CaCl₂, 0.5 mM MgCl₂). Sera from individual mice were pooled and heat inactivated for 1 h at 56°C prior to being added to wells in 2-fold dilutions starting at 1:8. Control wells contained either no serum or no complement. Following incubation for 1 h at 37°C, 10 μl from each well was plated onto BHI agar in triplicate, and the number of surviving bacteria was determined after overnight growth. The SBA was expressed as the reciprocal of the highest dilution of serum required to kill more than 50% of the bacteria relative to a no-serum control (49).

Statistical analysis. Statistical significance was calculated, using GraphPad Prism v.6.0, by unpaired Student *t* test or one-way ANOVA as indicated. A *P* value of <0.05 was considered statistically significant.

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/IAI.00305-17>.

SUPPLEMENTAL FILE 1, PDF file, 0.3 MB.

SUPPLEMENTAL FILE 2, PDF file, 0.1 MB.

SUPPLEMENTAL FILE 3, PDF file, 0.4 MB.

SUPPLEMENTAL FILE 4, PDF file, 0.3 MB.

ACKNOWLEDGMENTS

We are grateful to Rachel Exley for advice about experiments and the manuscript. We are grateful to Carina Brehony and Martin Maiden for providing *N. cinerea* strains and to Philip Ward and Susan Lea for providing purified CFH₆₋₇.

The work was supported by grants from Meningitis Now and the Wellcome Trust to C.M.T.

REFERENCES

- Knapp JS, Hook EW. 1988. Prevalence and persistence of *Neisseria cinerea* and other *Neisseria* spp. in adults. *J Clin Microbiol* 26: 896–900.
- Smith NH, Holmes EC, Donovan GM, Carpenter GA, Spratt BG. 1999. Networks and groups within the genus *Neisseria*: analysis of *argF*, *recA*, *rho*, and 16S rRNA sequences from human *Neisseria* species. *Mol Biol Evol* 16:773–783. <https://doi.org/10.1093/oxfordjournals.molbev.a026162>.
- Stephens DS, Greenwood B, Brandtzaeg P. 2007. Epidemic meningitis, meningococcaemia, and *Neisseria meningitidis*. *Lancet* 369:2196–2210. [https://doi.org/10.1016/S0140-6736\(07\)61016-2](https://doi.org/10.1016/S0140-6736(07)61016-2).
- Wong ES, Stamm WE. 1983. Urethral infections in men and women. *Annu Rev Med* 34:337–358. <https://doi.org/10.1146/annurev.me.34.020183.002005>.
- Gold R, Goldschneider I, Lepow ML, Draper TF, Randolph M. 1978. Carriage of *Neisseria meningitidis* and *Neisseria lactamica* in infants and children. *J Infect Dis* 137:112–121. <https://doi.org/10.1093/infdis/137.2.112>.
- Zaura E, Keijsers BJ, Huse SM, Crielaard W. 2009. Defining the healthy “core microbiome” of oral microbial communities. *BMC Microbiol* 9:259. <https://doi.org/10.1186/1471-2180-9-259>.
- Bennett JS, Jolley KA, Earle SG, Corton C, Bentley SD, Parkhill J, Maiden MC. 2012. A genomic approach to bacterial taxonomy: an examination and proposed reclassification of species within the genus *Neisseria*. *Microbiology* 158:1570–1580. <https://doi.org/10.1099/mic.0.056077-0>.
- Guibourdenche M, Popoff MY, Riou JY. 1986. Deoxyribonucleic acid relatedness among *Neisseria gonorrhoeae*, *N. meningitidis*, *N. lactamica*, *N. cinerea* and “*Neisseria polysaccharea*.” *Ann Inst Pasteur Microbiol* 137B:177–185.
- Liu G, Tang CM, Exley RM. 2015. Non-pathogenic *Neisseria*: members of an abundant, multi-habitat, diverse genus. *Microbiology* 161:1297–1312. <https://doi.org/10.1099/mic.0.000086>.
- Frasch CE, Zollinger WD, Poolman JT. 1985. Serotype antigens of *Neisseria meningitidis* and a proposed scheme for designation of serotypes. *Rev Infect Dis* 7:504–510. <https://doi.org/10.1093/clinids/7.4.504>.
- Pollard AJ, Levin M. 2000. Vaccines for prevention of meningococcal disease. *Pediatr Infect Dis J* 19:333–345. <https://doi.org/10.1097/00006454-200004000-00015>.

12. Jodar L, Feavers IM, Salisbury D, Granoff DM. 2002. Development of vaccines against meningococcal disease. *Lancet* 359:1499–1508. [https://doi.org/10.1016/S0140-6736\(02\)08416-7](https://doi.org/10.1016/S0140-6736(02)08416-7).
13. Harrison OB, Claus H, Jiang Y, Bennett JS, Bratcher HB, Jolley KA, Corton C, Care R, Poolman JT, Zollinger WD, Frascch CE, Stephens DS, Feavers I, Frosch M, Parkhill J, Vogel U, Quail MA, Bentley SD, Maiden MC. 2013. Description and nomenclature of *Neisseria meningitidis* capsule locus. *Emerg Infect Dis* 19:566–573. <https://doi.org/10.3201/eid1904.111799>.
14. Vipond C, Care R, Feavers IM. 2012. History of meningococcal vaccines and their serological correlates of protection. *Vaccine* 30(Suppl 2): B10–B17. <https://doi.org/10.1016/j.vaccine.2011.12.060>.
15. Kelly C, Arnold R, Galloway Y, O'Hallahan J. 2007. A prospective study of the effectiveness of the New Zealand meningococcal B vaccine. *Am J Epidemiol* 166:817–823. <https://doi.org/10.1093/aje/kwm147>.
16. Finne J, Leinonen M, Mäkelä PH. 1983. Antigenic similarities between brain components and bacteria causing meningitis. Implications for vaccine development and pathogenesis. *Lancet* ii:355–357.
17. Fredriksen JH, Rosenqvist E, Wedege E, Bryn K, Bjune G, Frøholm LO, Lindbak AK, Møgster B, Namork E, Rye U. 1991. Production, characterization and control of MenB-vaccine “Folkehelsa”: an outer membrane vesicle vaccine against group B meningococcal disease. *NIPH Ann* 14: 67–80.
18. Holst J, Martin D, Arnold R, Huergo CC, Oster P, O'Hallahan J, Rosenqvist E. 2009. Properties and clinical performance of vaccines containing outer membrane vesicles from *Neisseria meningitidis*. *Vaccine* 27(Suppl 2): B3–B12. <https://doi.org/10.1016/j.vaccine.2009.04.071>.
19. Tappero JW, Lagos R, Ballesteros AM, Plikaytis B, Williams D, Dykes J, Gheesling LL, Carlone GM, Hoiby EA, Holst J, Nøkleby H, Rosenqvist E, Sierra G, Campa C, Sotolongo F, Vega J, Garcia J, Herrera P, Poolman JT, Perkins BA. 1999. Immunogenicity of 2 serogroup B outer-membrane protein meningococcal vaccines: a randomized controlled trial in Chile. *JAMA* 281:1520–1527. <https://doi.org/10.1001/jama.281.16.1520>.
20. Tondella ML, Popovic T, Rosenstein NE, Lake DB, Carlone GM, Mayer LW, Perkins BA. 2000. Distribution of *Neisseria meningitidis* serogroup B serosubtypes and serotypes circulating in the United States. The Active Bacterial Core Surveillance Team. *J Clin Microbiol* 38:3323–3328.
21. Masignani V, Comanducci M, Giuliani MM, Bambini S, Adu-Bobie J, Arico B, Brunelli B, Pieri A, Santini L, Savino S, Serruto D, Litt D, Kroll S, Welsch JA, Granoff DM, Rappuoli R, Pizza M. 2003. Vaccination against *Neisseria meningitidis* using three variants of the lipoprotein GNA1870. *J Exp Med* 197:789–799. <https://doi.org/10.1084/jem.20021911>.
22. Fletcher LD, Bernfield L, Barniak V, Farley JE, Howell A, Knauf M, Ooi P, Smith RP, Weise P, Wetherell M, Xie X, Zagursky R, Zhang Y, Zlotnick GW. 2004. Vaccine potential of the *Neisseria meningitidis* 2086 lipoprotein. *Infect Immun* 72:2088–2100. <https://doi.org/10.1128/IAI.72.4.2088-2100.2004>.
23. Giuliani MM, Adu-Bobie J, Comanducci M, Arico B, Savino S, Santini L, Brunelli B, Bambini S, Biolchi A, Capecci B, Cartocci E, Ciocchi L, Di Marcello F, Ferlicca F, Galli B, Luzzi E, Masignani V, Serruto D, Veggi D, Contorni M, Morandi M, Bartalesi A, Cinotti V, Mannucci D, Titta F, Ovidi E, Welsch JA, Granoff D, Rappuoli R, Pizza M. 2006. A universal vaccine for serogroup B meningococcus. *Proc Natl Acad Sci U S A* 103: 10834–10839. <https://doi.org/10.1073/pnas.0603940103>.
24. Lucidarme J, Tan L, Exley RM, Findlow J, Borrow R, Tang CM. 2011. Characterization of *Neisseria meningitidis* isolates that do not express the virulence factor and vaccine antigen factor H binding protein. *Clin Vaccine Immunol* 18:1002–1014. <https://doi.org/10.1128/CVI.00055-11>.
25. Schneider MC, Prosser BE, Caesar JJ, Kugelberg E, Li S, Zhang Q, Quoraishi S, Lovett JE, Deane JE, Sim RB, Roversi P, Johnson S, Tang CM, Lea SM. 2009. *Neisseria meningitidis* recruits factor H using protein mimicry of host carbohydrates. *Nature* 458:890–893. <https://doi.org/10.1038/nature07769>.
26. Brehony C, Wilson DJ, Maiden MC. 2009. Variation of the factor H-binding protein of *Neisseria meningitidis*. *Microbiology* 155:4155–4169. <https://doi.org/10.1099/mic.0.027995-0>.
27. Kristensen T, Tack BF. 1986. Murine protein H is comprised of 20 repeating units, 61 amino acids in length. *Proc Natl Acad Sci U S A* 83: 3963–3967. <https://doi.org/10.1073/pnas.83.11.3963>.
28. Klickstein LB, Wong WW, Smith JA, Weis JH, Wilson JG, Fearon DT. 1987. Human C3b/C4b receptor (CR1). Demonstration of long homologous repeating domains that are composed of the short consensus repeats characteristics of C3/C4 binding proteins. *J Exp Med* 165:1095–1112.
29. Ripoché J, Day AJ, Harris TJ, Sim RB. 1988. The complete amino acid sequence of human complement factor H. *Biochem J* 249:593–602. <https://doi.org/10.1042/bj2490593>.
30. Weiler JM, Daha MR, Austen KF, Fearon DT. 1976. Control of the amplification convertase of complement by the plasma protein beta1H. *Proc Natl Acad Sci U S A* 73:3268–3272. <https://doi.org/10.1073/pnas.73.9.3268>.
31. Whaley K, Ruddy S. 1976. Modulation of the alternative complement pathways by beta 1 H globulin. *J Exp Med* 144:1147–1163. <https://doi.org/10.1084/jem.144.5.1147>.
32. Pangburn MK, Schreiber RD, Müller-Eberhard HJ. 1977. Human complement C3b inactivator: isolation, characterization, and demonstration of an absolute requirement for the serum protein beta1H for cleavage of C3b and C4b in solution. *J Exp Med* 146:257–270. <https://doi.org/10.1084/jem.146.1.257>.
33. Schmidt CQ, Herbert AP, Kavanagh D, Gandy C, Fenton CJ, Blaum BS, Lyon M, Uhrin D, Barlow PN. 2008. A new map of glycosaminoglycan and C3b binding sites on factor H. *J Immunol* 181:2610–2619. <https://doi.org/10.4049/jimmunol.181.4.2610>.
34. Schmidt CQ, Herbert AP, Hocking HG, Uhrin D, Barlow PN. 2008. Translational mini-review series on complement factor H: structural and functional correlations for factor H. *Clin Exp Immunol* 151:14–24. <https://doi.org/10.1111/j.1365-2249.2007.03553.x>.
35. Meri T, Amdahl H, Lehtinen MJ, Hyvärinen S, McDowell JV, Bhattacharjee A, Meri S, Marconi R, Goldman R, Jokiranta TS. 2013. Microbes bind complement inhibitor factor H via a common site. *PLoS Pathog* 9:e1003308. <https://doi.org/10.1371/journal.ppat.1003308>.
36. Caesar JJ, Lavender H, Ward PN, Exley RM, Eaton J, Chittock E, Malik TH, Goicoechea De Jorge E, Pickering MC, Tang CM, Lea SM. 2014. Competition between antagonistic complement factors for a single protein on *N. meningitidis* rules disease susceptibility. *eLife* 3. <https://doi.org/10.7554/eLife.04008>.
37. Muzzi A, Mora M, Pizzi M, Rappuoli R, Donati C. 2013. Conservation of meningococcal antigens in the genus *Neisseria*. *mBio* 4:e00163-13. <https://doi.org/10.1128/mBio.00163-13>.
38. Hadad R, Jacobsson S, Pizzi M, Rappuoli R, Fredlund H, Olcén P, Unemo M. 2012. Novel meningococcal 4CMenB vaccine antigens; prevalence and polymorphisms of the encoding genes in *Neisseria gonorrhoeae*. *APMIS* 120:750–760. <https://doi.org/10.1111/j.1600-0463.2012.02903.x>.
39. Jongerius I, Lavender H, Tan L, Ruivo N, Exley RM, Caesar JJ, Lea SM, Johnson S, Tang CM. 2013. Distinct binding and immunogenic properties of the gonococcal homologue of meningococcal factor H binding protein. *PLoS Pathog* 9:e1003528. <https://doi.org/10.1371/journal.ppat.1003528>.
40. Cann KJ, Rogers TR. 1989. Detection of antibodies to common antigens of pathogenic and commensal *Neisseria* species. *J Med Microbiol* 30: 23–31. <https://doi.org/10.1099/00222615-30-1-23>.
41. Snyder LA, Saunders NJ. 2006. The majority of genes in the pathogenic *Neisseria* species are present in non-pathogenic *Neisseria lactamica*, including those designated as ‘virulence genes’. *BMC Genomics* 7:128. <https://doi.org/10.1186/1471-2164-7-128>.
42. Evans CM, Pratt CB, Matheson M, Vaughan TE, Findlow J, Borrow R, Gorringer AR, Read RC. 2011. Nasopharyngeal colonization by *Neisseria lactamica* and induction of protective immunity against *Neisseria meningitidis*. *Clin Infect Dis* 52:70–77. <https://doi.org/10.1093/cid/ciq065>.
43. Wörmann ME, Horien CL, Johnson E, Liu G, Aho E, Tang CM, Exley RM. 2016. *Neisseria cinerea* isolates can adhere to human epithelial cells by type IV pilus-independent mechanisms. *Microbiology* 162:487–502. <https://doi.org/10.1099/mic.0.000248>.
44. Johnson S, Tan L, van der Veen S, Caesar J, Goicoechea De Jorge E, Harding RJ, Bai X, Exley RM, Ward PN, Ruivo N, Trivedi K, Cumber E, Jones R, Newham L, Staunton D, Ufret-Vincenty R, Borrow R, Pickering MC, Lea SM, Tang CM. 2012. Design and evaluation of meningococcal vaccines through structure-based modification of host and pathogen molecules. *PLoS Pathog* 8:e1002981. <https://doi.org/10.1371/journal.ppat.1002981>.
45. Loh E, Kugelberg E, Tracy A, Zhang Q, Gollan B, Ewles H, Chalmers R, Pellic V, Tang CM. 2013. Temperature triggers immune evasion by *Neisseria meningitidis*. *Nature* 502:237–240. <https://doi.org/10.1038/nature12616>.
46. Loh E, Lavender H, Tan F, Tracy A, Tang CM. 2016. Thermoregulation of meningococcal fHbp, an important virulence factor and vaccine antigen, is mediated by anti-ribosomal binding site sequences in the open reading frame. *PLoS Pathog* 12:e1005794. <https://doi.org/10.1371/journal.ppat.1005794>.
47. Sim E, Palmer MS, Puklavec M, Sim RB. 1983. Monoclonal antibodies

- against the complement control protein factor H (beta 1 H). *Biosci Rep* 3:1119–1131. <https://doi.org/10.1007/BF01120205>.
48. Des Prez RM, Bryan CS, Hawiger J, Colley DG. 1975. Function of the classical and alternate pathways of human complement in serum treated with ethylene glycol tetraacetic acid and MgCl₂-ethylene glycol tetraacetic acid. *Infect Immun* 11:1235–1243.
 49. Borrow R, Balmer P, Miller E. 2005. Meningococcal surrogates of protection—serum bactericidal antibody activity. *Vaccine* 23:2222–2227. <https://doi.org/10.1016/j.vaccine.2005.01.051>.
 50. Madico G, Welsch JA, Lewis LA, McNaughton A, Perlman DH, Costello CE, Ngampasutadol J, Vogel U, Granoff DM, Ram S. 2006. The meningococcal vaccine candidate GNA1870 binds the complement regulatory protein factor H and enhances serum resistance. *J Immunol* 177:501–510. <https://doi.org/10.4049/jimmunol.177.1.501>.
 51. Schneider MC, Exley RM, Chan H, Feavers I, Kang YH, Sim RB, Tang CM. 2006. Functional significance of factor H binding to *Neisseria meningitidis*. *J Immunol* 176:7566–7575. <https://doi.org/10.4049/jimmunol.176.12.7566>.
 52. Ngampasutadol J, Ram S, Gulati S, Agarwal S, Li C, Visintin A, Monks B, Madico G, Rice PA. 2008. Human factor H interacts selectively with *Neisseria gonorrhoeae* and results in species-specific complement evasion. *J Immunol* 180:3426–3435. <https://doi.org/10.4049/jimmunol.180.5.3426>.
 53. Ram S, McQuillen DP, Gulati S, Elkins C, Pangburn MK, Rice PA. 1998. Binding of complement factor H to loop 5 of porin protein 1A: a molecular mechanism of serum resistance of nonsialylated *Neisseria gonorrhoeae*. *J Exp Med* 188:671–680. <https://doi.org/10.1084/jem.188.4.671>.
 54. Persson CG, Erjefält I, Alkner U, Baumgarten C, Greiff L, Gustafsson B, Luts A, Pipkorn U, Sundler F, Svensson C. 1991. Plasma exudation as a first line respiratory mucosal defence. *Clin Exp Allergy* 21:17–24.
 55. Kaul TN, Welliver RC, Ogra PL. 1982. Appearance of complement components and immunoglobulins on nasopharyngeal epithelial cells following naturally acquired infection with respiratory syncytial virus. *J Med Virol* 9:149–158. <https://doi.org/10.1002/jmv.1890090210>.
 56. Prellner K, Stjernquist-Desatnik A. 1985. A simple method for collecting nasopharyngeal secretions, using cross-linked dextran. *Clin Otolaryngol Allied Sci* 10:85–87. <https://doi.org/10.1111/j.1365-2273.1985.tb01172.x>.
 57. Uhlén M, Fagerberg L, Hallström BM, Lindskog C, Oksvold P, Mardinoglu A, Sivertsson Å, Kampf C, Sjöstedt E, Asplund A, Olsson I, Edlund K, Lundberg E, Navani S, Szizyarto CA, Odeberg J, Djureinovic D, Takanen JO, Hober S, Alm T, Edqvist PH, Berling H, Tegel H, Mulder J, Rockberg J, Nilsson P, Schwenk JM, Hamsten M, von Feilitzen K, Forsberg M, Persson L, Johansson F, Zwahlen M, von Heijne G, Nielsen J, Pontén F. 2015. Proteomics tissue-based map of the human proteome. *Science* 347:1260419. <https://doi.org/10.1126/science.1260419>.
 58. Esparza-Gordillo J, Soria JM, Buil A, Almasy L, Blangero J, Fontcuberta J, Rodríguez de Córdoba S. 2004. Genetic and environmental factors influencing the human factor H plasma levels. *Immunogenetics* 56:77–82. <https://doi.org/10.1007/s00251-004-0660-7>.
 59. Sofat R, Mangione PP, Gallimore JR, Hakobyan S, Hughes TR, Shah T, Goodship T, D'Aiuto F, Langenberg C, Wareham N, Morgan BP, Pepys MB, Hingorani AD. 2013. Distribution and determinants of circulating complement factor H concentration determined by a high-throughput immunonephelometric assay. *J Immunol Methods* 390:63–73. <https://doi.org/10.1016/j.jim.2013.01.009>.
 60. Närkiö-Mäkelä M, Hellwage J, Tahkokallio O, Meri S. 2001. Complement-regulator factor H and related proteins in otitis media with effusion. *Clin Immunol* 100:118–126. <https://doi.org/10.1006/clim.2001.5043>.
 61. Li JF, Dai XP, Zhang W, Sun SH, Zeng Y, Zhao GY, Kou ZH, Guo Y, Yu H, Du LY, Jiang SB, Zhou YS. 2015. Upregulation of microRNA-146a by hepatitis B virus X protein contributes to hepatitis development by downregulating complement factor H. *mBio* 6:e02459-14. <https://doi.org/10.1128/mBio.02459-14>.
 62. Pandiripally V, Wei L, Skerka C, Zipfel PF, Cue D. 2003. Recruitment of complement factor H-like protein 1 promotes intracellular invasion by group A streptococci. *Infect Immun* 71:7119–7128. <https://doi.org/10.1128/IAI.71.12.7119-7128.2003>.
 63. Hammerschmidt S, Agarwal V, Kunert A, Haelbich S, Skerka C, Zipfel PF. 2007. The host immune regulator factor H interacts via two contact sites with the PspC protein of *Streptococcus pneumoniae* and mediates adhesion to host epithelial cells. *J Immunol* 178:5848–5858. <https://doi.org/10.4049/jimmunol.178.9.5848>.
 64. Hellwage J, Kühn S, Zipfel PF. 1997. The human complement regulatory factor-H-like protein 1, which represents a truncated form of factor H, displays cell-attachment activity. *Biochem J* 326:321–327. <https://doi.org/10.1042/bj3260321>.
 65. Lewis LA, Ngampasutadol J, Wallace R, Reid JE, Vogel U, Ram S. 2010. The meningococcal vaccine candidate neisserial surface protein A (NspA) binds to factor H and enhances meningococcal resistance to complement. *PLoS Pathog* 6:e1001027. <https://doi.org/10.1371/journal.ppat.1001027>.
 66. Lewis LA, Vu DM, Vasudhev S, Shaughnessy J, Granoff DM, Ram S. 2013. Factor H-dependent alternative pathway inhibition mediated by porin B contributes to virulence of *Neisseria meningitidis*. *mBio* 4:e00339-13. <https://doi.org/10.1128/mBio.00339-13>.
 67. Seib KL, Serruto D, Oriente F, Delany I, Adu-Bobie J, Veggi D, Aricò B, Rappuoli R, Pizza M. 2009. Factor H-binding protein is important for meningococcal survival in human whole blood and serum and in the presence of the antimicrobial peptide LL-37. *Infect Immun* 77:292–299. <https://doi.org/10.1128/IAI.01071-08>.
 68. Dürr UH, Sudheendra US, Ramamoorthy A. 2006. LL-37, the only human member of the cathelicidin family of antimicrobial peptides. *Biochim Biophys Acta* 1758:1408–1425. <https://doi.org/10.1016/j.bbame.2006.03.030>.
 69. Lysenko ES, Gould J, Bals R, Wilson JM, Weiser JN. 2000. Bacterial phosphorylcholine decreases susceptibility to the antimicrobial peptide LL-37/hCAP18 expressed in the upper respiratory tract. *Infect Immun* 68:1664–1671. <https://doi.org/10.1128/IAI.68.3.1664-1671.2000>.
 70. Veggi D, Gentile MA, Cantini F, Lo Surdo P, Nardi-Dei V, Seib KL, Pizza M, Rappuoli R, Banci L, Savino S, Scarselli M. 2012. The factor H binding protein of *Neisseria meningitidis* interacts with xenosiphophores in vitro. *Biochemistry* 51:9384–9393. <https://doi.org/10.1021/bi301161w>.
 71. Barnwal RP, Loh E, Godin KS, Yip J, Lavender H, Tang CM, Varani G. 2016. Structure and mechanism of a molecular rheostat, an RNA thermometer that modulates immune evasion by *Neisseria meningitidis*. *Nucleic Acids Res* 44:9426–9437.
 72. Bjornson AB, Mellencamp MA, Schiff GM. 1991. Complement is activated in the upper respiratory tract during influenza virus infection. *Am Rev Respir Dis* 143:1062–1066. https://doi.org/10.1164/ajrccm/143.5_Pt_1.1062.
 73. Gentile D, Doyle W, Whiteside T, Fireman P, Hayden FG, Skoner D. 1998. Increased interleukin-6 levels in nasal lavage samples following experimental influenza A virus infection. *Clin Diagn Lab Immunol* 5:604–608.
 74. Trotter CL, Andrews NJ, Kaczmarek EB, Miller E, Ramsay ME. 2004. Effectiveness of meningococcal serogroup C conjugate vaccine 4 years after introduction. *Lancet* 364:365–367. [https://doi.org/10.1016/S0140-6736\(04\)16725-1](https://doi.org/10.1016/S0140-6736(04)16725-1).
 75. Balmer P, Borrow R, Miller E. 2002. Impact of meningococcal C conjugate vaccine in the UK. *J Med Microbiol* 51:717–722. <https://doi.org/10.1099/0022-1317-51-9-717>.
 76. Read RC, Baxter D, Chadwick DR, Faust SN, Finn A, Gordon SB, Heath PT, Lewis DJ, Pollard AJ, Turner DP, Bazaz R, Ganguli A, Havelock T, Neal KR, Okike IO, Morales-Aza B, Patel K, Snape MD, Williams J, Gilchrist S, Gray SJ, Maiden MC, Toneatto D, Wang H, McCarthy M, Dull PM, Borrow R. 2014. Effect of a quadrivalent meningococcal ACWY glycoconjugate or a serogroup B meningococcal vaccine on meningococcal carriage: an observer-blind, phase 3 randomised clinical trial. *Lancet* 384:2123–2131. [https://doi.org/10.1016/S0140-6736\(14\)60842-4](https://doi.org/10.1016/S0140-6736(14)60842-4).
 77. Soeters HM, Whaley M, Alexander-Scott N, Kanadianian KV, MacNeil JR, Martin SW, McNamara LA, Sicard K, Vanner C, Vuong J, Wang X, Bandy U, Patel M, Rhode Island Meningococcal Carriage Evaluation Team. 2017. Meningococcal carriage evaluation in response to a serogroup B meningococcal disease outbreak and mass vaccination campaign at a college—Rhode Island, 2015–2016. *Clin Infect Dis* 64:1115–1122. <https://doi.org/10.1093/cid/cix091>.
 78. Robert-Guroff M. 2000. IgG surfaces as an important component in mucosal protection. *Nat Med* 6:129–130. <https://doi.org/10.1038/72206>.
 79. Reichhardt MP, Meri S. 2016. SALSA: a regulator of the early steps of complement activation on mucosal surfaces. *Front Immunol* 7:85. <https://doi.org/10.3389/fimmu.2016.00085>.
 80. Deasy AM, Guccione E, Dale AP, Andrews N, Evans CM, Bennett JS, Bratcher HB, Maiden MC, Gorringer AR, Read RC. 2015. Nasal inoculation of the commensal *Neisseria lactamica* inhibits carriage of *Neisseria meningitidis* by young adults: a controlled human infection study. *Clin Infect Dis* 60:1512–1520. <https://doi.org/10.1093/cid/civ098>.
 81. Zhang H, Morrison MA, Dewan A, Adams S, Andreoli M, Huynh N, Regan M, Brown A, Miller JW, Kim IK, Hoh J, Deangelis MM. 2008. The NEI/NCBI dbGAP database: genotypes and haplotypes that may specifically pre-

- dispose to risk of neovascular age-related macular degeneration. *BMC Med Genet* 9:51. <https://doi.org/10.1186/1471-2350-9-51>.
82. Prosser BE, Johnson S, Roversi P, Herbert AP, Blaum BS, Tyrrell J, Jowitt TA, Clark SJ, Tarelli E, Uhrin D, Barlow PN, Sim RB, Day AJ, Lea SM. 2007. Structural basis for complement factor H linked age-related macular degeneration. *J Exp Med* 204:2277–2283. <https://doi.org/10.1084/jem.20071069>.
83. Exley RM, Shaw J, Mowe E, Sun YH, West NP, Williamson M, Botto M, Smith H, Tang CM. 2005. Available carbon source influences the resistance of *Neisseria meningitidis* against complement. *J Exp Med* 201: 1637–1645. <https://doi.org/10.1084/jem.20041548>.
84. Mehr IJ, Seifert HS. 1998. Differential roles of homologous recombination pathways in *Neisseria gonorrhoeae* pilin antigenic variation, DNA trans-formation and DNA repair. *Mol Microbiol* 30:697–710. <https://doi.org/10.1046/j.1365-2958.1998.01089.x>.
85. Bennett JS, Jolley KA, Maiden MC. 2013. Genome sequence analyses show that *Neisseria oralis* is the same species as '*Neisseria mucosa* var. *heidelbergensis*'. *Int J Syst Evol Microbiol* 63:3920–3926. <https://doi.org/10.1099/ijs.0.052431-0>.
86. Maiden MC, Bygraves JA, Feil E, Morelli G, Russell JE, Urwin R, Zhang Q, Zhou J, Zurth K, Caugant DA, Feavers IM, Achtman M, Spratt BG. 1998. Multilocus sequence typing: a portable approach to the identification of clones within populations of pathogenic microorganisms. *Proc Natl Acad Sci U S A* 95:3140–3145. <https://doi.org/10.1073/pnas.95.6.3140>.