

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

Structural study of bioisosteric derivatives of 5-(1 H-indol-3-yl)-benzotriazole and their ability to form chalcogen bonds

Mirgaux, Manon; Scaillet, Tanguy; Kozlova, Arina; Tumanov, Nikolay; Frédérick, Raphael; Bodart, Laurie; Wouters, Johan

Published in: Acta Crystallographica. Section E: Crystallographic Communications

DOI: 10.1107/S2056989022002948

Publication date: 2022

Document Version Publisher's PDF, also known as Version of record

Link to publication

Citation for pulished version (HARVARD):

Mirgaux, M, Scaillet, T, Kozlova, A, Tumanov, N, Frédérick, R, Bodart, L & Wouters, J 2022, 'Structural study of bioisosteric derivatives of 5-(1 H-indol-3-yl)-benzotriazole and their ability to form chalcogen bonds', Acta Crystallographica. Section E: Crystallographic Communications, vol. 78, no. 4, pp. 418-424. https://doi.org/10.1107/S2056989022002948

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Received 9 March 2022 Accepted 16 March 2022

Edited by L. Van Meervelt, Katholieke Universiteit Leuven, Belgium

Keywords: crystal structure; bioisosterism; chalcogen; chalcogen interaction; tryptophan-2,3-dioxygenase inhibitors; X-ray crystallography.

Supporting information: this article has supporting information at journals.iucr.org/e

Structural study of bioisosteric derivatives of 5-(1*H*-indol-3-yl)-benzotriazole and their ability to form chalcogen bonds

Manon Mirgaux,^a* Tanguy Scaillet,^a Arina Kozlova,^b Nikolay Tumanov,^a Raphaël Frederick,^b Laurie Bodart^a and Johan Wouters^a*

^aNamur Institute of Structured Matter (NISM), Namur Research Institute for Life Science (NARILIS), Department of Chemistry, Laboratoire de Chimie Biologique Structurale (CBS) University of Namur (UNamur), 61 Rue de Bruxelles, 5000, Namur, Belgium, and ^bLouvain Drug Research Institute (LDRI), Université Catholique de Louvain (UCLouvain), Brussels B-1200, Belgium. *Correspondence e-mail: manon.mirgaux@unamur.be, johan.wouters@unamur.be

Recently, interest in the isosteric replacement of a nitrogen atom to selenium, sulfur or oxygen atoms has been highlighted in the design of potential inhibitors for cancer research. In this context, the structures of 5-(1H-indol-3-yl)-2,1,3benzotriazole derivatives [5-(1H-indol-3-yl)-2,1,3-benzothiadiazole (bS. C₁₄H₉N₃S) and 5-(1*H*-indol-3-yl)-2,1,3-benzoxadiazole (bO, C₁₄H₉N₃O)], as well as a synthesis intermediate of the selenated bioisostere [5-[1-(benzensulfonyl)-1*H*-indol-3-yl]-2,1,3-benzoselenadiazole (p-bSe, $C_{20}H_{13}N_3O_2SSe)$] were determined using single-crystal X-ray diffraction (SCXRD) analyses. Despite being analogues, different crystal packing, torsion angles and supramolecular features were observed, depending on the substitution of the central atoms of the benzotriazole. In particular, chalcogen interactions were described in the case of p-bSe and not in the bS and bO derivatives. An investigation by *ab initio* computational methods was therefore conducted to understand the effect of the substitution on the ability to form chalcogen bonds and the flexibility of the compounds.

1. Chemical context

Isosteric replacement is a common strategy in drug design to modulate the physicochemical properties of potential inhibitors. In 2021, Kozlova and co-workers (Kozlova *et al.*, 2021*a*) highlighted a series of bioisosteric derivatives acting as potential new inhibitors of the protein hTDO2, a therapeutic target in cancer research. These new molecules differ in the replacement of the central atom of benzotriazole by an oxygen, sulfur or selenium atom (Fig. 1). At this time, these inhibitors have not yet been crystallized or structurally char-





X = NH, O, S, Se

Figure 1 Structures of bioisosteres of 5-(1H-indol-3-yl)-benzotriazole with their torsion angle. X = NH, O, S, Se. acterized. In this context, the present work provides a structural characterization of the inhibitors described by Kozlova *et al.* (2021*b*) completed by *ab initio* calculations for their conformational characterization.

The contribution of an oxygen, a sulfur or a selenium atom instead of a nitrogen affects the ability of these inhibitors to participate in the formation of chalcogen bonds (Vogel et al., 2019). In particular, in these compounds, oxygen, sulfur and selenium atoms could act as chalcogen-bond donors. In recent years, the importance of chalcogen bonds in the stability and folding of proteins as well as their interaction with ligands has been highlighted by numerous investigations (Newberry & Raines, 2019; Kříž et al., 2018; Iwaoka et al., 2001; Iwaoka & Babe, 2015; Burling & Goldstein, 1992). In this article, the potential ability of the compounds to interact with aromatic groups, by chalcogen $-\pi$ interactions (Aakeroy *et al.*, 2019), was revealed by the crystallization of 5-[1-(benzensulfonyl)-1Hindol-3-yl] -2,1,3-benzoselenadiazole. Therefore, the effect of bioisosteric replacement on the ability to form chalcogen bonds has been studied by ab initio calculated electrostatic potential maps. This interesting series could be the starting point for the study of the effect of chalcogen interaction on protein stability and affinity.



2. Structural commentary

The compounds investigated in this study were kindly provided by the team of Raphaël Frédérick (UCLouvain, Belgium). Crystallization assays were performed, by slow evaporation at room temperature (293–298 K), in four different solvents [tetrahydrofuran (THF), chloroform, dichloromethane and *N*,*N*-dimethylformamide (DMF)]. Crystals of 5-(1*H*-indol-3-yl)-2,1,3-benzoxadiazole (bO) and of 5-(1*H*-indol-3-yl)-2,1,3-benzothiadiazole (bS) were obtained



Figure 2

Ellipsoid plots with atom labeling for (a) p-bSe (b) bS and (c) bO. Displacement ellipsoids are drawn at the 50% probability level.



Figure 3

Two perpendicular views (I. and II.) of the two mirror images observed in the crystal packings for (*a*) p-bSe (*b*) bS and (*c*) bO. Superposition of the mirror images with respect to the indole group. Torsion angles are ± 168.3 (2)° for p-bSe, ± 36.9 (2)° for bS and ± 146.7 (2)° for bO.

from chloroform. Despite numerous attempts, we were not able to crystallize the compound 5-(1H-indol-3-yl)-2,1,3-benzoselenadiazole (bSe). However, crystals of a synthesis intermediate -5-[1-(benzensulfonyl)-1H-indol-3-yl]-2,1,3-benzoselenadiazole (p-bSe) – were obtained in THF.

5-[1-(Benzensulfonyl)-1H-indol-3-yl]-2,1,3-benzoselenadiazole (p-bSe) crystallized in space group $P\overline{1}$ with one molecule of p-bSe in the asymmetric unit [Fig. 2(a)]. Interestingly, the molecule adopts an almost planar dihedral angle $[-168.3 (2)^{\circ}]$ between the indole and benzoselenadiazole ring (Fig. 3). 5-(1*H*-indol-3-yl)-2,1,3-benzothiadiazole (bS)and 5-(1H-indol-3-yl)-2,1,3-benzoxadiazole (bO) crystallized in space group Pbca [Fig. 2(b) and (c)]. The asymmetric units contain one molecule of bS or bO without disorder. In the three structures, two mirror images are observed in the crystal packing with a torsion angle of $\pm 168.3 (2)^{\circ}$ for p-bSe, $\pm 36.9 (2)^{\circ}$ for bS and $\pm 146.7 (2)^{\circ}$ for bO between the two aromatic parts of the molecules (Fig. 3). The isosteric replacement does not change significantly the planarity of the benzotriazole ring (r.m.s. deviation from planarity: 0.013 Å for p-bSe, 0.006 Å for bS and bO) or the indole ring (0.010 Å for p-bSe, 0.025 Å for bS and 0.011 Å for bO).

3. Supramolecular features

In the structure of p-bSe, a synthesis intermediate of the selenated bioisostere of 5-(1H-indol-3-yl)benzotriazole, the benzenesulfonyl contributes to the stabilization of the crystal

research communications

Table 1	
Hydrogen-bond geometry (Å, °) for p-bSe.	

$D - H \cdots A$	D-H	$H \cdot \cdot \cdot A$	$D \cdots A$	$D - H \cdots A$
$\begin{array}{c} C8-H8\cdots O2^{i}\\ C14-H14\cdots O1\end{array}$	0.93	2.46	3.380 (3)	168
	0.93	2.54	3.099 (4)	119

Symmetry code: (i) -x + 1, -y, -z + 1.

packing through weak hydrogen bonds [Table 1, Fig. 4(a)] and chalcogen $-\pi$ interactions. π -stacking interactions are observed between the selenadiazole and indole groups [centroid (Se/N1/C1/C6/N2)···centroid (N3/C7-C10) distance of 3.732 (2) Å, perpendicular distance of 3.587 (1) Å and horizontal displacement of 1.506 Å, Fig. 4(b)]. A second π stacking interaction is observed between the selenadiazole group (Se/N1/C1/C6/N2) and the indole group (C9-C14) [centroid ··· centroid distance of 3.915 (2) Å, perpendicular distance of 3.646 (1) Å and horizontal displacement of 1.331 Å, Fig. 4(b)]. A chalcogen $-\pi$ interaction between Se and the benzensulfonyl group (C15-C20) is also involved in crystal-packing stabilization [Se-.-centroid distance of 3.388 (1) Å and N2-Se \cdots centroid angle of 159.83 (8)°, Fig. 4(b)]. The presence of the protecting group (benzenesulfonyl) could explain the crystallization of the p-bSe compound with respect to the bSe compound. Indeed, in p-bSe, the orientation of the protecting group is ideal for allowing a chalcogen- π interaction whereas this type of interactions would be more difficult to set up in bSe.

In the structure of compound bS, π -interactions stabilize the crystal packing. π -stacking is observed between benzothiadiazole groups [centroid (C1–C6)···centroid (S1/N1/C1/C2/N2) distance of 3.689 (1) Å, perpendicular distance of 3.4989 (7) Å and horizontal displacement of 1.326 Å, Fig. 5(*a*)]. An N–H·· π interaction is also observed between indole groups [N3···centroid (C9–C14) distance of 3.345 (2) Å, H3N···centroid distance of 2.57 (2) Å and N3–H3N···centroid angle of 169 (2)°, Fig. 5(*b*)]. In this structure, no chalcogen interaction involving the sulfur atom is observed.

The crystal packing of bO is stabilized through π -interactions. π -stacking is observed between benzoxadiazole groups [centroid (O1/N1/C1–C6/N2)···centroid (O1/N1/C1– C6/N2) distance of 3.893 (1) Å, perpendicular distance of 3.5469 (8) Å and horizontal displacement of 1.570 Å, Fig. 6(*a*)]. An N–H·· π interaction is also observed between indole groups [N3···centroid (C9–C14) distance of 3.226 (2) Å, H3N···centroid distance of 2.57 (2) Å and N3– H3N···centroid angle of 138 (2)°, Fig. 6(*b*)]. No chalcogen interaction is observed.





Figure 4

Supramolecular organization of p-bSe: (*a*) hydrogen-bond interactions; (*b*) π -stacking interaction between the benzoselenadiazole group and the indole groups (centroids in red: Se/N1/C1/C6/N2 and N3/C7–C10 and centroids in green: Se/N1/C1/C6/N2 and C9–C14) as well as chalcogen- π interactions (in purple).

Figure 5

Supramolecular organization of bS. (a) π -stacking interaction between the benzothiadiazole groups and (b) N-H··· π interaction between two indole groups.

research communications





Figure 6

Supramolecular organization of bO. (a) π -stacking interaction between the benzoxadiazole groups and (b) N-H··· π interaction between two indole groups

4. Quantum *ab initio* studies of the bioisosteric substitution effect

As mentioned previously, the different derivatives vary mainly in their ability to interact through chalcogen bonds. In order to



Figure 7

Computed electrostatic potential (EPS) surfaces and associated molecular structures (a) bSe (b) bS (c) bO. The EPS color scale ranges from +8.160 volt to -5.440 volt.



Figure 8

Computed electrostatic potential (EPS) surfaces and associated molecular structures (a) bSe (b) p-bSe. The EPS color scale ranges from +8.160 volts to -5.440 volts.

characterize these differences in depth, quantum mechanics studies have been conducted. First, the presence of a σ -hole in the electron density was studied by means of electrostatic maps. Analysis indicates that the oxygen in benzoxadiazole [Fig. 7(c)] has a weakly positive environment. The σ -hole formation is enhanced by substitution of the central atom with sulfur [Fig. 7(b)] and selenium [Fig. 7(a)], with selenium having the most positive environment. The bioisosteric series thus has different characteristics in terms of the ability to form chalcogen bonds, with the selenium compound being the best chaclogen-bond donor in this bioisosteric series of molecules. These results may explain why chalcogen bonds are observed only in the supramolecular organization of the p-bSe molecule. The donor character of the selenium atom is not affected by the protected group [Fig. 8(a) and (b)]. The difficulty in crystallizing bSe (while p-bSe crystallized readily in THF) could be explained by the absence of the protecting group (benzenesulfonyl) in bSe. Indeed, the benzenesulfonyl group in p-bSe is electron-rich and acts as a well-oriented chalcogen-bond acceptor in p-bSe.

Secondly, in order to determine the effect of the substitution on the flexibility of the derivatives, conformational scans were performed around the torsion angle formed between the indole ring and the benzodiazole part (*T*1). As presented in Fig. 9, bO and bS are characterized by a very similar ΔE energy profile associated with the rotation around *T*1. For all three molecules (bO, bS and bSe), four minima are observed for each molecule, with symmetry on each side of the planar molecule. The energy transitions are low (maximum 15 kJ mol⁻¹) and the molecules are flexible. Moreover, the *T*1 torsion angles observed in the crystal structures of bS





Conformational scans and associated torsion angles calculated with Gaussian16a (ω B97XD, 6–31+G*).

CCDC refcode	Se label	Atoms of the π system	Secentroid distance	N−Se…centroid angle
QIBQUQ ^a	Se1	C18-C23	3.3676 (7)	147.08 (4)
OIBOUO ^a	Se1	C12-C17	3.0597 (8)	172.30 (5)
VOPMEV ^b	Se1	C5/C7/C10/C5B/C7B/C10B	3.8142 (3)	163.56 (4)
VOPNAS ^c	Se1	C1-C5/C10	3.802 (2)	162.5 (1)
VOPNAS ^c	Se2	C1-C5/C10	3.654 (2)	166.5 (1)
$YIWLOG^d$	Se5	C10-C15	4.032 (3)	144.6 (2)
$YIWLOG^d$	Se6	C34–C39	4.232 (4)	164.3 (4)

Table 2 Chalcogen bonds (Å, °) observed in benzoselenadiazole fragments in the CSD.

Notes: (a) Lee et al. (2018); (b) Lee et al., 2019); (c) Lee et al., 2019); (d) Tan et al. (2008).

 $[\pm 36.9 (2)^{\circ}]$ and bO $[\pm 146.7 (2)^{\circ}]$ are consistent with the energy minima determined by *ab initio* calculations with a relative deviation lower than 10%. Although the bioisosteric character of the flexibility is retained, two differences are observed between the bSe molecule and the bS and bO molecules. The first one is the energy at a torsion angle of 180° , which is lower for the molecule of bSe while it is slightly higher at 0° . The second one is a small shift observed between the angle associated with the energy minima of the bSe molecule and the bS and bO molecules.

The same calculations were performed for the protected molecule (p-bSe). The energy profile associated with the rotation around T1 in p-bSe is similar to those determined for bS and bO. There is a small shift of the values of the angle corresponding to the energy minimum with respect to bSe. This shift may be due to the protecting group that causes steric hindrance in p-bSe. The energies corresponding to minima A and C are higher than the energies for minima B and D, involving two local minima and a preference in the conformers. The maxima of the energies between A/B and C/D are also lower than in the case of the other compounds, supporting this hypothesis. In the case of p-bSe, the torsion angle observed in the crystallographic structure $[\pm 168.3 (2)^{\circ}]$ corresponds to an energy maximum on the energy profile associated with the rotation around T1 determined by ab initio calculations. The quasi-planarity of the molecule observed in the crystallographic structure could be encouraged by the formation of chalcogen bonds stabilizing this conformation of p-bSe.

5. Database survey

The Cambridge Structural Database (CSD version 5.42; Groom et al., 2016) was searched with ConQuest (version



Figure 10

Fragments searched for in the CSD during the database survey analysis: (a) bSe fragment (b) bS fragment (c) bO fragment.

2021.2.0; Bruno *et al.*, 2002) for benzoselenadiazole, benzothiadiazole and benzoxadiazole fragments (Fig. 10). The search for the benzoselenadiazole fragment resulted in 57 hits. Chalcogen bonds are observed in all of these hits. In particular, chalcogen– π interactions are observed in four structures [CSD refcodes: QIBQUQ (Lee *et al.*, 2018), VOPMEV (Lee *et al.*, 2019), VOPNAS (Lee *et al.*, 2019), and YIWLOG (Tan *et al.*, 2008)], listed in Table 2. The search for a benzothiadiazole fragment resulted in 34 hits. Chalcogen bonds but no chalcogen– π interactions are observed in these hits. The search for a benzoxadiazole fragment resulted in 24 hits but no chalcogen interactions are observed with the oxygen atom of benzoxadiazole as a chalcogen donor.

6. Synthesis and crystallization

The synthesis of the various compounds was reported by Kozlova *et al.* (2021*a*). Crystallization of the 5-(1*H*-indol-3-yl)benzotriazole derivatives were carried out by the solvent evaporation method. The compounds were dissolved in THF, chloroform, dichloromethane or DMF until complete dissolution. Slow evaporation of the solvent at room temperature (293–297 K) yielded colorless crystals that were then picked for XRD analysis. Crystals of 5-(1*H*-indol-3-yl)-2,1,3-benzothiadiazole and 5-(1*H*-indol-3-yl)-2,1,3-benzoxadiazole were obtained from chloroform while the protected benzoselenadiazole (5-[1-(benzensulfonyl)-1*H*-indol-3-yl]-2,1,3-benzoselenadiazole) was crystallized in THF.

7. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 3. In all of the structures, hydrogen atoms were placed in calculated positions and refined using a riding model [C–H bond length of 0.93 Å, with $U_{\rm iso}({\rm H}) =$ $1.2U_{\rm eq}({\rm C})$]. In the structure of 5-(1*H*-indol-3-yl)-2,1,3-benzothiadiazole, bS, the hydrogen on the nitrogen atom in the indole group was refined without constraint and the refined N–H distance is 0.78 (2) Å.

8. Quantum ab initio methodology

All the molecules investigated in the study (bS, bO, bSe, p-bSe) were optimized starting from the crystal coordinates using the density functional method (DFT) with the exchange-

Table 3Experimental details.

	p-bSe	bS	bO
Crystal data			
Chemical formula	$C_{20}H_{12}N_{2}O_{2}SSe$	$C_{14}H_0N_2S$	$C_{14}H_0N_2O$
<i>M</i> _	438.35	251.30	235.24
Crystal system, space group	Triclinic. P1	Orthorhombic, Phca	Orthorhombic, Phca
Temperature (K)	295	295	295
a, b, c (Å)	7.7760 (3), 9.9573 (4), 11.4124 (6)	7.5884 (1), 7.1060 (1), 43.2464 (7)	12.0256 (7), 7.7396 (5), 23.8551 (16)
α, β, γ (°)	90.970 (4), 92.771 (4), 94.283 (3)	90, 90, 90	90, 90, 90
$V(Å^3)$	879.95 (7)	2331.98 (6)	2220.3 (2)
Z	2	8	8
Radiation type	Cu Ka	Cu Ka	Cu Ka
$\mu (\text{mm}^{-1})$	4.18	2.32	0.75
Crystal size (mm)	$0.19 \times 0.10 \times 0.01$	$0.29 \times 0.18 \times 0.04$	$0.12 \times 0.09 \times 0.03$
Data collection			
Diffractometer	Xcalibur, Ruby, Gemini ultra	Xcalibur, Ruby, Gemini ultra R	Xcalibur, Ruby, Gemini ultra
Absorption correction	Analytical (<i>CrysAlis PRO</i> ; Rigaku OD, 2020)	Analytical (<i>CrysAlis PRO</i> ; Rigaku OD, 2020)	Analytical (<i>CrysAlis PRO</i> ; Rigaku OD, 2020)
T_{\min}, T_{\max}	0.663, 0.958	0.663, 0.920	0.941, 0.981
No. of measured, independent and observed $[I > 2\sigma(I)]$ reflections	9941, 3113, 2571	10906, 2072, 1825	6883, 1972, 1275
R _{int}	0.030	0.025	0.047
$(\sin \theta / \lambda)_{\rm max} ({\rm \AA}^{-1})$	0.598	0.598	0.597
Refinement			
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.037, 0.101, 1.04	0.035, 0.098, 1.07	0.050, 0.134, 1.06
No. of reflections	3113	2072	1972
No. of parameters	244	167	166
H-atom treatment	H-atom parameters constrained	H atoms treated by a mixture of independent and constrained refinement	H atoms treated by a mixture of independent and constrained refinement
$\Delta \rho_{\rm max}, \Delta \rho_{\rm min} \ ({\rm e} \ {\rm \AA}^{-3})$	0.46, -0.61	0.20, -0.30	0.14, -0.16

Computer programs: CrysAlis PRO (Rigaku OD, 2020), SHELXT2014/5 (Sheldrick, 2015a), SHELXL2016/6 (Sheldrick, 2015b), ShelXle (Hübschle et al., 2011), OLEX2 (Dolomanov et al., 2009), and Mercury (Macrae et al., 2020).

correlation functional ω B97XD and the 6-31+G* basis set. Because we were not able to crystallize the bSe compound, this molecule was created by substitution of the sulfur atom for a selenium atom from the coordinates of the bS molecule. The optimizations were performed with Gaussian16a (Frisch et al., 2016) the in gas phase. The electrostatic potential was calculated from the SCF-type density and was sliced by making 80 cubic points evenly distributed on a rectangular grid automatically generated by Gaussian16a. The resulting maps were visualized using DrawMol (Liegeois, 2021). For the conformational scans, the optimized structures were analyzed using relaxed scans around the torsion angle (T1) formed between the indole ring and the benzodiazole part from 0 to 360° by steps of 20° . The resulting conformations close to an energy minimum were extracted and refined by a new optimization at the same level of approximation. The preparation of the input files, as well as the visualization of the results was performed with the DrawMol and DrawSpectrum suite of programs (Liegeois, 2021). The graphs were drawn with the program Prism from GraphPad (one-way ANOVA followed by Dunnetts multiple comparisons test, Prism version 8.0.0 for Windows; GraphPad, 2021).

Funding information

This research used resources of the 'Plateforme Technologique de Calcul Intensif (PTCI)' (http://www.ptci.unamur.be), located at the University of Namur, Belgium, which is supported by the FNRS–FRFC, the Walloon Region, and the University of Namur (Conventions Nos. 2.5020.11, GEQU·G006.15, 1610468, and RW/GEQ2016). The PTCI is a member of the 'Consortium des Equipements de Calcul Intensif (CÉCI)' (http://www.ceci-hpc.be). This work is supported by the Belgian Fonds National de la Recherche Scientifique (FRS–FNRS; grant Nos. 3.05557.43, 28252254 and 32704190), the French Community of Belgium (ARC 21/26– 115), the Fonds spéciaux de recherche (FSR) at UCLouvain, and a J. Maisin Foundation grant. AK and MM acknowledge the Fonds de la Recherche Scientifique (FRS–FNRS, Belgium) for their Research Fellow grants.

References

- Aakeroy, C. B., Bryce, D. L., Desiraju, G. R., Frontera, A., Legon, A. C., Nicotra, F., Rissanen, K., Scheiner, S., Terraneo, G., Metrangolo, P. & Resnati, G. (2019). *Pure Appl. Chem.* **91**, 1889– 1892.
- Bruno, I. J., Cole, J. C., Edgington, P. R., Kessler, M., Macrae, C. F., McCabe, P., Pearson, J. & Taylor, R. (2002). *Acta Cryst.* B58, 389– 397.
- Burling, F. T. & Goldstein, B. M. (1992). J. Am. Chem. Soc. 114, 2313–2320.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). J. Appl. Cryst. 42, 339–341.
- Frisch, M. J., Trucks, G. W., Schlegel, H. B., Scuseria, G. E., Robb, M. A., Cheeseman, J. R., Scalmani, G., Barone, V., Petersson, G. A.,

Nakatsuji, H., Li, X., Caricato, M., Marenich, A. V., Bloino, J., Janesko, B. G., Gomperts, R., Mennucci, B., Hratchian, H. P., Ortiz, J. V., Izmaylov, A. F., Sonnenberg, J. L., Williams-Young, D., Ding, F., Lipparini, F., Egidi, F., Goings, J., Peng, B., Petrone, A., Henderson, T., Ranasinghe, D., Zakrzewski, V. G., Gao, J., Rega, N., Zheng, G., Liang, W., Hada, M., Ehara, M., Toyota, K., Fukuda, R., Hasegawa, J., Ishida, M., Nakajima, T., Honda, Y., Kitao, O., Nakai, H., Vreven, T., Throssell, K., Montgomery, J. A. Jr, Peralta, J. E., Ogliaro, F., Bearpark, M. J., Heyd, J. J., Brothers, E. N., Kudin, K. N., Staroverov, V. N., Keith, T. A., Kobayashi, R., Normand, J., Raghavachari, K., Rendell, A. P., Burant, J. C., Iyengar, S. S., Tomasi, J., Cossi, M., Millam, J. M., Klene, M., Adamo, C., Cammi, R., Ochterski, J. W., Martin, R. L., Morokuma, K., Farkas, O., Foresman, J. B. & Fox, D. J. (2016). *Gaussian16a*. Gaussian, Inc., Wallingford, CT, USA.

- GraphPad (2021). *GraphPad*. GraphPad Software, San Diego, California USA, www. graphpad. com
- Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). Acta Cryst. B72, 171–179.
- Hübschle, C. B., Sheldrick, G. M. & Dittrich, B. (2011). J. Appl. Cryst. 44, 1281–1284.
- Iwaoka, M. & Babe, N. (2015). Phosphorus Sulfur Silicon, 190, 1257– 1264.
- Iwaoka, M., Takemoto, S., Okada, M. & Tomoda, S. (2001). Chem. Lett. 30, 132–133.

- Kozlova, A., Thabault, L., Dauguet, N., Deskeuvre, M., Stroobant, V., Pilotte, L. & Frédérick, R. (2021a). Eur. J. Med. Chem. 227, 113892. https://doi.org/10.1016/j.ejmech.2021.113892
- Kozlova, A., Thabault, L., Liberelle, M., Klaessens, S., Prévost, J. R., Mathieu, C., Pilotte, L., Stroobant, V., Van den Eynde, B. & Frédérick, R. (2021b). J. Med. Chem. 64, 10967–10980.
- Kříž, K., Fanfrlík, J. & Lepšík, M. (2018). ChemPhysChem, 19, 2540– 2548.
- Lee, J., Lee, L. M., Arnott, Z., Jenkins, H., Britten, J. F. & Vargas-Baca, I. (2018). New J. Chem. 42, 10555–10562.
- Lee, L. M., Corless, V., Luu, H., He, A., Jenkins, H., Britten, J. F., Adam Pani, F. & Vargas-Baca, I. (2019). *Dalton Trans.* 48, 12541– 12548.
- Liegeois, V. (2021). DrawMol. University of Namur, Belgium. www. unamur. be/drawmol
- Macrae, C. F., Sovago, I., Cottrell, S. J., Galek, P. T. A., McCabe, P., Pidcock, E., Platings, M., Shields, G. P., Stevens, J. S., Towler, M. & Wood, P. A. (2020). J. Appl. Cryst. 53, 226–235.
- Newberry, R. W. & Raines, R. T. (2019). Chem. Biol. 14, 1677-1686.
- Rigaku OD (2020). CrysAlis PRO. Rigaku Oxford Diffraction Ltd, Yarnton, England.
- Sheldrick, G. M. (2015a). Acta Cryst. A71, 3-8.
- Sheldrick, G. M. (2015b). Acta Cryst. C71, 3-8.
- Tan, C. K., Wang, J., Leng, J. D., Zheng, L. L. & Tong, M. L. (2008). *Eur. J. Inorg. Chem.* pp. 771–778.
- Vogel, L., Wonner, P. & Huber, S. M. (2019). Angew. Chem. Int. Ed. 58, 1880–1891.

Acta Cryst. (2022). E78, 418-424 [https://doi.org/10.1107/S2056989022002948]

Structural study of bioisosteric derivatives of 5-(1*H*-indol-3-yl)-benzotriazole and their ability to form chalcogen bonds

Manon Mirgaux, Tanguy Scaillet, Arina Kozlova, Nikolay Tumanov, Raphaël Frederick, Laurie Bodart and Johan Wouters

Computing details

For all structures, data collection: *CrysAlis PRO* (Rigaku OD, 2020); cell refinement: *CrysAlis PRO* (Rigaku OD, 2020); data reduction: *CrysAlis PRO* (Rigaku OD, 2020); program(s) used to solve structure: *SHELXT2014/5* (Sheldrick, 2015a); program(s) used to refine structure: *SHELXL2016/6* (Sheldrick, 2015b), *ShelXle* (Hübschle *et al.*, 2011), *OLEX2* (Dolomanov *et al.*, 2009); molecular graphics: *Mercury* (Macrae *et al.*, 2020).

5-[1-(Benzenesulfonyl)-1H-indol-3-yl]-2,1,3-benzoselenadiazole (p-bSe)

Crystal data

 $C_{20}H_{13}N_{3}O_{2}SSe$ $M_{r} = 438.35$ Triclinic, *P*1 a = 7.7760 (3) Å b = 9.9573 (4) Å c = 11.4124 (6) Å $a = 90.970 (4)^{\circ}$ $\beta = 92.771 (4)^{\circ}$ $\gamma = 94.283 (3)^{\circ}$ $V = 879.95 (7) Å^{3}$

Data collection

Xcalibur, Ruby, Gemini ultra diffractometer Radiation source: fine-focus sealed X-ray tube, Enhance Ultra (Cu) X-ray Source Detector resolution: 5.1856 pixels mm⁻¹ ω scans Absorption correction: analytical (CrysAlisPro; Rigaku OD, 2020) $T_{min} = 0.663, T_{max} = 0.958$

Refinement

Refinement on F^2 Least-squares matrix: full $R[F^2 > 2\sigma(F^2)] = 0.037$ $wR(F^2) = 0.101$ S = 1.043113 reflections Z = 2 F(000) = 440 $D_x = 1.654 \text{ Mg m}^{-3}$ Cu K α radiation, $\lambda = 1.54184 \text{ Å}$ Cell parameters from 4183 reflections $\theta = 3.9-66.7^{\circ}$ $\mu = 4.18 \text{ mm}^{-1}$ T = 295 K Plate, colourless $0.19 \times 0.10 \times 0.01 \text{ mm}$

9941 measured reflections 3113 independent reflections 2571 reflections with $I > 2\sigma(I)$ $R_{int} = 0.030$ $\theta_{max} = 67.2^{\circ}, \theta_{min} = 3.9^{\circ}$ $h = -9 \rightarrow 9$ $k = -11 \rightarrow 9$ $l = -13 \rightarrow 13$

244 parameters 0 restraints Primary atom site location: dual Secondary atom site location: dual Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained	$(\Delta/\sigma)_{\rm max} = 0.001$
$w = 1/[\sigma^2(F_o^2) + (0.0588P)^2 + 0.2071P]$	$\Delta \rho_{\rm max} = 0.46 \text{ e } \text{\AA}^{-3}$
where $P = (F_0^2 + 2F_c^2)/3$	$\Delta \rho_{\rm min} = -0.61 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Structures were solved by the dual-space method of ShelXT (Sheldrick, 2015a) within Olex² (Dolomanov *et al.*, 2009). Structures were refined by the least- squares method implemented in SHELXL (Sheldrick, 2015b) within ShelXle (Hübschle *et al.*, 2011). Structures and crystal packings were visualized using Mercury (Macrae *et al.*, 2020).

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ((Å	2
---	----	---

	x	У	Ζ	$U_{ m iso}$ */ $U_{ m eq}$	
Se1	0.90079 (4)	0.81216 (3)	0.75959 (3)	0.07181 (16)	
S 1	0.52981 (9)	0.00138 (6)	0.25117 (6)	0.05354 (19)	
01	0.4385 (3)	-0.0082(2)	0.13980 (18)	0.0639 (5)	
O2	0.4503 (3)	-0.0486 (2)	0.35371 (18)	0.0650 (5)	
N1	0.9080 (3)	0.7558 (3)	0.6120 (2)	0.0674 (7)	
N2	0.7790 (4)	0.6657 (3)	0.8062 (2)	0.0685 (7)	
N3	0.5808 (3)	0.1632 (2)	0.2811 (2)	0.0539 (5)	
C1	0.8223 (3)	0.6341 (3)	0.6050(2)	0.0529 (6)	
C2	0.7984 (4)	0.5541 (3)	0.5005 (2)	0.0558 (6)	
H2	0.846444	0.585254	0.432073	0.067*	
C3	0.7060 (3)	0.4321 (2)	0.4993 (2)	0.0481 (6)	
C4	0.6317 (4)	0.3868 (3)	0.6070 (2)	0.0570 (7)	
H4	0.566610	0.304496	0.606199	0.068*	
C5	0.6530 (4)	0.4592 (3)	0.7089 (3)	0.0620(7)	
Н5	0.604349	0.426314	0.776555	0.074*	
C6	0.7506 (4)	0.5862 (3)	0.7115 (2)	0.0553 (6)	
C7	0.6741 (3)	0.3458 (2)	0.3939 (2)	0.0477 (6)	
C8	0.6014 (3)	0.2176 (3)	0.3945 (2)	0.0519 (6)	
H8	0.569709	0.172160	0.461603	0.062*	
C9	0.6494 (3)	0.2603 (3)	0.2037 (2)	0.0517 (6)	
C10	0.7064 (3)	0.3750 (3)	0.2714 (2)	0.0503 (6)	
C11	0.7763 (5)	0.4873 (3)	0.2134 (3)	0.0675 (8)	
H11	0.814269	0.565849	0.255173	0.081*	
C12	0.7881 (5)	0.4801 (3)	0.0936 (3)	0.0802 (10)	
H12	0.834370	0.554638	0.054712	0.096*	
C13	0.7326 (5)	0.3643 (3)	0.0297 (3)	0.0730 (9)	
H13	0.744430	0.362173	-0.050976	0.088*	
C14	0.6602 (4)	0.2520 (3)	0.0831 (2)	0.0616 (7)	
H14	0.620702	0.174545	0.040267	0.074*	
C15	0.7292 (4)	-0.0687(2)	0.2387 (2)	0.0545 (6)	
C16	0.8268 (4)	-0.0925 (3)	0.3403 (3)	0.0637 (7)	
H16	0.787011	-0.072995	0.413787	0.076*	
C17	0.9851 (4)	-0.1459 (3)	0.3296 (3)	0.0732 (9)	

H17	1.052470	-0.162969	0.396443	0.088*	
C18	1.0426 (4)	-0.1736 (3)	0.2205 (3)	0.0733 (9)	
H18	1.149596	-0.208273	0.214087	0.088*	
C19	0.9442 (5)	-0.1509 (3)	0.1210 (3)	0.0723 (8)	
H19	0.984606	-0.170801	0.047749	0.087*	
C20	0.7853 (4)	-0.0986 (3)	0.1287 (3)	0.0636 (7)	
H20	0.717547	-0.083800	0.061463	0.076*	

Atomic displacement parameters $(Å^2)$

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Se1	0.0753 (2)	0.0682 (2)	0.0690 (3)	-0.00422 (16)	-0.00371 (17)	-0.02183 (16)
S 1	0.0587 (4)	0.0467 (4)	0.0525 (4)	-0.0097 (3)	-0.0021 (3)	-0.0040 (3)
01	0.0655 (12)	0.0654 (12)	0.0575 (12)	-0.0055 (9)	-0.0107 (9)	-0.0115 (9)
O2	0.0722 (12)	0.0598 (11)	0.0604 (12)	-0.0146 (9)	0.0078 (10)	0.0011 (9)
N1	0.0683 (15)	0.0635 (15)	0.0673 (16)	-0.0116 (11)	0.0024 (12)	-0.0126 (12)
N2	0.0805 (17)	0.0695 (16)	0.0551 (15)	0.0090 (13)	-0.0007 (13)	-0.0112 (12)
N3	0.0681 (14)	0.0440 (11)	0.0473 (12)	-0.0086 (9)	0.0006 (10)	-0.0024 (9)
C1	0.0520 (14)	0.0522 (14)	0.0534 (16)	0.0035 (11)	-0.0050 (12)	-0.0072 (11)
C2	0.0604 (16)	0.0571 (15)	0.0486 (15)	-0.0046 (12)	0.0043 (12)	-0.0038 (11)
C3	0.0519 (14)	0.0454 (13)	0.0471 (14)	0.0061 (10)	-0.0019 (11)	-0.0009 (10)
C4	0.0734 (18)	0.0461 (14)	0.0511 (16)	0.0018 (12)	0.0011 (13)	0.0019 (11)
C5	0.081 (2)	0.0578 (16)	0.0477 (16)	0.0068 (14)	0.0042 (14)	0.0048 (12)
C6	0.0607 (16)	0.0553 (15)	0.0501 (15)	0.0136 (12)	-0.0058 (13)	-0.0059 (12)
C7	0.0507 (14)	0.0481 (13)	0.0440 (14)	0.0034 (10)	0.0012 (11)	-0.0017 (10)
C8	0.0604 (15)	0.0502 (14)	0.0443 (14)	-0.0010 (11)	0.0010 (12)	0.0005 (11)
C9	0.0562 (15)	0.0506 (14)	0.0477 (15)	0.0020 (11)	-0.0008 (12)	0.0019 (11)
C10	0.0567 (15)	0.0463 (14)	0.0483 (15)	0.0048 (11)	0.0041 (12)	0.0004 (11)
C11	0.096 (2)	0.0478 (15)	0.0566 (18)	-0.0091 (14)	0.0058 (16)	-0.0007 (12)
C12	0.120 (3)	0.0613 (18)	0.0573 (19)	-0.0120 (18)	0.0133 (18)	0.0073 (14)
C13	0.103 (2)	0.0705 (19)	0.0446 (16)	-0.0002 (16)	0.0070 (16)	0.0032 (13)
C14	0.0788 (19)	0.0597 (16)	0.0450 (15)	0.0000 (14)	-0.0025 (14)	-0.0026 (12)
C15	0.0639 (16)	0.0401 (13)	0.0568 (16)	-0.0098 (11)	-0.0022 (13)	-0.0009 (11)
C16	0.0712 (19)	0.0583 (16)	0.0591 (17)	-0.0064 (13)	-0.0054 (14)	0.0028 (13)
C17	0.072 (2)	0.0647 (19)	0.080 (2)	-0.0023 (15)	-0.0186 (17)	0.0078 (16)
C18	0.0690 (19)	0.0574 (18)	0.093 (3)	0.0020 (14)	0.0006 (18)	0.0007 (16)
C19	0.082 (2)	0.0645 (19)	0.071 (2)	0.0070 (16)	0.0059 (17)	-0.0070 (15)
C20	0.0748 (19)	0.0565 (16)	0.0581 (18)	0.0004 (13)	-0.0013 (15)	-0.0041 (13)

Geometric parameters (Å, °)

Se1—N1	1.772 (3)	С8—Н8	0.9300	
Se1—N2	1.782 (3)	C9—C14	1.384 (4)	
S101	1.424 (2)	C9—C10	1.399 (4)	
S1—O2	1.430 (2)	C10—C11	1.398 (4)	
S1—N3	1.657 (2)	C11—C12	1.376 (5)	
S1—C15	1.758 (3)	C11—H11	0.9300	
N1-C1	1.337 (4)	C12—C13	1.383 (5)	

	1 220 (4)	C12 1112	0.0200
N2—C6	1.329 (4)	C12—H12	0.9300
N3—C8	1.391 (3)	C13—C14	1.379 (4)
N3—C9	1.414 (4)	C13—H13	0.9300
C1—C2	1.420 (4)	C14—H14	0.9300
C1—C6	1.436 (4)	C15—C20	1.383 (4)
C2—C3	1.365 (4)	C15—C16	1.388 (4)
С2—Н2	0.9300	C16—C17	1.386 (5)
C3—C4	1,448 (4)	С16—Н16	0.9300
C3-C7	1 466 (4)	C17-C18	1 374 (5)
CA = C5	1.100(1) 1.355(4)	C17 H17	0.9300
$C_4 = C_3$	0.0200	C_{1}^{18} C_{10}^{10}	1 270 (5)
C4 - H4	0.9300	C_{10} U_{10}	1.370(3)
C5—C6	1.424 (4)		0.9300
C5—H5	0.9300	C19—C20	1.383 (5)
С7—С8	1.357 (4)	С19—Н19	0.9300
C7—C10	1.463 (4)	С20—Н20	0.9300
N1 Sol N2	05.07(11)	C14 C0 C10	$122 \in (2)$
N1 - Se1 - N2	95.07(11)	C14 = C9 = C10	123.0(3)
01 - 51 - 02	120.66 (12)	C14 - C9 - N3	129.3 (3)
01—S1—N3	107.53 (12)	C10—C9—N3	107.1 (2)
O2—S1—N3	104.65 (12)	C11—C10—C9	117.9 (3)
01—S1—C15	108.75 (13)	C11—C10—C7	134.3 (3)
O2—S1—C15	109.34 (13)	C9—C10—C7	107.8 (2)
N3—S1—C15	104.72 (12)	C12—C11—C10	119.0 (3)
C1—N1—Se1	106.3 (2)	C12—C11—H11	120.5
C6—N2—Se1	105.8 (2)	C10—C11—H11	120.5
C8—N3—C9	108.0 (2)	C11—C12—C13	121.5 (3)
C8—N3—S1	123.64 (18)	C11—C12—H12	119.3
C9-N3-S1	126.7(2)	C13 - C12 - H12	119.3
N1-C1-C2	120.7(2) 124.0(3)	C_{14} C_{13} C_{12} C_{12}	121.4(3)
N1 = C1 = C2	124.0(3)	$C_{14} = C_{13} = C_{12}$	110.3
N1 = C1 = C0	110.0(3)	$C_{14} = C_{13} = 1113$	119.3
$C_2 - C_1 - C_0$	120.0(2)		119.5
$C_3 = C_2 = C_1$	120.8 (3)	C13 - C14 - C9	116.6 (3)
С3—С2—Н2	119.6	С13—С14—Н14	121.7
C1—C2—H2	119.6	C9—C14—H14	121.7
C2—C3—C4	118.3 (2)	C20—C15—C16	121.6 (3)
C2—C3—C7	123.5 (2)	C20—C15—S1	119.6 (2)
C4—C3—C7	118.2 (2)	C16—C15—S1	118.8 (2)
C5—C4—C3	122.8 (3)	C17—C16—C15	118.4 (3)
C5—C4—H4	118.6	C17—C16—H16	120.8
C3—C4—H4	118.6	C15—C16—H16	120.8
C4—C5—C6	119.4 (3)	C18—C17—C16	120.2 (3)
С4—С5—Н5	120.3	C18—C17—H17	119.9
C6—C5—H5	120.3	C16—C17—H17	119.9
$N_2 - C_6 - C_5$	124 5 (3)	C19 - C18 - C17	120.8 (3)
N2-C6-C1	116.8 (3)	C19-C18-H18	110.6
12 - 0 - 01	110.0(3) 118.7(2)	$C_{17} = C_{10} = 1110$	119.0
$C_{2} = C_{1} = C_{1}$	10.7(2)	$C_{17} = C_{10} = C_{1110}$	120 4 (2)
0 - 0 - 0	100.2 (2)	10 - 19 - 120	120.4 (5)
$C_{0} - C_{-} - C_{3}$	123.9 (2)	C18—C19—H19	119.8

C10—C7—C3	129.8 (2)	C20—C19—H19	119.8
C7—C8—N3	110.8 (2)	C19—C20—C15	118.6 (3)
С7—С8—Н8	124.6	С19—С20—Н20	120.7
N3—C8—H8	124.6	C15—C20—H20	120.7
N2—Se1—N1—C1	-0.2 (2)	C8—N3—C9—C14	179.1 (3)
N1—Se1—N2—C6	-0.4(2)	S1—N3—C9—C14	13.3 (4)
O1—S1—N3—C8	151.5 (2)	C8—N3—C9—C10	-1.8 (3)
O2—S1—N3—C8	22.0 (3)	S1—N3—C9—C10	-167.7 (2)
C15—S1—N3—C8	-93.0 (2)	C14—C9—C10—C11	0.7 (4)
O1—S1—N3—C9	-44.8 (3)	N3-C9-C10-C11	-178.4 (3)
O2—S1—N3—C9	-174.2 (2)	C14—C9—C10—C7	-179.9 (3)
C15—S1—N3—C9	70.8 (3)	N3—C9—C10—C7	1.0 (3)
Se1—N1—C1—C2	-180.0 (2)	C8—C7—C10—C11	179.5 (3)
Se1—N1—C1—C6	0.7 (3)	C3—C7—C10—C11	0.8 (5)
N1—C1—C2—C3	-178.5 (3)	C8—C7—C10—C9	0.3 (3)
C6—C1—C2—C3	0.8 (4)	C3—C7—C10—C9	-178.5 (3)
C1—C2—C3—C4	0.5 (4)	C9-C10-C11-C12	-0.8 (5)
C1—C2—C3—C7	178.9 (2)	C7—C10—C11—C12	180.0 (3)
C2—C3—C4—C5	-1.3 (4)	C10-C11-C12-C13	-0.1 (6)
C7—C3—C4—C5	-179.8 (3)	C11—C12—C13—C14	1.2 (6)
C3—C4—C5—C6	0.7 (4)	C12—C13—C14—C9	-1.3 (5)
Se1—N2—C6—C5	-178.2 (2)	C10-C9-C14-C13	0.3 (5)
Se1—N2—C6—C1	0.9 (3)	N3-C9-C14-C13	179.3 (3)
C4—C5—C6—N2	179.7 (3)	O1—S1—C15—C20	11.5 (3)
C4—C5—C6—C1	0.6 (4)	O2—S1—C15—C20	145.2 (2)
N1—C1—C6—N2	-1.2 (4)	N3—S1—C15—C20	-103.2 (2)
C2-C1-C6-N2	179.5 (3)	O1—S1—C15—C16	-168.6 (2)
N1—C1—C6—C5	178.0 (3)	O2—S1—C15—C16	-35.0 (2)
C2—C1—C6—C5	-1.4 (4)	N3—S1—C15—C16	76.7 (2)
C2—C3—C7—C8	171.4 (3)	C20-C15-C16-C17	0.8 (4)
C4—C3—C7—C8	-10.2 (4)	S1—C15—C16—C17	-179.0(2)
C2—C3—C7—C10	-10.1 (4)	C15—C16—C17—C18	0.3 (4)
C4—C3—C7—C10	168.3 (3)	C16—C17—C18—C19	-0.9(5)
C10—C7—C8—N3	-1.4 (3)	C17—C18—C19—C20	0.4 (5)
C3—C7—C8—N3	177.4 (2)	C18—C19—C20—C15	0.6 (5)
C9—N3—C8—C7	2.1 (3)	C16—C15—C20—C19	-1.3 (4)

Hydrogen-bond geometry (Å, °)

D—H···A	<i>D</i> —Н	H···A	D····A	<i>D</i> —H··· <i>A</i>
C8—H8…O2 ⁱ	0.93	2.46	3.380 (3)	168
C14—H14…O1	0.93	2.54	3.099 (4)	119

Symmetry code: (i) -x+1, -y, -z+1.

5-[1-(Benzenesulfonyl)-1H-indol-3-yl]-2,1,3-benzothiadiazole (bS)

Crystal data

 $C_{14}H_9N_3S$ $M_r = 251.30$ Orthorhombic, Pbca a = 7.5884(1) Å b = 7.1060 (1) Åc = 43.2464 (7) Å V = 2331.98 (6) Å³ Z = 8F(000) = 1040

Data collection

Xcalibur, Ruby, Gemini ultra R	10906 measured reflections
diffractometer	2072 independent reflections
Radiation source: fine-focus sealed tube	1825 reflections with $I > 2\sigma(I)$
Detector resolution: 5.1856 pixels mm ⁻¹	$R_{\rm int} = 0.025$
ω scans	$\theta_{\rm max} = 67.2^{\circ}, \theta_{\rm min} = 4.1^{\circ}$
Absorption correction: analytical	$h = -9 \longrightarrow 8$
(CrysAlisPro; Rigaku OD, 2020)	$k = -8 \rightarrow 7$
$T_{\min} = 0.663, \ T_{\max} = 0.920$	$l = -51 \rightarrow 49$
Refinement	

Refinement on F^2	Secondary atom site location: dual
Least-squares matrix: full	Hydrogen site location: mixed
$R[F^2 > 2\sigma(F^2)] = 0.035$	H atoms treated by a mixture of independent
$wR(F^2) = 0.098$	and constrained refinement
S = 1.07	$w = 1/[\sigma^2(F_o^2) + (0.0478P)^2 + 0.7413P]$
2072 reflections	where $P = (F_0^2 + 2F_c^2)/3$
167 parameters	$(\Delta/\sigma)_{\rm max} = 0.001$
0 restraints	$\Delta ho_{ m max} = 0.20$ e Å ⁻³
Primary atom site location: dual	$\Delta \rho_{\rm min} = -0.30 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

 $D_{\rm x} = 1.432 \text{ Mg m}^{-3}$

 $\theta = 4.1 - 67.1^{\circ}$

 $\mu = 2.32 \text{ mm}^{-1}$

Plate, colourless

 $0.29 \times 0.18 \times 0.04 \text{ mm}$

T = 295 K

Cu Ka radiation, $\lambda = 1.54184$ Å

Cell parameters from 4620 reflections

Refinement. Structures were solved by the dual-space method of ShelXT (Sheldrick, 2015a) within Olex² (Dolomanov et al., 2009). Structures were refined by the least- squares method implemented in SHELXL (Sheldrick, 2015b) within ShelXle (Hübschle et al., 2011). Structures and crystal packings were visualized using Mercury (Macrae et al., 2020).

Fractional	atomic	coordinates	and is	otropic	or ed	auivalent	isotropi	ic dis	placement	parameters	$(Å^2$	2
1		000.00000000		on opre		100000000000000000000000000000000000000	1001. op1	·• ••••	p	pen ennerers	1	/

	x	У	Z	$U_{ m iso}$ */ $U_{ m eq}$	
S1	1.08880 (6)	0.65005 (9)	0.46249 (2)	0.0684 (2)	
N1	1.05463 (19)	0.6759 (2)	0.42591 (3)	0.0539 (4)	
N3	0.2880 (2)	0.8454 (2)	0.33487 (4)	0.0567 (4)	
H3N	0.206 (3)	0.904 (3)	0.3295 (5)	0.065 (6)*	
C9	0.5346 (2)	0.6723 (2)	0.33531 (4)	0.0369 (3)	
C5	0.6078 (2)	0.7131 (2)	0.39422 (3)	0.0384 (4)	
C6	0.7877 (2)	0.7053 (2)	0.39392 (3)	0.0396 (4)	

H6	0.849325	0.714886	0.375400	0.048*	
C7	0.5027 (2)	0.7429 (2)	0.36611 (4)	0.0395 (4)	
N2	0.8896 (2)	0.6472 (3)	0.47547 (4)	0.0649 (5)	
C14	0.6590(2)	0.5510(2)	0.32215 (4)	0.0453 (4)	
H14	0.750683	0.502873	0.334031	0.054*	
C10	0.3973 (2)	0.7415 (2)	0.31639 (4)	0.0441 (4)	
C1	0.8784 (2)	0.6826 (2)	0.42215 (4)	0.0415 (4)	
C4	0.5157 (2)	0.6949 (3)	0.42314 (4)	0.0480 (4)	
H4	0.393193	0.698955	0.422964	0.058*	
C2	0.7846 (2)	0.6662 (3)	0.45063 (4)	0.0476 (4)	
C3	0.5985 (2)	0.6722 (3)	0.45054 (4)	0.0543 (5)	
H3	0.534867	0.660972	0.468820	0.065*	
C13	0.6439 (3)	0.5037 (3)	0.29140 (4)	0.0567 (5)	
H13	0.725749	0.422248	0.282623	0.068*	
C11	0.3836 (3)	0.6946 (3)	0.28513 (4)	0.0547 (5)	
H11	0.293096	0.742288	0.272936	0.066*	
C8	0.3506 (2)	0.8457 (3)	0.36445 (4)	0.0518 (4)	
H8	0.297698	0.906861	0.381037	0.062*	
C12	0.5077 (3)	0.5761 (3)	0.27307 (4)	0.0594 (5)	
H12	0.501492	0.542837	0.252302	0.071*	

Atomic displacement parameters $(Å^2)$

T 711	1 /22	1/33	1712	1713	1.723
U	0	0	0.2	0.5	0
0.0416 (3)	0.1158 (5)	0.0479 (3)	-0.0007 (3)	-0.00962 (19)	-0.0036 (3)
0.0354 (7)	0.0777 (11)	0.0486 (8)	-0.0024 (7)	-0.0023 (6)	-0.0024 (7)
0.0500 (9)	0.0557 (9)	0.0645 (10)	0.0187 (8)	-0.0130 (7)	0.0043 (7)
0.0349 (8)	0.0340 (7)	0.0417 (8)	-0.0034 (6)	-0.0024 (6)	0.0053 (6)
0.0371 (8)	0.0380 (8)	0.0399 (8)	-0.0005 (6)	0.0003 (6)	-0.0026 (6)
0.0363 (8)	0.0460 (9)	0.0365 (8)	0.0000 (7)	0.0031 (6)	-0.0003 (6)
0.0363 (8)	0.0375 (8)	0.0448 (8)	0.0013 (7)	-0.0009 (6)	0.0014 (7)
0.0484 (9)	0.1062 (14)	0.0401 (8)	0.0008 (9)	-0.0042 (7)	-0.0029 (8)
0.0431 (9)	0.0462 (9)	0.0466 (9)	0.0039 (7)	-0.0008 (7)	0.0019 (7)
0.0456 (9)	0.0373 (8)	0.0494 (9)	-0.0009 (7)	-0.0065 (7)	0.0082 (7)
0.0336 (8)	0.0490 (9)	0.0421 (8)	-0.0013 (7)	0.0012 (6)	-0.0035 (7)
0.0323 (8)	0.0651 (11)	0.0465 (9)	0.0003 (8)	0.0048 (7)	-0.0020 (8)
0.0423 (9)	0.0634 (11)	0.0370 (8)	0.0001 (8)	-0.0005 (7)	-0.0032 (7)
0.0424 (10)	0.0812 (13)	0.0393 (9)	-0.0010 (9)	0.0090 (7)	-0.0006 (8)
0.0638 (12)	0.0558 (11)	0.0504 (10)	0.0032 (9)	0.0050 (9)	-0.0068 (8)
0.0603 (11)	0.0553 (10)	0.0486 (10)	-0.0060 (9)	-0.0165 (8)	0.0119 (8)
0.0473 (10)	0.0537 (10)	0.0543 (10)	0.0137 (8)	-0.0026 (8)	-0.0043 (8)
0.0739 (13)	0.0635 (12)	0.0409 (9)	-0.0103 (11)	-0.0051 (9)	-0.0008(8)
	$\begin{array}{c} U^{11} \\ 0.0416 (3) \\ 0.0354 (7) \\ 0.0500 (9) \\ 0.0349 (8) \\ 0.0371 (8) \\ 0.0363 (8) \\ 0.0363 (8) \\ 0.0484 (9) \\ 0.0431 (9) \\ 0.0431 (9) \\ 0.0436 (8) \\ 0.0323 (8) \\ 0.0323 (8) \\ 0.0423 (9) \\ 0.0424 (10) \\ 0.0638 (12) \\ 0.0603 (11) \\ 0.0473 (10) \\ 0.0739 (13) \end{array}$	U^{11} U^{22} 0.0416 (3) 0.1158 (5) 0.0354 (7) 0.0777 (11) 0.0500 (9) 0.0557 (9) 0.0349 (8) 0.0340 (7) 0.0371 (8) 0.0380 (8) 0.0363 (8) 0.0460 (9) 0.0363 (8) 0.0375 (8) 0.0484 (9) 0.1062 (14) 0.0431 (9) 0.0462 (9) 0.0323 (8) 0.0490 (9) 0.0323 (8) 0.0651 (11) 0.0423 (9) 0.0634 (11) 0.0424 (10) 0.0812 (13) 0.0638 (12) 0.0558 (11) 0.0473 (10) 0.0537 (10) 0.0739 (13) 0.0635 (12)	U^{11} U^{22} U^{33} 0.0416 (3)0.1158 (5)0.0479 (3)0.0354 (7)0.0777 (11)0.0486 (8)0.0500 (9)0.0557 (9)0.0645 (10)0.0349 (8)0.0340 (7)0.0417 (8)0.0371 (8)0.0380 (8)0.0399 (8)0.0363 (8)0.0460 (9)0.0365 (8)0.0363 (8)0.0375 (8)0.0448 (8)0.0484 (9)0.1062 (14)0.0401 (8)0.0431 (9)0.0462 (9)0.0466 (9)0.0356 (8)0.0490 (9)0.0421 (8)0.0323 (8)0.0651 (11)0.0465 (9)0.0423 (9)0.0634 (11)0.0393 (9)0.0638 (12)0.0558 (11)0.0486 (10)0.0603 (11)0.0537 (10)0.0448 (10)0.0473 (10)0.0537 (10)0.0543 (10)0.0739 (13)0.0635 (12)0.0409 (9)	U^{11} U^{22} U^{33} U^{12} 0.0416 (3)0.1158 (5)0.0479 (3)-0.0007 (3)0.0354 (7)0.0777 (11)0.0486 (8)-0.0024 (7)0.0500 (9)0.0557 (9)0.0645 (10)0.0187 (8)0.0349 (8)0.0340 (7)0.0417 (8)-0.0034 (6)0.0371 (8)0.0380 (8)0.0399 (8)-0.0005 (6)0.0363 (8)0.0375 (8)0.0460 (9)0.0365 (8)0.0363 (8)0.0375 (8)0.0448 (8)0.0013 (7)0.0484 (9)0.1062 (14)0.0401 (8)0.0008 (9)0.0431 (9)0.0462 (9)0.0466 (9)-0.0039 (7)0.0363 (8)0.0490 (9)0.0421 (8)-0.0013 (7)0.0323 (8)0.0651 (11)0.0465 (9)0.0003 (8)0.0423 (9)0.0634 (11)0.0370 (8)0.0001 (8)0.0424 (10)0.0812 (13)0.0393 (9)-0.0010 (9)0.0638 (12)0.0558 (11)0.0486 (10)-0.0060 (9)0.0473 (10)0.0537 (10)0.0543 (10)0.0137 (8)0.0739 (13)0.0635 (12)0.0409 (9)-0.0103 (11)	U^{11} U^{22} U^{33} U^{12} U^{13} 0.0416 (3)0.1158 (5)0.0479 (3) $-0.0007 (3)$ $-0.00962 (19)$ 0.0354 (7)0.0777 (11)0.0486 (8) $-0.0024 (7)$ $-0.0023 (6)$ 0.0500 (9)0.0557 (9)0.0645 (10)0.0187 (8) $-0.0130 (7)$ 0.0349 (8)0.0340 (7)0.0417 (8) $-0.0034 (6)$ $-0.0024 (6)$ 0.0371 (8)0.0380 (8)0.0399 (8) $-0.0005 (6)$ 0.0003 (6)0.0363 (8)0.0460 (9)0.0365 (8)0.0000 (7)0.0031 (6)0.0363 (8)0.0375 (8)0.0448 (8)0.0013 (7) $-0.0009 (6)$ 0.0484 (9)0.1062 (14)0.0401 (8)0.0008 (9) $-0.0042 (7)$ 0.0431 (9)0.0462 (9)0.0466 (9) $-0.0039 (7)$ $-0.0008 (7)$ 0.0323 (8)0.0490 (9) $0.0421 (8)$ $-0.0013 (7)$ $0.0012 (6)$ 0.0323 (8)0.0651 (11) $0.0370 (8)$ $0.0001 (8)$ $-0.0005 (7)$ 0.0423 (9) $0.0634 (11)$ $0.0370 (8)$ $0.0001 (8)$ $-0.0005 (7)$ 0.0638 (12) $0.0558 (11)$ $0.0544 (10)$ $0.0032 (9)$ $0.0050 (9)$ 0.0603 (11) $0.0537 (10)$ $0.0490 (9)$ $-0.0137 (8)$ $-0.0026 (8)$ $0.0473 (10)$ $0.0537 (12)$ $0.0409 (9)$ $-0.0103 (11)$ $-0.026 (8)$

Geometric parameters (Å, °)

S1—N2	1.6128 (17)	N2—C2	1.344 (2)
S1—N1	1.6136 (16)	C14—C13	1.376 (2)
N1—C1	1.348 (2)	C14—H14	0.9300

N3—C8	1.365 (2)	C10-C11	1.396 (2)
N3—C10	1.368 (2)	C1—C2	1.428 (2)
N3—H3N	0.79 (2)	C4—C3	1.351 (2)
C9-C14	1 399(2)	C4—H4	0.9300
C_{0}	1.377(2)	$C_1 = C_2$	1,412,(2)
C9—C10	1.413 (2)		1.415 (5)
C9_C7	1.444 (2)	C3—H3	0.9300
C5—C6	1.367 (2)	C13—C12	1.400 (3)
C5—C4	1.438 (2)	С13—Н13	0.9300
C5—C7	1.469 (2)	C11—C12	1.366 (3)
C6—C1	1.411 (2)	C11—H11	0.9300
С6—Н6	0.9300	С8—Н8	0.9300
C7—C8	1 368 (2)	C12—H12	0.9300
07 00	1.500 (2)		0.9500
NO SI NI	101 07 (9)	N1 C1 C6	126 26 (15)
$N_2 = S_1 = N_1$	101.07(6)	NI = CI = CO	120.30 (13)
CI-NI-SI	106.37 (12)	NI - CI - C2	112.81 (15)
C8—N3—C10	109.73 (15)	C6-C1-C2	120.82 (15)
C8—N3—H3N	123.5 (16)	C3—C4—C5	123.18 (16)
C10—N3—H3N	126.6 (16)	C3—C4—H4	118.4
C14—C9—C10	118.42 (15)	C5—C4—H4	118.4
C14—C9—C7	134.62 (15)	N2—C2—C3	126.70 (17)
С10—С9—С7	106.86 (14)	N2—C2—C1	113.72 (16)
C6C5C4	119.35 (15)	$C_{3}-C_{2}-C_{1}$	119.58 (15)
C6-C5-C7	122 68 (14)	C4 - C3 - C2	118 11 (16)
C_{1} C_{2} C_{3} C_{4}	122.00(14) 117.07(14)	$C_4 = C_5 = C_2$	120.0
$C_{4} C_{5} C_{7}$	117.97(14)	C_{4}	120.9
	118.95 (14)	С2—С3—Н3	120.9
С5—С6—Н6	120.5	C14-C13-C12	121.24 (18)
С1—С6—Н6	120.5	C14—C13—H13	119.4
C8—C7—C9	106.16 (14)	С12—С13—Н13	119.4
C8—C7—C5	125.33 (15)	C12—C11—C10	117.74 (17)
С9—С7—С5	128.51 (14)	C12—C11—H11	121.1
C2—N2—S1	106.03 (13)	C10-C11-H11	121.1
C13—C14—C9	119.16 (16)	N3—C8—C7	110.01 (16)
C13—C14—H14	120.4	N3—C8—H8	125.0
C9-C14-H14	120.4	С7—С8—Н8	125.0
N_3 C_{10} C_{11}	130 51 (16)	C_{11} C_{12} C_{13}	121.0 121.26(17)
$N_2 = C_{10} = C_0$	107.22(15)		110.4
	107.23(13)		119.4
C11—C10—C9	122.17 (16)	C13-C12-H12	119.4
	0.05 (15)		
N2 - S1 - N1 - C1	-0.35 (15)	C5-C6-C1-N1	-1/8.66 (17)
C4—C5—C6—C1	-1.0(2)	C5—C6—C1—C2	0.5 (2)
C7—C5—C6—C1	177.87 (14)	C6—C5—C4—C3	0.8 (3)
C14—C9—C7—C8	-175.18 (18)	C7—C5—C4—C3	-178.16 (17)
C10—C9—C7—C8	0.92 (18)	S1—N2—C2—C3	-179.70 (18)
C14—C9—C7—C5	4.5 (3)	S1—N2—C2—C1	-0.2 (2)
C10—C9—C7—C5	-179.43 (15)	N1—C1—C2—N2	-0.1(2)
C6-C5-C7-C8	-143.47 (18)	C6-C1-C2-N2	-179.37 (16)
C4-C5-C7-C8	354(2)	N1-C1-C2-C3	179 49 (18)
$C_{1} C_{2} C_{1} C_{2} C_{3}$	35.7(2)	$C_{1} C_{1} C_{2} C_{3}$	(10)
U - U - U	JU.J (J)	0 - 0 - 0 - 0 - 0 - 0 - 0 - 0 - 0 - 0 -	0.2 (3)

C4—C5—C7—C9	-144.19 (17)	C5—C4—C3—C2	0.0 (3)
N1—S1—N2—C2	0.31 (16)	N2-C2-C3-C4	179.05 (19)
C10-C9-C14-C13	0.2 (2)	C1—C2—C3—C4	-0.4 (3)
C7—C9—C14—C13	175.93 (18)	C9—C14—C13—C12	0.6 (3)
C8—N3—C10—C11	176.87 (18)	N3-C10-C11-C12	-175.41 (19)
C8—N3—C10—C9	0.3 (2)	C9-C10-C11-C12	0.7 (3)
C14—C9—C10—N3	176.06 (14)	C10—N3—C8—C7	0.3 (2)
C7—C9—C10—N3	-0.78 (18)	C9—C7—C8—N3	-0.7 (2)
C14—C9—C10—C11	-0.8 (2)	C5-C7-C8-N3	179.60 (15)
C7—C9—C10—C11	-177.65 (15)	C10-C11-C12-C13	0.1 (3)
S1—N1—C1—C6	179.53 (14)	C14—C13—C12—C11	-0.7 (3)
S1—N1—C1—C2	0.27 (19)		

5-(1H-Indol-3-yl)-2,1,3-benzoxadiazole (bO)

Crystal data

C₁₄H₉N₃O $M_r = 235.24$ Orthorhombic, *Pbca* a = 12.0256 (7) Å b = 7.7396 (5) Å c = 23.8551 (16) Å V = 2220.3 (2) Å³ Z = 8F(000) = 976

Data collection

Xcalibur, Ruby, Gemini ultra diffractometer Radiation source: fine-focus sealed X-ray tube, Enhance Ultra (Cu) X-ray Source Detector resolution: 5.1856 pixels mm⁻¹ ω scans Absorption correction: analytical (CrysAlisPro; Rigaku OD, 2020) $T_{\min} = 0.941, T_{\max} = 0.981$

Refinement

Refinement on F^2 Least-squares matrix: full $R[F^2 > 2\sigma(F^2)] = 0.050$ $wR(F^2) = 0.134$ S = 1.061972 reflections 166 parameters 0 restraints Primary atom site location: dual

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

 $D_x = 1.407 \text{ Mg m}^{-3}$ Cu K α radiation, $\lambda = 1.54184 \text{ Å}$ Cell parameters from 1395 reflections $\theta = 7.4-66.3^{\circ}$ $\mu = 0.75 \text{ mm}^{-1}$ T = 295 KPlate, clear yellow $0.12 \times 0.09 \times 0.03 \text{ mm}$

6883 measured reflections 1972 independent reflections 1275 reflections with $I > 2\sigma(I)$ $R_{int} = 0.047$ $\theta_{max} = 67.1^{\circ}, \ \theta_{min} = 3.7^{\circ}$ $h = -14 \rightarrow 12$ $k = -6 \rightarrow 9$ $l = -28 \rightarrow 19$

Secondary atom site location: dual Hydrogen site location: mixed H atoms treated by a mixture of independent and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0633P)^2 + 0.0873P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{max} < 0.001$ $\Delta\rho_{max} = 0.14$ e Å⁻³ $\Delta\rho_{min} = -0.16$ e Å⁻³ **Refinement**. Structures were solved by the dual-space method of ShelXT (Sheldrick, 2015a) within Olex² (Dolomanov *et al.*, 2009). Structures were refined by the least- squares method implemented in SHELXL (Sheldrick, 2015b) within ShelXle (Hübschle *et al.*, 2011). Structures and crystal packings were visualized using Mercury (Macrae *et al.*, 2020).

	x	У	Ζ	$U_{ m iso}$ */ $U_{ m eq}$
01	0.26211 (19)	0.2142 (3)	0.43917 (8)	0.0909 (6)
N1	0.1692 (2)	0.2567 (3)	0.46968 (9)	0.0801 (7)
N2	0.3589 (2)	0.2507 (3)	0.46824 (9)	0.0822 (7)
N3	0.52478 (16)	0.6406 (3)	0.70573 (9)	0.0640 (6)
H3N	0.581 (2)	0.696 (3)	0.7137 (11)	0.077*
C1	0.2075 (2)	0.3199 (3)	0.51701 (10)	0.0629 (6)
C2	0.3259 (2)	0.3161 (3)	0.51611 (10)	0.0616 (6)
C3	0.3883 (2)	0.3735 (3)	0.56268 (10)	0.0610 (6)
H3A	0.465597	0.368956	0.562224	0.073*
C4	0.33330 (18)	0.4356 (3)	0.60815 (9)	0.0520 (6)
C5	0.21290 (18)	0.4396 (3)	0.60788 (10)	0.0566 (6)
Н5	0.176549	0.482253	0.639392	0.068*
C6	0.1509 (2)	0.3849 (3)	0.56443 (10)	0.0658 (7)
H6	0.073675	0.389643	0.565567	0.079*
C7	0.39274 (17)	0.5038 (3)	0.65679 (9)	0.0506 (5)
C8	0.49217 (18)	0.5924 (3)	0.65382 (10)	0.0610 (6)
H8	0.530990	0.615615	0.620952	0.073*
C9	0.36413 (16)	0.4987 (2)	0.71516 (9)	0.0473 (5)
C10	0.44927 (17)	0.5855 (3)	0.74484 (10)	0.0513 (6)
C11	0.4497 (2)	0.6039 (3)	0.80242 (11)	0.0624 (6)
H11	0.506433	0.663393	0.820582	0.075*
C12	0.3635 (2)	0.5313 (3)	0.83195 (11)	0.0654 (7)
H12	0.361933	0.541208	0.870791	0.078*
C13	0.27815 (19)	0.4427 (3)	0.80449 (11)	0.0606 (6)
H13	0.220822	0.394500	0.825468	0.073*
C14	0.27709 (17)	0.4252 (3)	0.74691 (10)	0.0527 (5)
H14	0.219735	0.365798	0.729246	0.063*

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (A^2)

Atomic displacement parameters $(Å^2)$

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
01	0.1045 (15)	0.1109 (16)	0.0571 (10)	-0.0033 (12)	-0.0023 (11)	-0.0055 (10)
N1	0.0843 (15)	0.0991 (17)	0.0569 (14)	-0.0107 (13)	-0.0056 (13)	0.0038 (12)
N2	0.0843 (16)	0.1040 (17)	0.0582 (14)	0.0035 (13)	0.0010 (12)	-0.0021 (12)
N3	0.0455 (10)	0.0674 (13)	0.0791 (15)	-0.0127 (9)	-0.0027 (10)	0.0060 (11)
C1	0.0670 (15)	0.0709 (15)	0.0507 (15)	-0.0074 (12)	-0.0041 (12)	0.0099 (12)
C2	0.0682 (15)	0.0693 (15)	0.0473 (14)	0.0029 (12)	0.0075 (12)	0.0066 (12)
C3	0.0510(13)	0.0737 (15)	0.0585 (15)	0.0027 (12)	0.0037 (12)	0.0057 (12)
C4	0.0474 (11)	0.0533 (12)	0.0552 (13)	-0.0006 (10)	0.0027 (11)	0.0089 (11)
C5	0.0466 (12)	0.0683 (14)	0.0547 (14)	0.0018 (11)	0.0013 (11)	0.0035 (11)
C6	0.0506 (13)	0.0811 (17)	0.0655 (16)	-0.0038 (12)	-0.0020 (12)	0.0079 (13)

C7	0.0437 (11)	0.0512 (11)	0.0570 (14)	0.0000 (10)	-0.0030 (10)	0.0048 (11)
C8	0.0485 (12)	0.0708 (14)	0.0636 (15)	-0.0056 (11)	0.0031 (11)	0.0118 (12)
C9	0.0414 (10)	0.0439 (10)	0.0567 (13)	0.0029 (9)	0.0002 (10)	0.0028 (10)
C10	0.0440 (11)	0.0474 (12)	0.0626 (15)	0.0012 (10)	-0.0038 (11)	0.0010 (11)
C11	0.0602 (14)	0.0566 (13)	0.0704 (16)	0.0025 (12)	-0.0105 (13)	-0.0099 (12)
C12	0.0720 (16)	0.0630 (15)	0.0613 (15)	0.0119 (13)	0.0023 (13)	-0.0066 (12)
C13	0.0551 (13)	0.0618 (14)	0.0648 (15)	0.0076 (11)	0.0122 (12)	0.0040 (12)
C14	0.0424 (11)	0.0514 (12)	0.0642 (14)	0.0018 (10)	0.0023 (11)	0.0031 (11)

Geometric parameters (Å, °)

O1—N1	1.373 (3)	С5—Н5	0.9300	
O1—N2	1.384 (3)	С6—Н6	0.9300	
N1—C1	1.314 (3)	C7—C8	1.380 (3)	
N2—C2	1.310 (3)	С7—С9	1.435 (3)	
N3—C8	1.351 (3)	C8—H8	0.9300	
N3—C10	1.370 (3)	C9—C14	1.411 (3)	
N3—H3N	0.83 (3)	C9—C10	1.415 (3)	
C1—C6	1.413 (3)	C10—C11	1.381 (3)	
C1—C2	1.424 (4)	C11—C12	1.374 (3)	
C2—C3	1.413 (3)	C11—H11	0.9300	
C3—C4	1.358 (3)	C12—C13	1.397 (4)	
С3—НЗА	0.9300	C12—H12	0.9300	
C4—C5	1.448 (3)	C13—C14	1.380 (3)	
C4—C7	1.461 (3)	C13—H13	0.9300	
C5—C6	1.345 (3)	C14—H14	0.9300	
N1—O1—N2	111.69 (18)	C8—C7—C9	105.7 (2)	
C1—N1—O1	105.1 (2)	C8—C7—C4	124.2 (2)	
C2—N2—O1	105.1 (2)	C9—C7—C4	130.03 (19)	
C8—N3—C10	110.23 (19)	N3—C8—C7	110.0 (2)	
C8—N3—H3N	126.4 (19)	N3—C8—H8	125.0	
C10—N3—H3N	123.3 (19)	С7—С8—Н8	125.0	
N1—C1—C6	130.7 (2)	C14—C9—C10	117.4 (2)	
N1—C1—C2	109.2 (2)	C14—C9—C7	135.2 (2)	
C6—C1—C2	120.1 (2)	C10—C9—C7	107.39 (18)	
N2—C2—C3	130.2 (2)	N3—C10—C11	130.0 (2)	
N2—C2—C1	108.9 (2)	N3—C10—C9	106.7 (2)	
C3—C2—C1	120.8 (2)	C11—C10—C9	123.3 (2)	
C4—C3—C2	118.7 (2)	C12-C11-C10	117.7 (2)	
C4—C3—H3A	120.6	C12—C11—H11	121.1	
С2—С3—Н3А	120.6	C10—C11—H11	121.1	
C3—C4—C5	119.4 (2)	C11—C12—C13	121.0 (2)	
C3—C4—C7	121.6 (2)	C11—C12—H12	119.5	
C5—C4—C7	119.0 (2)	C13—C12—H12	119.5	
C6—C5—C4	123.4 (2)	C14—C13—C12	121.4 (2)	
С6—С5—Н5	118.3	C14—C13—H13	119.3	
С4—С5—Н5	118.3	C12—C13—H13	119.3	

C5—C6—C1	117.5 (2)	C13—C14—C9	119.2 (2)
C5—C6—H6	121.2	C13—C14—H14	120.4
С1—С6—Н6	121.2	С9—С14—Н14	120.4
N2-01-N1-C1	0.4 (3)	C5—C4—C7—C9	35.3 (3)
N1-01-N2-C2	-0.4 (3)	C10—N3—C8—C7	0.3 (3)
01-N1-C1-C6	179.9 (3)	C9—C7—C8—N3	-0.1 (2)
01-N1-C1-C2	-0.3 (3)	C4—C7—C8—N3	179.06 (19)
01—N2—C2—C3	178.7 (2)	C8—C7—C9—C14	-177.6 (2)
01—N2—C2—C1	0.2 (3)	C4—C7—C9—C14	3.3 (4)
N1—C1—C2—N2	0.0 (3)	C8—C7—C9—C10	-0.1 (2)
$C6-C1-C2-N2 \\N1-C1-C2-C3 \\C6-C1-C2-C3 \\N2-C2-C3-C4$	179.9 (2)	C4—C7—C9—C10	-179.2 (2)
	-178.6 (2)	C8—N3—C10—C11	179.1 (2)
	1.2 (3)	C8—N3—C10—C9	-0.4 (2)
	-179.5 (3)	C14—C9—C10—N3	178.34 (18)
$\begin{array}{c} C1 - C2 - C3 - C4 \\ C1 - C2 - C3 - C4 \\ C2 - C3 - C4 - C5 \\ C2 - C3 - C4 - C7 \end{array}$	-1.1 (3)	C7—C9—C10—N3	0.3 (2)
	0.6 (3)	C14—C9—C10—C11	-1.2 (3)
	-177.4 (2)	C7—C9—C10—C11	-179.3 (2)
C3-C4-C5-C6	-0.1 (3)	N3-C10-C11-C12	-178.4 (2)
C7-C4-C5-C6	177.9 (2)	C9-C10-C11-C12	1.0 (3)
C4-C5-C6-C1	0.2 (3)	C10-C11-C12-C13	-0.3 (3)
N1-C1-C6-C5	179.0 (3)	C11-C12-C13-C14	-0.1 (3)
$\begin{array}{c} C2-C1-C6-C5\\ C3-C4-C7-C8\\ C5-C4-C7-C8\\ C3-C4-C7-C9\\ \end{array}$	-0.8 (3) 34.3 (3) -143.6 (2) -146.7 (2)	C12—C13—C14—C9 C10—C9—C14—C13 C7—C9—C14—C13	-0.1 (3) 0.7 (3) 178.0 (2)