

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

Hormonal therapies and venous thrombosis

Douxfils, Jonathan; Morimont, Laure; Creinin, Mitchell D; Gaspard, Ulysse; Foidart, Jean-Michel

Published in:

Research and Practice in Thrombosis and Haemostasis

DOI:

[10.1016/j.rpth.2023.100155](https://doi.org/10.1016/j.rpth.2023.100155)

Publication date:

2023

Document Version

Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (HARVARD):

Douxfils, J, Morimont, L, Creinin, MD, Gaspard, U & Foidart, J-M 2023, 'Hormonal therapies and venous thrombosis: considerations for prevention and management-a reappraisal', *Research and Practice in Thrombosis and Haemostasis*, vol. 7, no. 4, 100155, pp. 100155. <https://doi.org/10.1016/j.rpth.2023.100155>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LETTER TO THE EDITOR

Hormonal therapies and venous thrombosis: considerations for prevention and management—a reappraisal

To the Editor,

We appreciate the response of LaVasseur et al. [1] to our Letter to the Editor on their article on the risk of venous thromboembolism (VTE) with hormonal therapies, in which they remark on a potential link of drospirenone (DRSP) with an increased risk of thromboembolism because of its antiminerocorticoid effect, which could interfere with the effect of aldosterone on hemostasis. These observations are debated in the literature [2] because aldosterone administration enhances thrombus burden, an effect that was prevented by cotreatment with eplerenone. Additionally, aldosterone treatment decreases bleeding time, induces platelet activation and degranulation, and increases the expression of plasminogen activator inhibitor-1, whereas antiminerocorticoids increase bleeding time, decrease platelet activation, and inhibit the expression of plasminogen activator inhibitor-1, fibrinogen, P-selectin, and interleukin-1 β in a variety of rodent models [2]. Further, a recent retrospective cohort study reported that spironolactone treatment in adults with acne did not increase the risk of VTE (odds ratio, 0.57; 95% CI, 0.31-1.06) or pulmonary embolism (odds ratio, 0.60; 95% CI, 0.26-1.37) compared with that in patients being treated with tetracycline antibiotics [3]. DRSP, as a progestin-only contraceptive at a dose of 4 mg, has not been categorized by regulatory bodies as having a different risk of VTE from other progestin-only contraceptives. In addition, hemostatic investigations of individual coagulation factors did not reveal the clinically relevant impact of this DRSP at 4 mg (Figure A) [4]. Interestingly, estetrol (E4), in combination with DRSP, has a coagulation profile similar to DRSP alone, whereas a clear difference is observed with the ethinylestradiol (EE)/DRSP combination.

Evidence is also accumulating regarding the correlation between coagulation markers, such as thrombin generation-based assays, and the risk of VTE observed in phase 4 studies. Haverinen et al. [5] also reported that thrombin generation was lower after exposure to estradiol valerate/dienogest compared with EE/dienogest, the latter being associated with a higher risk of VTE compared with EE/levonorgestrel (adjusted hazard ratio, 1.57; 95% CI, 1.07-2.30) [6], whereas the former has a similar lower risk of VTE (adjusted hazard ratio, 0.40; 95% CI, 0.20-1.10) [7]. We also observed that EE-containing combined oral contraceptives (COCs) have higher

thrombin generation than E4/DRSP and that the magnitude of the impact of EE-containing COCs can be considered clinically relevant (Figure B).

Lastly, the plasma coagulation inhibitor subcommittee of the ISTH SSC has recently recommended the use of the endogenous thrombin potential-based activated protein C (APC) resistance assay to evaluate hormone-induced acquired APC resistance, an independent risk factor for VTE [8]. The model presented in our initial response is in line with this expert recommendation because we showed that acquired APC resistance is correlated with the risk of VTE with different COC preparations. This validates our observation that E4/DRSP leads to lower acquired APC resistance compared with EE-containing COC, reducing the burden of this independent risk factor [9].

Therefore, we reiterate our conclusion that the choice of estrogen matters when considering the risk of VTE and that classification of COC should be reappraised considering the latest scientific evidence.

Jonathan Douxfils PharmD, PhD^{1,2} 

Laure Morimont PharmD, PhD^{1,2}

Mitchell D. Creinin MD³

Ulysse Gaspard MD, PhD⁴

Jean-Michel Foidart MD, PhD^{5,6}

¹Qualiblood sa, Namur, Belgium

²Department of Pharmacy, Faculty of Medicine, University of Namur, Namur Research Institute for Life Sciences, Clinical Pharmacology Research Group, Namur, Belgium

³Department of Obstetrics and Gynecology, University of California, Davis, Sacramento, California, USA

⁴Department of Obstetrics and Gynecology, University of Liège, Belgium

⁵Estetra SRL, An affiliate Company of Mithra Pharmaceuticals, Liège, Belgium

⁶University of Liège, Liège, Belgium

Handling Editor: Dr Cihan Ay




FIGURE (A) A comparison of the absolute change from baseline in a common series of coagulation factors between 75-µg desogestrel and 4-mg drospirenone; 2 progestin-only contraceptives; and the combined oral contraceptives (COCs) 15-mg estetrol (E4) + 3-mg drospirenone, 20-µg ethinylestradiol + 3-mg drospirenone, and 30-µg ethinylestradiol + 150-µg levonorgestrel. Different profiles were observed for products containing ethinylestradiol, whereas progestin-only contraceptives and E4-containing pills are either neutral or have decreased hypercoagulability. (B) Comparison of thrombin generation profiles of ethinylestradiol-containing COCs versus those of E4-containing COCs. Products that contain ethinylestradiol showed mean thrombograms outside the reference range, whereas E4 does not clinically impact thrombin generation. COC, combined oral contraceptive; EE, ethinylestradiol.

Correspondence

Jonathan Douxfils, Qualiblood sa, Namur, and University of Namur, Faculty of Medicine, Department of Pharmacy, Namur Research Institute for Life Sciences, Clinical Pharmacology Research Group, Rue du Séminaire 20a, 5000 Namur, Belgium.
Email: jonathan.douxfils@qualiblood.eu

REFERENCES

- [1] LaVasseur C, Shatzel J, Kartika T. Re: hormonal therapies and venous thrombosis: the estrogen matters! Morimont et al. *Res Pract Thromb Haemost.* 2023;7:100029. <https://doi.org/10.1016/j.rpth.2022.100029>
- [2] Moss ME, Jaffe IZ. Mineralocorticoid receptors in the pathophysiology of vascular inflammation and atherosclerosis. *Front Endocrinol.* 2015;6:153. <https://doi.org/10.3389/fendo.2015.00153>
- [3] Shields A, Flood K, Barbieri JS. Spironolactone use for acne is not associated with an increased risk of venous thromboembolism: a matched, retrospective cohort study. *J Am Acad Dermatol.* 2023. <https://doi.org/10.1016/j.jaad.2023.02.028>

TWITTER

Jonathan Douxfils @douxfilj

- [4] Palacios S, Colli E, Regidor PA. Metabolic and laboratory effects of a progestin-only pill containing drospirenone 4 mg in comparison to desogestrel 75 microg: a double-blind, double-dummy, prospective, randomised study. *Eur J Contracept Reprod Health Care*. 2021;26:454–61. <https://doi.org/10.1080/13625187.2021.1957094>
- [5] Haverinen AH, Luiro KM, Szanto T, Kangasniemi MH, Hiltunen L, Sainio S, et al. Combined oral contraceptives containing estradiol valerate vs ethinylestradiol on coagulation: a randomized clinical trial. *Acta Obstet Gynecol Scand*. 2022;101:1102–11. <https://doi.org/10.1111/aogs.14428>
- [6] Dinger J. COCs containing dienogest and 30 µg ethinylestradiol may carry a higher VTE risk compared with corresponding preparations with levonorgestrel: a meta-analysis of four large cohort studies. *Front Womens Health*. 2020;5:5. <https://doi.org/10.15761/fwh.1000175>
- [7] Grandi G, Facchinetti F, Bitzer J. Confirmation of the safety of combined oral contraceptives containing oestradiol on the risk of venous thromboembolism. *Eur J Contracept Reprod Health Care*. 2022;27:83–4. <https://doi.org/10.1080/13625187.2022.2029397>
- [8] Moore GW, Castoldi E, Teruya J, Morishita E, Adcock DM. Factor V Leiden-independent activated protein C resistance: communication from the plasma coagulation inhibitors subcommittee of the International Society on Thrombosis and Haemostasis Scientific and Standardisation Committee. *J Thromb Haemost*. 2023;21:164–74. <https://doi.org/10.1016/j.jth.2022.10.014>
- [9] Morimont L, Creinin MD, Gaspard U, Foidart JM, Douxfils J. Hormonal therapies and venous thrombosis: estrogen matters. *Res Pract Thromb Haemost*. 2023;7:100021. <https://doi.org/10.1016/j.rpth.2022>