

## RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

### Quasi-ordered photonic structures colour the bluespotted ribbontail ray

Bouchat, J.; Cortesi, F.; Cheney, K.; Vukusic, P.; Justin Marshall, N.; Deparis, O.; Mouchet, S. R.

*Published in:*

International Conference on Metamaterials, Photonic Crystals and Plasmonics

*Publication date:*

2023

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication](#)

*Citation for pulished version (HARVARD):*

Bouchat, J, Cortesi, F, Cheney, K, Vukusic, P, Justin Marshall, N, Deparis, O & Mouchet, SR 2023, Quasi-ordered photonic structures colour the bluespotted ribbontail ray. in *International Conference on Metamaterials, Photonic Crystals and Plasmonics*. International Conference on Metamaterials, Photonic Crystals and Plasmonics, pp. 1126-1127, 13th International Conference on Metamaterials, Photonic Crystals and Plasmonics, META 2023, Paris, France, 18/07/23.

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Quasi-ordered photonic structures colour the bluespotted ribbontail ray

J. Bouchat<sup>1</sup>, F. Cortesi<sup>2</sup>, K. Cheney<sup>2,3</sup>, P. Vukusic<sup>4</sup>, N. Justin Marshall<sup>2</sup>, O. DeParis<sup>1</sup>, and S. R. Mouchet<sup>1,2,4\*</sup>

<sup>1</sup>Department of Physics and Namur Institute of Structured Matter (NISM), University of Namur, Rue de Bruxelles 61, 5000 Namur, Belgium

<sup>2</sup>Queensland Brain Institute, The University of Queensland, Brisbane 4072, Queensland, Australia

<sup>3</sup>School of Biological Sciences, The University of Queensland, Brisbane 4072, Queensland, Australia

<sup>4</sup>School of Physics, University of Exeter, Stocker Road, Exeter EX4 4QL, UK

\*corresponding author: [sebastien.mouchet@unamur.be](mailto:sebastien.mouchet@unamur.be)

**Abstract:** Due to the scarcity of blue colour exhibited by natural organisms, highlighting the underlying this colour mechanisms is always very impactful for the understanding of the natural world. In this research, the colour of the blue rounded spots occurring in the skin of *Taeniura lymma* stingray was unveiled by a combination of experimental and numerical techniques. Our results demonstrated that this blue colour arises from coherent scattering in quasi-ordered photonic structures occurring in the skin of this stingray.

Blue is often described as the rarest colour in natural organisms [1]. Elucidating the underlying mechanism(s) giving rise to such a hue is always very impactful for the understanding of the natural world in which human beings live. In this research, the colour of the blue rounded spots occurring in the skin of the bluespotted ribbontail ray *Taeniura lymma* was unveiled by a combination of experimental and numerical techniques. Histological observations were carried out with optical, fluorescence and transmission electron microscopies. Optical characterisations were performed by spectrometry and microspectrofluorimetry. Numerical simulations were based on two-dimensional fast Fourier transforms of electron micrographs of the observed structures and Benedek's theory relating Fourier transform of spatial variation in refractive index to the intensity of coherent scattering [2]. Our results demonstrated that the blue colour of this ray arises from coherent scattering in quasi-ordered photonic structures occurring in the skin of this animal. This type of structures made of collagen fibres is mostly unknown in marine species. In addition, structural blue colours had never been reported in elasmobranches.



Figure 1. The blue spots occurring in the integuments of *T. lymma* arise from quasi-ordered photonic structures. Photograph by Taken reproduced from

<https://pixabay.com/photos/ray-stingray-fish-sea-ocean-539788/>

**Acknowledgements**

SRM was supported by the Belgian National Fund for Scientific Research (FRS-FNRS) (91400/1.B.309.18F), the Maturation Fund of the Walloon Region, and a BEWARE Fellowship (Convention n°2110034) of the Walloon Region (Marie Skłodowska-Curie Actions of the European Union - COFUND - contract~847587), as a Postdoctoral Researcher.

**References**

1. Mouchet, S. R. and O. Deparis, *Natural Photonics & Bioinspiration*, Boston: Artech House, 2021.
2. Benedek, G. B., "Theory of Transparency of the Eye," *Applied Optics*, Vol. 10, No. 3, 459-473, 1971.