

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

Exponentially long time stability near an equilibrium point for non-linearizable analytic vector fields

Carletti, Timoteo

Published in:

Zeitschrift für angewandte Mathematik und Physik

Publication date:

2005

Document Version

Early version, also known as pre-print

[Link to publication](#)

Citation for pulished version (HARVARD):

Carletti, T 2005, 'Exponentially long time stability near an equilibrium point for non-linearizable analytic vector fields', *Zeitschrift für angewandte Mathematik und Physik*, vol. 56, pp. 1-13.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Exponentially long time stability near an equilibrium point for non-linearizable analytic vector fields

Timoteo Carletti

Abstract. We study the orbit behaviour of a germ of an analytic vector field of $(\mathbb{C}^n, 0)$, $n \geq 2$. We prove that if its linear part is semisimple, non-resonant and verifies a Bruno-like condition, then the origin is effectively stable: stable for finite but exponentially long times.

Mathematics Subject Classification (2000). Primary 37C75, 34A25.

Keywords. linearization vector field, Gevrey class, Bruno condition, effective stability, Nekhoroshev theorem.

1. Introduction

Let us consider the germ of analytic vector field, $X_F = \sum_{1 \leq j \leq n} F_j(z) \frac{\partial}{\partial z_j}$, of $(\mathbb{C}^n, 0)$ $n \geq 2$, whose components $(F_j)_{1 \leq j \leq n}$ are analytic functions in a neighbourhood of the origin, vanishing at $0 \in \mathbb{C}^n$.

Let us consider the associated Ordinary Differential Equation:

$$\frac{dz}{dt} = F(z); \quad (1.1)$$

under the above assumptions $z(t; 0) = 0$ for all t is an equilibrium solution¹! We are interested in studying the stability of orbits of X_F in a neighbourhood of this equilibrium point.

We use the standard definition of *stability* (see [12]) for an equilibrium solution: $z = 0$ it is stable in the past and in the future, if for any neighbourhood U of 0 there exists a neighbourhood V , containing the origin, s.t. $z(0; z_0) \in V$ implies $z(t; z_0) \in U$ for all $t \in \mathbb{R}$.

In a coordinates system centred at the equilibrium point, the j -th component of the vector field will take the form: $F_j(z) = (Az)_j + f_j(z)$, with A a $n \times n$ complex matrix and f_j analytic function such that $f_j(0) = Df_j(0) = 0$, for all $1 \leq j \leq n$.

¹ Here and throughout the paper by $z(t; z_0)$ we mean the solution at time t of (1.1) s.t. $z(0; z_0) = z_0$. When the value of z_0 will not be relevant we'll just write $z(t)$.

Following the idea of Poincaré to study the orbit of (1.1) in a neighbourhood of the origin, we will try to find an analytic change of coordinates, through an analytic diffeomorphisms $z \mapsto H(z) = w$, the *linearization*, s.t. in the new coordinates the vector field X_F is conjugate to its linear part: $H^*X_F H^{-1} = X_A$, where $X_A = \sum (Az)_j \frac{\partial}{\partial z_j}$. Hence equation (1.1) rewrites:

$$\frac{dw}{dt} = Aw. \quad (1.2)$$

This change of coordinates must solve:

$$AH(z) = DH(z) \cdot (Az + f(z)), \quad (1.3)$$

and it is unique by assuming $DH(0) = \mathbb{I}$.

Clearly if the linear system (1.2) is stable and (1.1) is analytically linearizable, then also the latter is stable. It is a remarkable result that this condition is also necessary, as the following Theorem states:

Theorem 1.1. (Carathéodory–Cartan 1932) *Necessary and sufficient condition for the stability of the solution $z = 0$ of (1.1) for all real t is that:*

1. *A is diagonalizable with purely imaginary eigenvalues;*
2. *there exists an holomorphic function $z = K(w) = w + \mathcal{O}(|w|^2)$, $w \in \mathbb{C}^n$, which brings (1.1) into the linear system:*

$$\frac{dw}{dt} = Aw.$$

The problem is classical but for the shake of completeness we briefly recall some known results. As already mentioned the problem has been considered by Poincaré in his Thesis [13], where also a first positive result has been provided: there exists a germ of analytic change of coordinates, $z \mapsto H(z)$ which brings (1.1) into the associated linear one, if the eigenvalues $(\omega_j)_{1 \leq j \leq n}$ of $DF(0)$ are *non-resonant*, i.e. $\alpha \cdot \omega \neq \omega_j$, for all $\alpha \in \mathbb{N}^n$ s.t. $|\alpha| = \alpha_1 + \dots + \alpha_n \geq 2$ and for all $j \in \{1, \dots, n\}$, and if the eigenvalues belong to the *Poincaré domain*, i.e. the *complex hull* of $(\omega_j)_{1 \leq j \leq n}$, in \mathbb{C} , does not contain the origin.

Removing the non-resonance hypothesis, one could not in general bring the full system into a linear one, even at formal level. Dulac [7] proved that if the eigenvalues belong to the Poincaré domain then system (1.1) is analytically conjugated to a polynomial *normal form*, i.e. to a particular vector field containing only resonant monomials, moreover this polynomial is a linear one if the eigenvalues are non-resonant. Recently a new proof of these two statements has been given [6] by adapting the smooth normalization argument used by Sternberg [16] and using a geometrical interpretation of the Poincaré domain.

When A belongs to the *Siegel domain*, i.e. the origin is contained in the convex hull of the eigenvalues plotted as points in the complex plane, the situation is harder because *small divisors* are involved: the existence of an analytic linearization is strictly related to the arithmetic property of approximation of the vector $\omega = (\omega_1, \dots, \omega_n)$, with vectors of \mathbb{N}^n .

In 1942, C.L. Siegel proved [15], assuming non-resonant hypothesis and eigenvalues belonging to the Siegel domain, that system (1.1) can be analytically linearized provided $(\omega_j)_{1 \leq j \leq n}$ are not "well approximable" by integer component vectors:

$$\exists \gamma > 0 \text{ and } \tau > n - 1 : \forall \alpha \in \mathbb{N}^n \text{ and } \forall j \in \{1, \dots, n\} : |\alpha \cdot \omega - \omega_j| \geq \gamma |\alpha|^{-\tau},$$

in this case we say that $A = \text{diag}(\omega_1, \dots, \omega_n)$ verifies a *Diophantine condition* of exponent τ and constant γ , for short $A \in CD(\gamma, \tau)$.

A.D. Bruno [2], weakened the arithmetical condition of Siegel by assuming the, today called, *Bruno condition*:³

$$\sum_{k \geq 0} \frac{\log \Omega^{-1}(2^{k+1})}{2^k} < +\infty, \tag{1.4}$$

where for all positive integer p :

$$\Omega(p) = \min\{|\alpha \cdot \omega - \omega_j| : j \in \{1, \dots, n\}, \alpha \in \mathbb{N}^n, 0 < |\alpha| < p\}.$$

Thus if A is non-resonant, semisimple and verifies the above condition, then there exists an analytic linearization which brings (1.1) into (1.2). In [2] also the resonant case has been considered, and additional conditions must be imposed to ensure existence of an analytical normal form.

So let us assume A to verify hypothesis of Theorem 1.1: let $(\omega_j)_{1 \leq j \leq n} \subset \mathbb{R}$, with at least two of them have opposite signs, and $A = \text{diag}(i\omega_1, \dots, i\omega_n)$, namely A belongs to the *Siegel domain* ⁴

Let us make a step backward and consider the following problem [4]. Let $\mathcal{A}_1 \subset \mathcal{A}_2 \subset \mathbb{C}^n[[z_1, \dots, z_n]]$ be two classes of formal power series closed w.r.t. to derivation and composition. Let $\hat{f} \in \mathcal{A}_1$ s.t. $\hat{f} = \sum_{|\alpha| \geq 2} f_\alpha z^\alpha$, let $A \in GL(n, \mathbb{C})$ and consider the following (formal) ODE:

$$\frac{dz}{dt} = Az + \hat{f}(z). \tag{1.5}$$

We say that (1.5) is *linearizable in \mathcal{A}_2* if there exists $\hat{H} \in \mathcal{A}_2$, normalized with $\hat{H} = z + \mathcal{O}(|z|^{|\alpha|})$, $|\alpha| \geq 2$, s.t. *formally* we have:

$$w = \hat{H}(z) \quad \text{and} \quad \frac{dw}{dt} = Aw.$$

If both \mathcal{A}_1 and \mathcal{A}_2 coincide with the ring of formal power series we already know that, generically, the problem has solution if and only if A is *non-resonant*,

³ The Bruno condition can be rewritten using a general increasing sequence of integer numbers, $(p_k)_{k \geq 0}$, infact Bruno proved [2] at pag. 222, that (1.4) is equivalent to:

$$\sum_{k \geq 0} \frac{\log \Omega^{-1}(p_{k+1})}{p_k} < +\infty.$$

⁴ According to the classification of [2] this case is *Poincaré domain 1.d*, but we prefer consider it as a Siegel case because the obstructions to the linearizability are very similar to those encountered in the Siegel domain.

which will be assumed from now. In the other cases of general algebras of formal power series, new arithmetical conditions on A have to be imposed if we are in the Siegel domain. Whereas for the Poincaré domain the classical Poincaré proof can be adapted to handle this situation. The former case has been considered in details in [4] section 5, to which we refer for all details⁵. There author proved that the Bruno condition is still sufficient to linearize whenever $\mathcal{A}_1 = \mathcal{A}_2$, otherwise new Bruno-like conditions are introduced, weaker than the original Bruno condition. This point of view has been introduced in [4] to understand the intermediate cases "between" the analytical and the formal one, and to determine an arithmetical condition which "interpolates" between the two extremal condition: Bruno's condition and non-resonance condition.

An interesting case is when \mathcal{A}_1 is the ring of convergent power series in some neighbourhood of the origin, and \mathcal{A}_2 is the algebra of *Gevrey- s* , $s > 0$, formal power series. Namely we are considering the *Gevrey linearization of analytic vector fields*.

Let $\hat{F} = \sum f_\alpha z^\alpha$, $(f_\alpha)_{\alpha \in \mathbb{N}^n} \subset \mathbb{C}^n$ be a formal power series, then we say that it is *Gevrey- s* [1, 14], $s > 0$, if there exist two positive constants C_1, C_2 such that:

$$|f_\alpha| \leq C_1 C_2^{-s|\alpha|} (|\alpha|!)^s \quad \forall \alpha \in \mathbb{N}^n. \quad (1.6)$$

We denote the class of formal vector valued power series *Gevrey- s* by \mathcal{G}_s . It is closed w.r.t. derivation and composition.

In the *Gevrey- s* case the arithmetical condition introduced in [4], called *Bruno- s condition*, $s > 0$, for short \mathcal{B}_s , reads:

$$\limsup_{|\alpha| \rightarrow +\infty} \left(2 \sum_{m=0}^{\kappa(\alpha)} \frac{\log \Omega^{-1}(p_{m+1})}{p_m} - s \log |\alpha| \right) < +\infty, \quad (1.7)$$

for some increasing sequence of positive integer $(p_k)_{k \geq 0}$ and $\kappa(\alpha)$ is defined by $p_{\kappa(\alpha)} \leq |\alpha| < p_{\kappa(\alpha)+1}$.

Remark 1.2. This definition recall the classical one of Bruno [2], where first one suppose the existence of a strictly increasing sequence of positive integer such that (1.7) holds, then one can prove (see [2] §IV page 222) that one can take an exponentially growing sequence, e.g. $p_k = 2^k$. This holds also in our case, in fact we can prove that (1.7) is equivalent to:

$$\limsup_{N \rightarrow +\infty} \left(\sum_{l=0}^N \frac{\log \Omega^{-1}(2^{l+1})}{2^l} - sN2 \log 2 \right) < +\infty.$$

A proof of this claim can be found in [5].

When $n = 2$, under the above condition (non-resonance and Siegel domain), rescaling time by $-\omega_2$ (we recall that the non-resonance condition implies $\omega_2 \neq 0$),

⁵ See also [3] where a similar problem for germs of diffeomorphisms of $(\mathbb{C}, 0)$ has been studied.

the ODE associated to the vector field can be rewritten as:

$$\begin{cases} \dot{z}_1 = \omega z_1 + h.o.t. \\ \dot{z}_2 = -z_2 + h.o.t. \end{cases}, \tag{1.8}$$

where $\omega = -\omega_1/\omega_2 \in (\mathbb{R} \setminus \mathbb{Q})^+$ and high order terms means $\mathcal{O}(|z|^{|\alpha|})$ with $|\alpha| \geq 2$, namely only the ratio of the eigenvalues enters. Then the Bruno- s condition can be slightly weakened (see [3]):

$$\limsup_{n \rightarrow +\infty} \left(\sum_{j=0}^{k(n)} \frac{\log q_{j+1}}{q_j} - s \log n \right) < +\infty, \tag{1.9}$$

where $k(n)$ is defined by $q_{k(n)} \leq n < q_{k(n)+1}$ and $(q_n)_n$ are the denominators of the convergents [10] to ω .

We remark that in both cases new conditions (1.7) and (1.9) are weaker than Bruno's condition, which is recovered when $s = 0$. When $n = 2$ we prove that the set $\bigcup_s \mathcal{B}_s$ is $PSL(2, \mathbb{Z})$ -invariant (see remark 3.1).

The main result of [4] in the case of Gevrey- s classes reads:

Theorem 1.3. (Gevrey- s linearization) *Let $\omega_1, \dots, \omega_n$ be real numbers and $A = \text{diag}(i\omega_1, \dots, i\omega_n)$; let $D_1 = \{z \in \mathbb{C}^n : |z_i| < 1, 1 \leq i \leq n\}$ be the isotropic polydisk of radius 1 and let $F : D_1 \rightarrow \mathbb{C}^n$ be an analytic function, such that $F(z) = Az + f(z)$, with $f(0) = Df(0) = 0$. If A is non-resonant and verifies a Bruno- s , $s > 0$, condition (1.7) (or condition (1.9) if $n = 2$), then there exists a formal Gevrey- s linearization \hat{H} .*

The aim of this paper is to show that the Gevrey character of the formal linearization can give information concerning the dynamics of the analytic vector field. Let $F(z) = Az + f(z)$ verify hypotheses of Theorem 1.3, assume moreover X_F not to be analytically linearizable. We will show that even if there is not a *Stable domain*, where the dynamics of X_F is conjugate to the dynamics of its linear part, we have an open neighbourhood of the origin which "behaves as a Stable domain" for the flow of X_F for finite but long time, which results exponentially long: the *effective stability* [8, 9] of the equilibrium solution.

In the case of analytic linearization, i.e. \hat{H} is convergent in a neighbourhood of the origin, let $z \mapsto H(z)$ be the germ of analytic diffeomorphism (change of coordinates $w = H(z)$) whose Taylor series at the origin coincides with \hat{H} . Then $w_j(t) = H_j(z(t))$ for all $j = 1, \dots, n$, thus by (1.2) $w_j(t) = e^{it\omega_j} w_j(0)$ and so $|H_j(z(t))| = |H_j(z(0))|$, i.e. $(|H_j(z)|)_{1 \leq j \leq n}$, is *constant along the orbits*, namely it is a *first integral* and the flow of (1.1) is bounded for all t and sufficiently small $|z_0|$.

We will prove that any non-zero z_0 belonging to a polydisk of sufficiently small radius $r > 0$, can be followed up to a time $T = \mathcal{O}(\exp\{r^{-1/s}\})$, being $s > 0$ the Gevrey exponent of the formal linearization, and we can find an *almost first integral*: a function which varies by a quantity of order r during this interval of

time. More precisely we prove the following

Theorem 1.4. *Let $n \in \mathbb{N}$, $n \geq 2$. Given real $\omega_1, \dots, \omega_n$, with at least two of them of opposite sign, consider $A = \text{diag}(i\omega_1, \dots, i\omega_n)$; let $F : D_1 \rightarrow \mathbb{C}^n$ be an analytic function, such that $F(z) = Az + f(z)$, with $f(0) = Df(0) = 0$. If A is non-resonant and verifies a Bruno- s , $s > 0$, condition (1.7) (or (1.9) if $n = 2$), then for all sufficiently small $0 < r_{**} < 1$, there exist positive constants A_{**}, B_{**}, C_{**} such that for all $0 < |z_0| < r_{**}/2$, the solutions $z(t; z_0)$ are well defined and verify $|z(t; z_0)| \leq C_{**}r_{**}$, for all $|t| \leq T_* = A_{**}^{-1} \exp\{B_{**}(r_{**}/|z_0|)^{1/s}\}$.*

We want to stress here that, when $s \rightarrow 0$ the stability time goes to infinity, because the *critical exponent of stability time* is $1/s$: namely we obtain *stability*. At the same time the Bruno- s condition “tends” to the classical Bruno condition, which is a sufficient condition to ensure analytic linearizability and hence stability under our assumptions. Thus we recover the classical stability result as limit of longer and longer effective stability times.

Results similar to Theorem 1.4 have been obtained in [8] for hamiltonian vector fields and in [9] for real reversible systems of coupled harmonic oscillators. In both papers effective stability is proved by assuming the linear part of the vector field to verify some Diophantine condition $CD(\gamma, \tau)$, for some $\gamma > 0$ and $\tau > n - 1$, and the critical exponent of stability time is $1/\tau$. In our result, too, the critical exponent of stability time depends on some arithmetical property of the linear part of the vector field but in a more general way in fact we assume $A \in \mathcal{B}_s \supset CD(\gamma, \tau)$, for all $\gamma > 0$ and $\tau \geq n - 1$.

The main difference between the present result and the one of [8, 9] is that here we first *linearize formally* the system and then using properties of the formal linearization we conclude using the Poincaré summation at the smallest term (see Lemma 2.2), whereas in [8, 9] authors obtain effective stability using a finite order normal form, then working on it and using again the Poincaré summation at the smallest term, they conclude their proof.

Hence the main difference is that we have to conjugate the analytic vector field with a linear one, whereas a non-linear normal form is used in [8, 9]. This introduces our main drawback: we must assume A to be non-resonant (to be formally linearizable) and this prevents us from considering real vector fields and hamiltonian ones, where an “intrinsic” resonance is present.

In section 3 we discuss the relation between the Bruno- s condition and other arithmetical conditions.

2. Proof of the main Theorem

In this part we will prove our main result, Theorem 1.4. The proof will be divided into three steps: first we use the Gevrey- s character of the formal linearization \hat{H} , given by Theorem 1.3, to find an approximate solution of the conjugacy equa-

tion (1.3) up to a (exponentially) small correction (paragraph 2.1); then we prove a Lemma allowing us to control how the small error introduced in the solution propagates (paragraph 2.2). Finally we collect all the informations to conclude the proof (paragraph 2.3).

2.1. Determination of an approximate solution

Let F verifies hypotheses of Theorem 1.4 and let us consider the first order differential equation in \mathbb{C}^n , $n \geq 2$:

$$\frac{dz}{dt} = F(z). \tag{2.1}$$

By Theorem 1.3 this system can be put in linear form by a formal power series \hat{H} which belongs to \mathcal{G}_s and it solves (formally):

$$\frac{d}{dt}\hat{H}(z) = A\hat{H}(z), \tag{2.2}$$

we observe that one can choose $\hat{H}(z) = z + \mathcal{O}(|z|^2)$.

Since $\hat{H} = \sum h_\alpha z^\alpha \in \mathcal{G}_s$, there exist positive constants A_1 and B_1 such that

$$|h_\alpha| \leq A_1 B_1^{-s|\alpha|} (|\alpha|!)^s \quad \forall |\alpha| \geq 1. \tag{2.3}$$

For any positive integer N we consider the *vectorial polynomial*, sum of homogeneous vector monomials of degree $1 \leq l \leq N$, defined by: $\mathcal{H}_N(z) = \sum_{l=1}^N \sum_{|\alpha|=l} h_\alpha z^\alpha$ and the *Remainder Function*:

$$\mathcal{R}_N(z) = D\mathcal{H}_N(z) \cdot F(z) - A\mathcal{H}_N(z). \tag{2.4}$$

Clearly $\mathcal{H}_N(z)$ doesn't solve the linearization problem, but:

$$\frac{d}{dt}\mathcal{H}_N(z) = A\mathcal{H}_N(z) + \mathcal{R}_N(z), \tag{2.5}$$

hence the remainder function gives the difference from the true solution and the approximate one.

The following Proposition collects some properties of the remainder function.

Proposition 2.1. *Let $\mathcal{R}_N(z)$ be the remainder function defined in (2.4) and let $\alpha \in \mathbb{N}^n$, then:*

- 1) $\partial_z^\alpha \mathcal{R}_N(0) = 0$ if $|\alpha| \leq N$.
- 2) For all $0 < r < 1$ there exist positive constants A_2 and B_2 such that if $|\alpha| \geq N + 1$, then:

$$\left| \frac{1}{\alpha!} \partial_z^\alpha \mathcal{R}_N(0) \right| \leq A_2 r^{-|\alpha|} B_2^{-sN} (N!)^s.$$

3) For all $0 < r < 1$ and $|z| < r/4$ there exist positive constants A_3, B_3 such that:

$$|\mathcal{R}_N(z)| \leq A_3 B_3^{-sN} (N!)^s \left(\frac{|z|}{r} \right)^{N+1}. \quad (2.6)$$

Where we used the compact notation $\frac{1}{\alpha!} \partial_z^\alpha = \frac{1}{\alpha_1! \dots \alpha_n!} \frac{\partial^{|\alpha|}}{\partial z_1^{\alpha_1} \dots \partial z_n^{\alpha_n}}$.

Proof. Statement 1) is an immediate consequence of the definition of \mathcal{R}_N .

To prove 2) we observe that $\mathcal{R}_N(z)$ is an analytic function on D_1 , being obtained with product of analytic functions, then one gets by Cauchy's estimates for all $0 < r < 1$ and for all $|\alpha| \geq N + 1$:

$$\left| \frac{1}{\alpha!} \partial_z^\alpha \mathcal{R}_N(0) \right| \leq \frac{1}{(2\pi)^n} \frac{1}{r^{|\alpha|+1}} \max_{|z|=r} |D\mathcal{H}_N \cdot F(z)|, \quad (2.7)$$

because $\partial^\alpha \mathcal{H}_N = 0$ for $|\alpha| \geq N + 1$. Recalling the Gevrey estimate (2.3) for \mathcal{H}_N and the analyticity of F we obtain:

$$\left| \frac{1}{\alpha!} \partial_z^\alpha \mathcal{R}_N(0) \right| \leq A_2 B_2^{-sN} (N!)^s r^{-|\alpha|}, \quad (2.8)$$

for some positive constants A_2 and B_2 depending on the previous constants, on the dimension n and on F .

To prove 3) let us write the Taylor series $\mathcal{R}_N(z) = \sum_{|\alpha| \geq N+1} \frac{1}{\alpha!} \partial_z^\alpha \mathcal{R}_N(0) z^\alpha$: the bound on derivatives (2.8) implies the estimate (2.6) for all $|z| < r/4$ and for some positive constants A_3 and B_3 . \square

The bound (2.6) on $\mathcal{R}_N(z)$ depends on the positive integer N , so we can determine the value of N for which the right hand side of (2.6) attains its minimum, that's Poincaré's idea of *summation at the smallest term*.

Lemma 2.2. (Summation at the smallest term)

Let $\mathcal{R}_N(z)$ defined as before and let $0 < r_* < 1/4$ then there exist positive constants A_4, B_4 such that for all $0 < |z| < r_*$ we have:

$$|\mathcal{R}_{\bar{N}}(z)| \leq A_4 \exp \left\{ -B_4 \left(\frac{r_*}{|z|} \right)^{1/s} \right\}, \quad (2.9)$$

where $\bar{N} = \lfloor B_4 (r_*/|z|)^{1/s} \rfloor$ and $\lfloor x \rfloor$ denotes the integer part of $x \in \mathbb{R}$.

Proof. Let us fix $0 < r_* < 1/4$, then for $0 < |z| < r_*$ by Stirling formula we obtain:

$$|\mathcal{R}_N(z)| \leq A_4 \left(N B_3^{-1} (|z|/r_*)^{1/s} \right)^{Ns} e^{-sN}, \quad (2.10)$$

for some positive constant A_4 . The right hand side of (2.10) attains its minimum at $\bar{N} = B_3 (r_*/|z|)^{1/s}$, evaluating the value of this minimum we get (2.9) with $B_4 = B_3$. \square

2.2. Control of the “errors”

Let us define $\mathcal{H}(z) = \mathcal{H}_{\bar{N}}(z)$ and $\mathcal{R}(z) = \mathcal{R}_{\bar{N}}(z)$, being \bar{N} the “optimal value” obtained in Lemma 2.2. We remark that $\mathcal{H}(z)$ doesn’t solve (2.2) but the “error”, $\mathcal{R}(z)$, can be made very small: exponentially small. We will prove that for initial conditions in a sufficiently small disk, one can follow the flows for an exponential long time without leaving a disk of comparable size.

Lemma 2.3. (Control the flow) *Let a, b, α and R be positive real numbers. Let $T = Ra^{-1}e^{b/(2R)^\alpha}$ and let us consider the Cauchy problem:*

$$\begin{cases} \frac{d}{dt}x(t) = ae^{-b/x^\alpha} \\ x(0) = R \end{cases} .$$

Then $0 < x(t) < 2R$ for all $|t| < T$.

Proof. Let us write the Cauchy problem in integral form:

$$x(t) = R + \int_0^t ae^{-b/(x(s))^\alpha} ds ,$$

$x(t)$ is trivially monotonically increasing, hence the same holds for the function $t \mapsto ae^{-b/(x(t))^\alpha}$. Let us suppose that there exists $0 < t_0 < T$, for which $x(t_0) = 2R$; then:

$$2R = x(t_0) = R + \int_0^{t_0} ae^{-b/(x(s))^\alpha} ds < R + t_0ae^{-b/(x(t_0))^\alpha} ,$$

namely $t_0 > Ra^{-1}e^{b/(2R)^\alpha} = T$, which gives a contradiction. Hence either $x(t_0) > 2R$ for all $0 < t < T$ or $x(t_0) < 2R$, but the first case have to be excluded because $x(0) = R < 2R$.

The case $t < 0$ can be handled in a similar way by showing that $t \mapsto x(t)$ doesn’t decrease too much. □

Let r_* as in Lemma 2.2, define $\rho(z) = |\mathcal{H}(z)|$ for all $0 < |z| < r_*$, then Lemma 2.2 admits the following Corollary, which allows us to control the evolution of $\rho(z)$.

Corollary 2.4. *Let $0 < r_* < 1/4$, then there exists $0 < r_{**} \leq r_*$ and positive constants A_*, B_* such that for all $0 < |z| < r_{**}$ we have:*

$$\frac{d}{dt}\rho(z) \leq A_* \exp\left\{ -B_* \left(\frac{r_*}{\rho(z)}\right)^{1/s} \right\} . \tag{2.11}$$

Proof. Let $j \in \{1, \dots, n\}$ and let us consider the time evolution of $|\mathcal{H}_j(z(t))|$. If \mathcal{H} was a solution this would be a constant of motion, this is not the case but its evolution is nevertheless very slow. In fact thanks to (2.5) we get:

$$\frac{d}{dt}|\mathcal{H}_j(z)| \leq |\mathcal{R}_j(z)| ,$$

hence a similar statement holds for $\rho(z) = |\mathcal{H}(z)| = \max_{1 \leq j \leq n} |\mathcal{H}_j(z)|$.

We want now to express the exponential smallness of $|\mathcal{R}(z)|$ in terms of $\rho(z)$ instead of $|z|$. $\mathcal{H}(z)$ is tangent to the identity close to zero and then locally invertible. The inverse is still tangent to the identity, vanishing at zero and analytic in a neighbourhood of the origin, then sufficiently close to the origin we have $|\mathcal{H}^{-1}(w)| \leq C|w|$, for some $C > 0$. Finally we can take $|z|$ sufficiently small, say $|z| < r_{**}$ for some $0 < r_{**} \leq r_*$, s.t. $|z| \leq C|\mathcal{H}(z)| < r_*$, hence:

$$|\mathcal{R}(z)| \leq A_4 \exp\left\{-B_4 \left(\frac{r_*}{|z|}\right)^{1/s}\right\} \leq A_4 \exp\left\{-B_4 C^{-1/s} \left(\frac{r_*}{\rho(z)}\right)^{1/s}\right\},$$

and the claim follows with $A_* = A_4$ and $B_* = B_4 C^{-1/s}$. \square

2.3. End of the proof

We are now able to conclude the proof of the main Theorem 1.4. Take any $0 < |z_0| < r_{**}/2$ and let $\rho_0 = |\mathcal{H}(z(0; z_0))|$, then there exists a positive constant C_1 s.t. $\rho_0 \leq C_1|z_0|$. By Corollary 2.4 we have

$$\frac{d}{dt}\rho(z(t; z_0)) \leq A_* \exp\left\{-B_* (r_*/\rho(z(t; z_0)))^{1/s}\right\}. \quad (2.12)$$

Let us call $R = \rho_0$, $a = A_*$, $b = B_* r_*^{1/s}$ and $\alpha = 1/s$ then we can apply Lemma 2.3, to conclude:

$$\rho(z(t; z_0)) \leq 2\rho_0 < C_1 r_{**} \quad \forall |t| \leq T_* = \rho_0 A_*^{-1} \exp\left\{B_* \left(\frac{r_*}{2\rho_0}\right)^{1/s}\right\}. \quad (2.13)$$

This implies that $\mathcal{H}(z(t; z_0))$ is well defined in this interval of time, it is not constant and $|\mathcal{H}(z(t; z_0))| \leq 3C_1 r_{**}$. Recalling that $\mathcal{H}(z)$ is tangent to the identity close to zero, we have $|z| \leq C_3 |\mathcal{H}(z)|$ for some positive C_3 .

Then setting $A_{**} = 2A_* r_{**}^{-1}$, $B_{**} = B_* (r_*/(2r_{**}))^{1/s}$ and $C_{**} = C_1 C_3$, we get:

$$|z(t; z_0)| \leq C_{**} r_{**},$$

for all $|t| \leq A_{**}^{-1} \exp\left\{B_{**} \left(\frac{r_{**}}{|z_0|}\right)^{1/s}\right\}$.

3. Arithmetical conditions

In this paper we proved that any analytic germs of vector field of $(\mathbb{C}^n, 0)$ with diagonal, non-resonant linear part of Siegel type has an *effective stability* domain, i.e. stable up to finite but “long times”, close to the stationary point, provided the linear part verifies an arithmetical Bruno-like condition depending on a parameter

$s > 0$, which in the case of 2 dimensional vector fields can be put in the form:

$$\limsup_{n \rightarrow +\infty} \left(\sum_{j=0}^{k(n)} \frac{\log q_{j+1}}{q_j} - s \log n \right) < +\infty.$$

The aim of this section is to compare our Bruno- s conditions with other arithmetical conditions used in literature.

Let $\gamma > 0, \tau > 1$ then for all $x \in CD(\gamma, \tau)$, one has:

$$\sum_{j=0}^{+\infty} \frac{\log q_{j+1}}{q_j} \leq C_1 + C_2(\tau + 1),$$

for some positive constants C_1 and C_2 thus, $CD(\gamma, \tau) \subset \mathcal{B} \subset \mathcal{B}_s$, for all $s > 0$.

The group $PSL(2, \mathbb{Z})$ play an important role in number theory being the generator of the (Gauss) continued fraction algorithm, hence it is important to control its action on \mathcal{B}_s .

Remark 3.1. (Invariance of $\bigcup_{s>0} \mathcal{B}_s, n = 1$ under the action of $PSL(2, \mathbb{Z})$) The continued fraction development [10, 11] of an irrational number ω gives us the sequences: $(a_k)_{k \geq 0}$ and $(\omega_k)_{k \geq 0}$. Then we introduce $(\beta_k)_{k \geq -1}$ defined by $\beta_{-1} = 1$ and for all integer $k \geq 0$: $\beta_k = \prod_{j=0}^k \omega_j$, which verifies: $1/2 < \beta_k q_{k+1} < 1$ and $q_n \beta_{n-1} + q_{n-1} \beta_n = 1$, where q_k 's are the denominators of the continued fraction development of ω . We claim that condition Bruno- s (1.9) is equivalent to the following one:

$$\limsup_{k \rightarrow +\infty} \left(\sum_{j=0}^k \beta_{j-1} \log \omega_j^{-1} + s \log \beta_{k-1} \right) < +\infty. \tag{3.1}$$

This can be proved by using the relations between β_l and q_l , to obtain the bound, for all integer $k > 0$:

$$\begin{aligned} & \left| \sum_{l=0}^k \left(\beta_{l-1} \log \omega_l + \frac{\log q_{l+1}}{q_l} \right) \right| \\ & \leq \sum_{l=0}^k \left| \beta_{l-1} \log \beta_l q_{l+1} \right| + \left| \beta_{l-1} \log \beta_{l-1} \right| + \left| \frac{q_{l-1}}{q_l} \beta_l \log q_{l+1} \right| \leq 18, \end{aligned}$$

where we used the convergence of series $\sum q_l^{-1}$ and $\sum q_l^{-1} \log q_l$ (see [11] page 272).

To prove the invariance of $\bigcup_s \mathcal{B}_s$ under the action of $PSL(2, \mathbb{Z})$, is enough to consider its generators: $T\omega = \omega + 1$ and $S\omega = 1/\omega$. For any irrational ω , T acts trivially being $\beta_k(T\omega) = \beta_k(\omega)$ for all k , whereas for S we have $\beta_k(\omega) = \omega \beta_{k-1}(S\omega)$ for all $k \geq 1$. Let ω be an irrational and let $\omega' = \omega^{-1}$, let us also denote with a $'$ quantities given by the continued fraction algorithm applied to ω' , then using (3.1)

one can prove:

$$\omega_0 \left(\sum_{j=0}^k \beta'_{j-1} \log \omega'_j{}^{-1} + s \omega_0^{-1} \log \beta'_{k-1} \right) = C(\omega, s) + \sum_{j=0}^{k+1} \beta_{j-1} \log \omega_j^{-1} + s \log \beta_k,$$

where $C(\omega, s) = \omega_0 (\log \omega_1^{-1} - s \log \omega_0) + \sum_{l=0}^1 \beta_{l-1} \log \omega_l^{-1}$, from which the claim follows.

Let us consider a slightly stronger version of the Bruno- s condition: $\omega \in (0, 1) \setminus \mathbb{Q}$ belongs to $\tilde{\mathcal{B}}_s$ if:

$$\lim_{n \rightarrow +\infty} \left(\sum_{l=0}^k \frac{\log q_{l+1}}{q_l} - s \log q_k \right) < +\infty, \quad (3.2)$$

where $(q_n)_n$ are the convergents to ω .

Remark 3.2. This new condition is stronger than Bruno- s , because the existence of the limit is required. One can construct numbers ω which verify \mathcal{B}_s but not $\tilde{\mathcal{B}}_s$. For a simple proof we refer the interested reader to [5].

Finally let us introduce a second arithmetical condition denoted by \mathcal{B}'_s to be the set of irrational numbers whose convergents verify:

$$\lim_{k \rightarrow +\infty} \frac{\log q_{k+1}}{q_k \log q_k} = s. \quad (3.3)$$

We state without proof the following proposition, and we refer to [5], to all details:

Proposition 3.3. *Let $s > 0$ and let $\omega \in (0, 1) \cap \tilde{\mathcal{B}}_s$. Then if ω is not a Bruno number then $\omega \in \mathcal{B}'_s$, otherwise $\omega \in \mathcal{B}'_0$.*

Therefore if $\omega \in \tilde{\mathcal{B}}_s \setminus \mathcal{B}$ then the denominators of the convergent to ω can grow as a factorial, more precisely, $q_{k+1} = \mathcal{O}((q_k!)^s)$, is allowed.

References

- [1] W. Balsler, *From Divergent Power Series to Analytic Functions. Theory and Applications of Multisummable Power Series*, Lectures Notes in Mathematics, **1582**, Springer, (1994).
- [2] A.D. Bruno, Analytical form of differential equations, *Transactions Moscow Math. Soc.* **25** (1971), 131–288.
- [3] T. Carletti and S. Marmi, Linearization of analytic and non-analytic germs of diffeomorphisms of $(\mathbb{C}, 0)$, *Bull. Soc. Math. Française* **128** (2000), 69–85.
- [4] T. Carletti, *The Lagrange inversion formula on non-Archimedean fields. Non-analytical form of differential and finite difference equations*, DCDS series A, Vol. **9** N. 4 (2003), 835–858.
- [5] T. Carletti, Exponentially long time stability for non-linearizable analytic germs of $(\mathbb{C}^n, 0)$, *Ann. Inst. Fourier*, **54** (2004), 1–15.

- [6] T. Carletti, A. Margheri e M. Villarini, Normalization of Poincar Singularities via Variation of Constants, preprint submitted to Publicacions Matemàtiques (2004).
- [7] H. Dulac, Solutions d'un systeme d'equations differentielles dans le voisinage de valeurs singulieres, *Bull. Soc. Math. Fr.* **40** (1912), 324–383.
- [8] A. Giorgilli, A. Fontich, L. Galgani and C. Simó, Effective stability for a Hamiltonian system near an elliptic equilibrium point, with an application to the restricted three body problem, *J. of Differential Equations* **77** (1989), 167–198.
- [9] A. Giorgilli and A. Posilicano, Estimates for normal forms of differential equations near an equilibrium point, *J. of Appl. Math. and Phys. (ZAMP)*, **39** (1989), 713–732.
- [10] G.H.Hardy and E.M.Wright, *An introduction to the theory of numbers*, 5th edition Oxford Univ. Press, 1979.
- [11] S.Marmi, P.Moussa and J.-C.Yoccoz, The Brjuno functions and their regularity properties, *Communications in Mathematical Physics* **186** (1997), 265–293.
- [12] J. Moser, Stable and random motions in dynamical systems, *Annals of Math. Stud.* **77**, Princeton Univ. Press, (1977).
- [13] H. Poincaré, *Œuvres, tome I*, Gauthier–Villars, Paris 1917.
- [14] J.-P. Ramis, Séries divergentes et Théorie asymptotiques, *Publ. Journées X-UPS* (1991), 1–67.
- [15] C.L. Siegel, Iteration of analytic functions, *Annals of Mathematics* **43** (1942), 807–812.
- [16] S. Sternberg, The structure of local homeomorphisms II, III, *Amer. J. Math.* **80** (1980), 623–632 and **81** 578–604.

Timoteo Carletti
Scuola Normale Superiore
piazza dei Cavalieri 7
56126 Pisa
Italia
e-mail: t.carletti@sns.it

(Received: July 5, 2004; revised: December 9, 2004)



To access this journal online:
<http://www.birkhauser.ch>
