

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

Vaccination of calves using the BRSV nucleocapsid protein in a DNA prime-protein boost strategy stimulates cell-mediated immunity and protects the lungs against BRSV replication and pathology

Letellier, Carine; Boxus, Mathieu; Rosar, Laurent; Toussaint, Jean-François; Walravens, Karl; Roels, Stefan; Meyer, Gilles; Letesson, Jean-Jacques; Kerkhofs, Pierre

Published in: Vaccine

Publication date: 2008

Document Version Early version, also known as pre-print

Link to publication

Citation for pulished version (HARVARD):

Letellier, C, Boxus, M, Rosar, L, Toussaint, J-F, Walravens, K, Roels, S, Meyer, G, Letesson, J-J & Kerkhofs, P 2008, 'Vaccination of calves using the BRSV nucleocapsid protein in a DNA prime-protein boost strategy stimulates cell-mediated immunity and protects the lungs against BRSV replication and pathology', Vaccine, vol. 26, no. 37, pp. 4840-4848.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
You may not further distribute the material or use it for any profit-making activity or commercial gain
You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Vaccine 26 (2008) 4840-4848



Contents lists available at ScienceDirect

Vaccine



journal homepage: www.elsevier.com/locate/vaccine

Vaccination of calves using the BRSV nucleocapsid protein in a DNA prime-protein boost strategy stimulates cell-mediated immunity and protects the lungs against BRSV replication and pathology

Letellier Carine^{a,*}, Boxus Mathieu^b, Rosar Laurent^a, Toussaint Jean-François^a, Walravens Karl^a, Roels Stefan^a, Meyer Gilles^c, Letesson Jean-Jacques^d, Kerkhofs Pierre^a

^a Veterinary and Agrochemical Research Centre, Brussels, Belgium

^b FUSAGx, Molecular and Cellular Biology, Gembloux, Belgium

^c Ecole Nationale Vétérinaire, UMR INRA/ENVT 1225 Interactions Hôtes-Agents Pathogènes, Toulouse, France

^d Facultés Universitaires Notre-Dame de la Paix, Unité de Recherche en Biologie Moléculaire, Namur, Belgium

ARTICLE INFO

Article history: Received 23 May 2007 Received in revised form 25 June 2008 Accepted 29 June 2008 Available online 21 July 2008

Keywords: BRSV Prime-boost Nucleocapsid gene

ABSTRACT

Respiratory syncytial virus (RSV) is a major cause of respiratory disease in both cattle and young children. Despite the development of vaccines against bovine (B)RSV, incomplete protection and exacerbation of subsequent RSV disease have occurred. In order to circumvent these problems, calves were vaccinated with the nucleocapsid protein, known to be a major target of CD8⁺ T cells in cattle. This was performed according to a DNA prime–protein boost strategy. The results showed that DNA vaccination primed a specific T-cell-mediated response, as indicated by both a lymphoproliferative response and IFN- γ production. These responses were enhanced after protein boost. After challenge, mock-vaccinated by successive administrations of plasmid DNA and protein exhibited protection against the development of pneumonic lesions and the viral replication in the BAL fluids and the lungs. The protection correlated to the cell-mediated immunity and not to the antibody response.

© 2008 Elsevier Ltd. All rights reserved.

1. Introduction

RSV is a major cause of hospitalization of infants less than 2 years of age [1]. Like its human counterpart, bovine respiratory syncytial virus (BRSV) is a major cause of calf morbidity and mortality [2]. Calves less than 1 year of age are particularly susceptible to the disease [3]. Maternal antibodies against respiratory syncytial virus do not prevent infection in calves but antibodies seem to decrease the severity of infection [4–6].

Furthermore, BRSV is one of the viruses contributing to bovine respiratory disease complex [7]. Protection against BRSV afforded by vaccination has been described and numerous vaccines are licensed and widely used in the field [8–10]. However, exacerbation of respiratory disease has been induced in calves and in infants vaccinated with formalin-inactivated BRSV vaccine [11,12]. In mice, enhanced disease has an immunopathological basis, characterized by polarized type 2-helper (TH2) response. Increased IL-4 production has been associated with pulmonary histopathology

and eosinophilia [13–15]. In calves, enhanced pathology has been associated with elevated IgE titers and marked eosinophilia, suggesting a TH2-mediated immune response [16–18]. In addition, some authors have reported high mortality in calves subsequent to the use of inactivated vaccine in the field [19,20]. There is thus a need for protective vaccines, able to confer protection in neonates, even in the presence of maternal antibodies and lacking disease exacerbating side effects.

It has previously been shown that the F and G surface glycoproteins, and the N protein, are the major protective antigens of BRSV [21]. Neutralizing antibodies to the F protein can mediate protection [22]. However, serum antibodies provide greater contribution in reducing BRSV induced lower respiratory disease than upper respiratory tract virus shedding [4,23]. The cell-mediated immunity is also a critical parameter in the outcome of RSV infection [24,25]. The F, P, G, M, M2 and N proteins are antigenic targets recognized by bovine CD8⁺ T cells, the latter two proteins being also specific targets for murine (M2 protein) or murine and human (N protein) CD8⁺ T cells [26–29]. These cells constitute the major lymphocyte subpopulation in the respiratory tract of calves recovering from BRSV infection and depletion of those cells also induces delayed virus clearance from the lungs and nasopharynges in calves [23,30,31].

^{*} Corresponding author. Tel.: +32 2 379 05 64; fax: +32 2 379 06 70. *E-mail address:* calet@var.fgov.be (L. Carine).

⁰²⁶⁴⁻⁴¹⁰X/\$ – see front matter 0 2008 Elsevier Ltd. All rights reserved. doi:10.1016/j.vaccine.2008.06.100

In mice, diminished CTL activation and migration to the lungs by treatment with anti-LFA-1 delayed viral clearance [32]. Furthermore, priming of the RSV-specific CD8⁺ response suppressed the eosinophilia induced after challenge of animals vaccinated with the G glycoprotein [33].

DNA vaccination is an effective way of generating humoral and cell-mediated response [34]. Intramuscular vaccination of mice with plasmid DNA expressing the F or G proteins of RSV has shown to induce a strong TH1 response and to reduce the viral load in the lungs after RSV challenge, without inducing enhanced pulmonary inflammatory response or eosinophilia [35,36]. However, it is generally recognized that DNA vaccines are often less effective in large animals than in mice [37]. Improvement of the immune response can be achieved by boosting animals previously vaccinated with plasmid DNA with inactivated vaccine. This approach has successfully been used to protect calves against bovine herpesvirus-1 (BoHV-1) challenge [38]. This strategy was also successfully used in a previous BRSV experiment. Calves vaccinated with codonoptimized plasmid DNA expressing the F and N BRSV proteins and boosted with an inactivated commercial vaccine were protected against clinical signs, gross pneumonic lesions and virus replication [39]. In that study, the humoral and cell-mediated immune responses were primed by DNA injection and enhanced by the protein boost. They reduced the clinical signs and the virus replication after BRSV challenge. The use of the F protein of RSV in a prime-boost protocol also represents an interesting strategy for early life immunization [40].

The purpose of the work presented here was to investigate the protection of calves against BRSV, by focusing mainly on the



Fig. 1. Polyacrylamide gel electrophoresis of the N protein, purified from recombinant baculovirus infected Sf9 insect cells (lane 2) and a molecular weight marker (Sigma, lane 1). The proteins were stained with Coomassie blue.

stimulation of the BRSV cell-mediated immunity. Therefore, a combination of plasmid and protein vaccination was performed with the N protein as immunogen. The protection conferred by this vaccination scheme was evaluated after BRSV challenge.

2. Materials and methods

2.1. Expression of the N protein

The pcDNA3 plasmid expressing synthetic DNA sequence coding for the ORF of the N protein of BRSV (pNsyn) has been described elsewhere [39]. Large scale plasmid DNA was purified from transformed TOP 10 bacteria cells by affinity chromatography on anion-exchange resin (Plasmid Giga kit, Qiagen).

The purified N protein was obtained from a recombinant baculovirus expressing the coding sequence of the N protein of BRSV strain RB94. Briefly, the N gene was cloned in the pBlueBacHis2A vector (Invitrogen). Recombinant baculovirus expressing the N protein (BacN) was obtained after transfection of Sf9 insect cells with the recombinant plasmid and purification of the β-galactosidaseexpressing plaques as recommended by the manufacturer. Sf9 cells were infected with BacN at a moi of 2. Cells were harvested 48-72 h pi, when at least 95% of the cells were lysed, washed with PBS and resuspended in Tris-HCl 20 mM pH 7.9, NaCl 0.5 M, Imidazole 5 mM buffer. The 6His-tagged N protein was purified on Ni-NTA resin (Qiagen) in batch, as recommended by the manufacturer. The protein concentration was determined by BCA protein assay (Pierce) according to the manufacturer's recommendations. As expected, a protein of 43 kDa was purified and analyzed by PAGE (Fig. 1) [41]. Prior to immunization, QuilA adjuvant was added to the N protein at a final concentration of 1 mg/ml.

2.2. Cell culture and virus

The BRSV RB90 virus was grown on calf primary kidney cells in Minimal Eagle medium (MEM, GIBCO BRL) supplemented with 10% fetal calf serum, 10 mg/ml gentamycin and 10⁶ IU penicillin.

Challenge inoculum's consisted in lung lavage fluid of a calf collected after intratracheal inoculation with the BRSV strain VRS3761, as described previously [39,42]. This inoculum contained 10^{3.4}TCID₅₀/ml and was free of bovine viral diarrhea virus, bovine herpesvirus-1, bovine parainfluenza virus-3, bovine coronavirus, bovine adenovirus-5, endotoxins, bacteria and mycoplasmas. Back titration confirmed the amount of virus inoculated.

2.3. Study design

Fourteen cross-bred dairy calves between 4 and 6 weeks of age were randomly allocated in 4 experimental groups of 3 or 4 animals and housed in 3 identically conditioned stables. All calves were shown to be seronegative for BRSV, BoHV-1 and Bovine viral diarrhea virus at the start of the experiment.

As shown in Table 1, 7 calves were vaccinated twice at a 4 weeks interval (day 0 and 29) with 500 μ g of pNsyn plasmid DNA in saline. The first vaccine dose was injected intramuscularly and the second dose was administrated intradermally in 4 sites in the neck. Seven other calves were subjected to the same protocol, but with the pcDNA3 control plasmid. Four weeks after the second DNA administration (day 60), respectively, 4 and 3 calves injected with the pNsyn and the control plasmids were subjected to intramuscular injection of 300 μ g of purified N protein, adjuvanted with Quil A in a 2 ml volume. The other calves were treated with saline. Three weeks after the third vaccination (day 80), all the calves were challenged with the BRSV 3761 strain by intratracheal (10 ml) injection combined to intranasal nebulization (5 ml). The calves were examined Gro

Mo

pNs Npu

pNs

Table 1	
Experimental de	esign

ир	n	Vaccination			Challenge	n calves
		Day 0/V1	Day 29/V2	Day 60/V3	Day 80	Day 86
:k	4	pcDNA3	pcDNA3	Saline	BRSV	2
yn	3	pNsyn	pNsyn	Saline	BRSV	1
r	3	pcDNA3	pcDNA3	Npur	BRSV	1
yn-Npur	4	pNsyn	pNsyn	Npur	BRSV	2

daily after challenge for the following clinical signs: cough, nasal discharge, anorexia, depression, rectal temperature and respiratory rate.

Sera and heparin treated blood were collected prior to each vaccination, on the day of challenge and on days 6 (day 86) and 12 after challenge (day 92) for examination of humoral and cell-mediated responses. Bronchoalveolar lung (BAL) fluids were collected every 2 or 3 days from day 3 before to day 11 after challenge by instillation and aspiration of 50 ml of PBS as described previously [39]. On days 6 and 12 after challenge, one or two calves of each group were euthanized. Lungs were excised immediately after euthanasia. Macroscopic lesions were recorded to score the extent of pneumonic consolidation and seven pieces of tissue from right cranial (cranial and caudal portions), middle and caudal lobes were excised.

2.4. Lymphocyte proliferation assay (LPA)

Heparinised blood samples were ten-fold diluted in RPMI 1640 medium (Gibco, Belgium) supplemented with 2 mM glutamine, 5×10^{-5} M β -2-Mercaptoethanol, 100 U/ml of gentamycin and 2.5 µg/ml fungizone. Two hundred µl of diluted blood samples were incubated during 6 days with 20 µl of BRSV antigen (Sonicated and heat-inactivated culture supernatant of primary calf kidney cells infected with RB90 virus, containing 10⁵TCID₅₀/ml), or 20 µl of control antigen (Sonicated and heat-inactivated culture supernatant of primary mock-infected calf kidney cells). After the 6-days incubation, 0.8 μ Ci of methyl [³H] thymidine in 25 μ l RPMI 1640 medium was added to each well. Cells were collected after a subsequent 18 h incubation using a cell harvester and radioactivity incorporated into the DNA was measured by liquid scintillation counting (Betaplate, Pharmacia, Sweden). The results were expressed as stimulation indexes (SI) corresponding to the number of c.p.m. obtained with the BRSV antigen divided by the number of c.p.m. obtained with the control preparation. The SI cutoff of 2.4 was determined as the mean+3 standard deviations of the SI values of all animals before vaccination.

2.5. IFN- γ assay

The gamma-interferon (IFN- γ) production was measured after a 24-h in vitro stimulation of 1 ml of heparinised blood with 20 µl of BRSV or control antigen. The amounts of IFN- γ in the plasma were quantified by ELISA with the *Bovine* γ -*interferon EASIA* kit (Biosource Europe). The results were expressed as stimulation indexes (SI) corresponding to the OD obtained with the BRSV antigen divided by the OD obtained with the control preparation. The cut-off SI value of 1.6 was calculated as 3 standard deviations above the mean SI value of all calves before vaccination.

2.6. Serological response

Serum samples were collected from all calves before each vaccination, on the day of challenge and on days 6 and 12 after the challenge. They were subjected to the indirect immunofluorescence test as described previously [43]. Briefly, the sera were serially three-fold diluted starting from a 1/15 dilution and they were incubated on both slides covered with BRSV infected and with non-infected primary kidney cells. After a washing step, the cells were incubated with FITC-conjugated rabbit antibovine IgG and observed on a fluorescence microscope. Slides were examined without any information on the group assignments.

Euthanized Day 92

2

2

2

Local humoral priming was investigated by testing the presence of N-specific IgG and IgA in the BAL after challenge. IgG antibodies were determined in an indirect ELISA, using the purified N protein as coating antigen. After incubation with two-fold dilutions of LBA, starting from 1:10, the test was revealed by using HRP-conjugated rabbit anti-bovine serum and TMB as substrate. N-specific IgA antibodies were determined essentially as described [44]. Briefly, plates were coated with Mab against bovine IgA (Cedi-diagnostics, The Netherlands) and incubated with two-fold dilutions of LBA starting from 1:2. Reaction with the purified N protein was revealed by using biotin-conjugated bovine serum against BRSV, HRP-conjugated streptavidin and TMB.

2.7. Viral replication

The viral load in the BAL fluids and in the lungs was investigated by quantitative real-time RT-PCR. Briefly, for each animal, pieces of approximately 1 cm³ of each lung lobe were mixed together in 10 ml of PBS using an Ultraturrax homogenizer. Two hundred μ l of BAL fluid and 200 μ l of homogenate extracts were subjected to RNA extraction and cDNA synthesis as described previously [45]. Realtime RT-PCR specific for BRSV and for bovine beta-actin (ACTB) were performed along with the corresponding standard curves allowing quantification. The results were expressed as the number of BRSV RNA copies per 10³ ACTB copies in the BAL fluids and per 10⁶ ACTB copies in the lungs.

2.8. Histopathology

A necropsy sample of lesion was taken in each pulmonary lobe. The samples were stored in 10% neutral buffered formalin, embedded in paraffin, sectioned at $5\,\mu m$ and stained with hematoxylin and eosin for histologic examination. Slides were then examined without any information on the group assignments.

2.9. Statistics

The data collected in the 4 groups were compared by one-way analysis of variance and Tukey's test with a family error rate of 0.05. Logarithmic transformation was applied to the lymphocyte proliferation SI, the IFN- γ SI and the IFI titers in order to fulfil the conditions of variances homogeneity (checked by Bartlett's test) and normality (checked by Ryan and Joiner's test).



Fig. 2. In vitro BRSV-specific lymphoproliferation following vaccination and challenge. Blood samples were tested at the time of each vaccination (day 0, 29 and 60), on the day of challenge (week 80) and on days 6 and 12 after challenge (days 86 and 92). The results are expressed as stimulation index. The bars represent the mean value for each group and the vertical bars represent the standard deviation of the means. The horizontal dashed bar represents the cut-off value. The stars indicate the means that differ significantly from the mean of the mock-vaccinated group (p < 0.05).

3. Results

3.1. BRSV-specific cellular response

BRSV-specific lymphoproliferative response was detected in 4 of the 7 calves vaccinated twice with pNsvn plasmid (Fig. 2). While injection of the N protein was not able to induce detectable lymphoproliferative response in non-primed animals, this protein boosted the responses in calves previously vaccinated with pNsyn. After the boost, all the calves of the pNsyn-Npur group had a stimulation index above the cut-off and a considerable enhancement of the BRSV-specific proliferative response. At that time, 2 out of the 3 calves of the pNsyn group still had a SI value above the cut-off. On day 80, the mean SI value of the pNsyn-Npur group was significantly different from the mean SI value of the 3 other groups (p < 0.05) and it remained higher than those of both the mock-vaccinated and Npur groups after challenge (p < 0.05). The BRSV challenge did not induce a detectable response in the mockvaccinated group. In contrast, the lymphoproliferative responses of the calves vaccinated with pNsyn were enhanced and the mean SI value of this group was significantly different from those of the control group and the group vaccinated with the N protein alone (p < 0.05). Twelve days after challenge, the calves of the Npur group also showed low levels of proliferative response.

After 2 plasmid vaccinations, the IFN- γ SI of all animals remained below the cut-off (Fig. 3). However, after the protein boost, the animals subjected to the pNsyn plasmid and the N protein vaccinations displayed significant IFN- γ production compared to the 3 other groups (day 80, p < 0.05). The animals vaccinated with pNsyn alone had also been primed for IFN- γ production, as shown by significant SI values 1 week after the challenge (day 86). On day 86, the IFN- γ SI of the 2 groups vaccinated with pNsyn were significantly different from those of the 2 other groups (p < 0.05). The mock-vaccinated groups remained negative till the end of the



Fig. 3. In vitro IFN- γ production after BRSV antigenic stimulation. Blood samples were tested before at the time of each vaccination (days 0, 29 and 60), at challenge (day 80) and at 6 and 12 days after challenge (days 86 and 92). The results are expressed as stimulation index. The bars represent the mean value for each group and the vertical bars represent the standard deviation of the means. The horizontal dashed bar represents the cut-off value. The stars indicate the means that differ significantly from the mean of the mock-vaccinated group.

experiment, while one animal of the Npur group show a SI above the cut-off on day 92.

3.2. BRSV-specific antibody response

BRSV-specific antibody response was measured by indirect immunofluorescence (Fig. 4). The pNsyn priming elicited low levels of antibody that increased either after the protein boost in all animals of the pNsyn–Npur group or after the challenge in animals of the pNsyn group. Injection of the N protein alone also induced specific seroconversion. At the time of challenge, the mean antibody titers of the 3 vaccinated groups were not significantly different from each other (p > 0.05). However, the antibody titers of the 3 vaccinated groups were significantly higher than those of the mockvaccinated group (p < 0.05). Seroconversion of the mock-vaccinated animals was only observed 12 days after challenge, consistent with a priming effect of the challenge.



Fig. 4. BRSV-specific antibody response. Sera taken at the time of each vaccination (days 0, 29 and 60), on the day of challenge (day 80) and on days 6 and 12 after challenge (days 86 and 92) were analyzed by indirect immunofluorescence. The results are expressed as the log 3 of the last positive dilution and they are presented as the mean value for each group. The vertical bars represent the standard deviation of the means.



Fig. 5. Total RNA was extracted from BAL fluids collected 3 days before and 3, 5, 7 and 11 days after challenge. The viral load was examined by quantitative real-time RT-PCR and expressed as the number of BRSV RNA copies per 10³ ACTB RNA copies. The bars represent the mean value for each group and the vertical bars represent the standard deviation of the means. The stars indicate the means that differ significantly from the mean of the control group.

The priming of a local response in the different vaccinated groups was investigated by measuring N-specific IgG and IgA in the BAL after challenge. None of the animals exhibited N-specific IgG response 3 days before challenge, and all the animals of the mock-and-pNsyn-vaccinated groups remained negative till the end of the experiment. IgG antibodies were detected in the Npur group from day 5 to 11 after the challenge with a maximum mean titer of 1/20 at D+5. In the pNsyn–Npur group, positive animals were recorded from 3 to 7 days after the challenge, with a maximum mean titer of 1/30, 5 days after the challenge (data not shown). In contrast, no N-specific IgA response was detected in the BAL, whatever the vaccinated group (data not shown).

3.3. Protection against BRSV challenge

3.3.1. Clinical signs

After challenge, no fever or increase of the respiratory rate were recorded. Three animals of the mock-vaccinated group experienced a mild cough lasting for at least 1 day after challenge, but this was not considered as significant (data not shown).

3.3.2. Viral RNA load in the BAL

The replication of challenged BRSV was followed in the BAL fluids collected every 2 or 3 days by quantitative real-time RT-PCR (Fig. 5). As expected, all the calves were negative 3 days before the challenge. Viral RNA was detected in the mock-vaccinated calves from day 3 after the challenge till the end of the experiment. However, at day 11, the infection has almost resolved, as the viral load in the BAL fluids was about 100 copies of viral RNA. Similar kinetic was observed in the calves vaccinated with the N protein, excepted that the virus was cleared on day 11. All animals primed with pNsyn showed lower BRSV RNA copy numbers in their BAL fluids than the 2 other groups. Indeed, for each animal and at each time point, the viral load was below 200 RNA copies/10³ ACTB copies. Also, each animal was positive at maximum 2 successive time points and all calves had cleared the virus on day 11 post-challenge. However, the great individual variability and the low number of animals in each group prevented the differences between the 2 groups vaccinated with pNsyn and the 2 other groups to be statistically significant, except on day 5 after challenge (p < 0.05).

3.3.3. Postmortem examination

One or 2 animals of each group were killed on day 6 after challenge while the 2 remaining animals of each group were killed on day 12.

The lungs were examined and the presence of pneumonic consolidation was recorded for each lobe. The number of affected lobes in each group is summarized in Table 2. In the mock-vaccinated group, all the calves showed lesions in the caudal part of the cranial lobe, and in the middle and accessory lobes. The cranial part of the cranial lobe was affected in 3 calves and the caudal one in 2 calves. In contrast, in the group vaccinated with both pNsyn and the N protein, only one calf showed one spot of consolidation in the cranial lobe (cranial part) and another in the caudal lobe. This group was significantly different from mock-vaccinated group (p < 0.05). The 2 other groups gave intermediate results (8 affected lobes/15), and the 6 animals showed lesions in the cranial part of the cranial lobe. However, the difference with the control group was not statistically supported. These lesions were also less severe in extent in these 2 groups than in the control group with the exception of one animal vaccinated with the N protein. All the lobes of this animal were affected and the extent of the lesions was comparable to those observed in the mock-vaccinated group.

Histological examination of the lungs revealed typical BRSV lesions. Histopathological lesions of the bronchi and bronchioles including lymphoid peribronchial cuffing (Fig. 6a and b), lymphoid infiltration, hyperplasia and metaplasia of the epithelium with desquamation and the presence of syncytial cells in the lumen (Fig. 6c and d). Infiltration of mononuclear inflammatory cells with consolidation and emphysema were recorded in the alveoli (Fig. 6e and f). However, irrespective of the slaughtering time post-challenge (day 6 or day 12), the lesions were less severe and more limited in the group vaccinated with pNsyn and Npur (Fig. 6a, c and e) than in the 3 other groups (Fig. 6b, d and f).

The viral replication in the lungs was analyzed by quantitative real-time RT-PCR (Table 3). No BRSV RNA was detected in the lungs of the calves vaccinated with pNsyn alone and in the calves primed with pNsyn and boosted with the N protein. Two out of the 3 calves vaccinated with the N protein alone were also protected. In contrast, BRSV RNA was detected in the lungs of all the mock-vaccinated animals.

4. Discussion

Numerous advantages have been described for the use of DNA vaccines. They theoretically combine the efficacy of live vaccines and the safety afforded by the inactivated vaccine [46]. They can circumvent the inhibitory effects of maternal antibodies [47] and they prime the TH1 component of the immune response [48]. This is of particular interest for overcoming the TH2 bias when vaccinating neonates against BRSV [49].

Several studies described the vaccination of mice with plasmids expressing the F or G glycoproteins [35,50,51]. These 2 antigens have also been tested as vaccine candidates in calves. A plasmid expressing the G glycoprotein afforded only partial reduction of viral shedding after challenge [52,53]. More recently,

Table 2		
Macroscopic	pneumonic	lesions

	Mock	Npur	pNsyn	pNsyn–Npur
Cranial: cranial part	3/4	3/3	3/3	1/4
Cranial: caudal part	4/4	1/3	1/3	0/4
Middle	4/4	2/3	3/3	0/4
Caudal	2/4	1/3	1/3	1/4
Accessory	4/4	1/3	0/3	0/4
Total	17/20	8/15	8/15	2/20

For each animal, all the pulmonary lobes were examined for the presence of pneumonic consolidation. The ratios of animals presenting lesions are indicated for each lobe. The total number of affected lobes in each group is also presented.



Fig. 6. Pulmonary histopathology after BRSV challenge in the pNsyn–Npur vaccinated group (a, c and e) and in the other groups: mock-vaccinated group (b and d), pNsyn-vaccinated group (f). (b) Lymphoid peribronchial cuffing (arrows). (d) Epithelial hyperplasia and metaplasia (arrows) with desquamation of the cells in the lumen (arrowhead). (f) Consolidation of the alveoli with infiltration of mononuclear inflammatory cells. (a) Absence of lymphoid peribronchial cuffing. (c) Absence of epithelial proliferation. (e) Absence of infiltration of mononuclear inflammatory cells.

DNA vaccine encoding the BRSV fusion protein induced significant protection against the virus challenge, by reducing the nasopharyngeal excretion of the virus and the extent of gross pneumonic lesions. However, complete protection against BRSV was not conferred [54]. Finally, DNA immunization of infant rhesus monkeys or young calves against both the BRSV F and nucleocapsid (N) proteins stimulated both humoral and cell-mediated immunity against the virus and reduced viral replication, clinical signs and pulmonary lesions after challenge [39,55]. Reports describing the use of the N protein are rather scarce. However, N was shown to be a major target of the immune response against RSV. Vaccination of mice with recombinant vaccinia virus encoding the (H)RSV N protein induced partial resistance [56,57]. Similarly, immunization of young calves with a recombinant vaccinia virus expressing the BRSV N protein induced non-neutralizing antibodies and primed the BRSV-specific proliferative response and IFN- γ production that resulted in reduction of viral replication in the upper and lower respiratory tract [21,28]. Overall, these data suggest that the protection afforded by the N protein might be largely mediated by cellular immunity.

The efficacy of DNA immunization against the N protein alone had never been studied before the present study. The purpose of this work was thus first to stimulate the cellular immune response using a DNA vaccine and then to enhance this response with a proteic boost. Therefore, we used of the nucleocapsid (N) protein of BRSV in a "prime-boost" vaccination strategy and we compare it to the use of DNA or protein vaccination alone.

In the present study, we show that both cellular and humoral arms of the immune response were stimulated by the vaccine combination. Injection of either pNsyn or Npur induced a BRSV-specific humoral response. The neutralizing activity of the antibodies was not investigated here, but antibodies against the N protein were shown non-neutralizing by others [21]. At the time of challenge, however, there was no significant difference between the vaccinated groups. Furthermore, vaccination did not prime for a local N-specific IgA response. This is in accordance with the results

Table 3

The viral loads in the lungs were examined by conducting quantitative real-time RT-PCR on total RNA extracted from homogenates of pieces from the different pulmonary lobes

Experimental group	Calf	BRSV RNA copies/10 ⁶ ACTB RNA copies
	6601	$5.5 imes 10^3$
. K − −1−	6604	$2.0 imes 10^4$
IVIOCK	6621	$5.2 imes 10^5$
	6628	1.2×10^3
	6615	-
Npur	6622	$6.8 imes 10^4$
	6626	-
	6596	-
pNsyn	6610	-
	6624	-
	6603	-
DNeve Neur	6611	-
piasyn-iapui	6612	-
	6620	-

(-) Not detected.

published by others. Indeed, vaccination with a recombinant vaccinia virus expressing the N protein induced local IgG1 but not IgA response [21]. A lymphoproliferative response was only induced in calves that received the pNsyn plasmid. This response was increased when calves were boosted with the N protein, resulting in significantly higher stimulation index compared to the 3 other groups. In contrast, only the prime–boost combination induced the production of IFN- γ .

After the challenge, the clinical and virological protections were evaluated. No clinical signs were recorded, even in the mock-vaccinated animals. This is in contrast with the previous experiment, where the same stock of BRSV 3761 was used [39]. The reasons are unclear but could be linked to the age or the genetic background of the animals [58]. However, the virus replicated in the respiratory tract of the mock-vaccinated animals, the highest viral load being observed in the BAL on day 5 post-infection. At autopsy, the viral load in the lungs was as high as previously described [39] and gross pneumonic as well as histopathological lesions were observed. After injection of the N protein, 2 out the 3 calves were protected against BRSV replication in the lungs. However, high viral loads were detected in the BAL fluids, and protection against pneumonic lesions was only partial. DNA vaccination alone was able to protect against viral replication in the lungs and to strongly reduce the viral replication in the BAL but the calves were not totally protected against the development of gross pneumonic lesions. The presence of lung consolidations at autopsy indicates the spread of the virus in the lungs. Indeed, pathological changes have been shown to appear when replication of BRSV spreads to the alveoli and may continue after clearance of the virus [59]. In contrast, protection against both the viral replication in the lungs and the gross and histological pneumonic lesions was afforded by the combination of DNA and protein vaccination, suggesting that infectious virus was cleared before infecting the alveoli.

The role of the cellular immunity in the protection against RSV and the pathogenesis induced by the virus is controversial. Bovine IFN- γ has been shown to induce a TH1 response, characterized by IgG2 production [60]. On one hand, both CD4⁺ and CD8⁺ T cells can eliminate the virus but they also cause immunopathology in mice [61]. IFN- γ was identified as a key molecule involved in both the virus control and the immunopathology induced by CD8⁺ T cells [62]. In mice, TH1 and TH2 biased T cell responses could either co-exist or down regulate each other [14,63]. On the other hand, impaired viral clearance and enhanced disease observed after vac-

cination with FI vaccine have been correlated with decreased IFN- γ production [64]. Formulation of FI-BRSV with CpG ODN resulted in increased IFN- γ secretion and reduction of both gross lung pathology and viral replication in calves and in mice [65,66]. Furthermore, Modified Live vaccine induced cellular immunity, in particular secretion of IFN- γ by peripheral blood leukocytes, appeared to be a more consistent correlate of protection than pre-challenge serum antibody [67]. Our data are consistent with these results. Indeed, the only animals that were protected against both viral replication in the lungs and the development of macroscopic lung lesions resulting of BRSV challenge were those that had been vaccinated successively with pNsyn and N protein and that exhibited significant IFN- γ production at the time of the challenge.

Injection of the purified N protein, mixed with Quil A did not induce measurable lymphoproliferative response. This was not anticipated because other authors showed that insect cells infected with recombinant baculovirus expressing the F protein and mixed with Quil A stimulate the cellular-mediated immune response in lambs [68]. However, as baculovirus has been shown as stimulator of immune response, this could contribute to the overall effect of the immunogen [69]. In our experiment, the N protein was nevertheless able to prime for a lymphoproliferative response, detected after BRSV challenge, and this might be attributed to a TH1 triggering of the adjuvant. After vaccination with the N protein, 2 out of the 3 calves were protected against viral replication in the lungs. Virus was detected in the lungs of the remaining calf (calf 6622). Interestingly, this calf had also higher viral load in BAL fluids and more extended lesions of the lungs compared to the 2 other calves of this group. The 3 calves showed identical serological responses. However, when the lymphoproliferative response was individually examined on blood collected 6 days after challenge, the SI was below 1 for calf 6622 and of about 3 for the 2 other calves (data not shown). We hypothesize that the development of pulmonary lesions and the viral replication in the lungs were correlated with the absence of cell-mediated immunity.

The prime-boost strategy used here was superior to DNA or protein vaccination alone in term of protection for large animals. This is in accordance with the results published by others. Indeed, this strategy was recently shown to induce stronger humoral immune response and better protection against BVDV in cattle compared to immunization with DNA or protein alone [70]. Boosting DNA vaccinated mice with recombinants vectors can elicit strong CD4 and CD8 immune responses [71]. Furthermore, it has been shown that the longevity of T-cell stimulation was higher after DNA prime-MLV boost than after MLV vaccination alone against BoHV-1 in cattle [72]. Boosting CD8 response with inactivated virus or recombinant proteins is more controversial. While inactivated viral boost was shown to enhance the CD4 and antibody responses but not the CD8 response against severe acute respiratory syndrome coronavirus in mice, other authors described an enhancement of CD8⁺ CTL and CD4⁺ T-helper cells against HIV after a protein boost in nonhuman primates [73,74]. Here also, although the cytotoxic T cell response was not investigated, increased lymphoproliferative response and IFN- γ production were observed, suggesting an enhancement of the cellular response after the protein boost.

In conclusion, the results presented here show protection against BRSV replication in the lung and against lung pathology after nucleocapsid-based DNA prime–protein boost protocol. This vaccination strategy elicited both humoral and cellular immune responses. The observed protection correlated with the lymphoproliferative response and the interferon- γ production. The results also confirm the efficacy of the prime–boost strategy in a large animal model. This strategy is promising for the protection against RSV in cattle and it could also open perspectives for vaccinating young infants as well. Further work should be performed to confirm the present results in neonatal calves harboring maternal antibodies. Finally, further development would combine the nucleocapsid and the fusion proteins in a DNA prime–protein boost strategy in order to stimulate protective cellular immunity and neutralizing antibodies at once.

Acknowledgments

This study was supported by the Federal Public Service Health, Food chain safety and environment (Brussels, Belgium). The authors are grateful to C. Giannoulis for the purification of plasmid DNA, to B. Lambrecht for providing the Sf9 cells, to Stéphanie Durand, Caroline Rodeghiero, P. Van Muylem and Rita Geeroms for their skilled technical assistance and to the staff of the animal facilities of the INRA of Nouzilly for taking care of the animals.

References

- Brandenburg AH, Neijens HJ, Osterhaus ADME. Pathogenesis of RSV lower respiratory tract infection: implications for vaccine development. Vaccine 2001;19:2769–82.
- [2] Larsen LE. Bovine respiratory syncytial virus (BRSV): a review. Acta Vet Scand 2000;41:1–24.
- [3] Van der Poel WH, Brand A, Kramps JA, Van Oirschot TJ. Respiratory syncytial virus infections in human beings and in cattle. J Infect 1994;29:215–28.
- [4] Belknap EB, Baker JC, Patterson JS, Walker RD, Haines DM, Clark EG. The role of passive immunity in bovine respiratory syncytial virus-infected calves. J Infect Dis 1991;163:470–6.
- [5] Kimman TG, Westenbrink F, Schreuder BEC, Straver PJ. Local and systemic antibody response to bovine respiratory syncytial virus infection and reinfection in calves with and without maternal antibodies. J Clin Microbiol 1987;25:1097–106.
- [6] Kimman TG, Zimmer GM, Westenbrink F, Mars J, van Leeuwen E. Epidemiological study of bovine respiratory syncytial virus infections in calves: Influence of maternal antibodies on the outcome of disease. Vet Rec 1988;123:104–9.
- [7] Ellis JA. The immunology of the bovine respiratory disease complex. Vet Clin North Am Food Anim Pract 2001;17:535–50.
- [8] Peters AR, Thevasagayam SJ, Wieseman A, Salt JS. Duration of immunity of a quadrivalent vaccine against respiratory diseases caused by BHV-1, PIV3, BVDV, and BRSV in experimentally infected calves. Prev Vet Med 2004;66:63–77.
- [9] Patel JR, Didlick SA. Evaluation of efficacy of an inactivated vaccine against BRSV in calves with maternal antibodies. Am J Vet Res 2004;65:417–21.
- [10] Ellis J, Gow S, West K, Waldner C, Rhodes C, Mutwiri G, et al. Response of calves to challenge exposure with virulent BRSV following intranasal administration of vaccines formulated for parenteral administration. J Am Vet Med Assoc 2007;15:233–43.
- [11] Kim HW, Canchola JG, Brandt CD, Pyles G, Chanock RM, Jensen K, et al. Respiratory syncytial virus disease in infants despite prior administration of antigenic inactivated vaccine. Am J Epidemiol 1969;89:422–34.
- [12] Gershwin LJ, Schelegle ES, Gunther RA, Anderson ML, Woolums AR, Larochelle DR, et al. A bovine model of vaccine enhanced respiratory syncytial virus pathophysiology. Vaccine 1998;16:1225–36.
- [13] Hussell T, Baldwin CJ, O'Garra A, Openshaw PJ. CD8+ T cells control Th2driven pathology during pulmonary respiratory syncytial virus infection. Eur J Immunol 1997;27:3341–9.
- [14] Varga SM, Wang X, Welsh RM, Braciale TJ. Immunopathology in RSV infection is mediated by a discrete oligoclonal subset of antigen-specific CD4(+) T cells. Immunity 2001;15:637–46.
- [15] Connors M, Giese NA, Kulkarni AB, Firestone CY, Morse III HC, Murphy BR. Enhanced pulmonary histopathology induced by respiratory syncytial virus (RSV) challenge of formalin-inactivated RSV-immunized BALB/c mice is abrogated by depletion of interleukin-4 (IL-4) and IL-10. J Virol 1994;68:5321–5.
- [16] Antonis AFG, Schrijver RS, Daus F, Steverink PJGM, Stockhofe N, Hensen EJ, et al. Vaccine-induced immunopathology during bovine respiratory syncytial virus infection: exploring the parameters of pathogenesis. J Virol 2003;77:12067–73.
- [17] Gershwin LJ, Gunther RA, Anderson ML, Woolums AR, McArthur- Vaughan K, Randel KE, et al. Bovine respiratory syncytial virus-specific IgE is associated with interleukin-2 and -4, and interferon-gamma expression in pulmonary lymph of experimentally infected calves. Am J Vet Res 2000;61:291–8.
- [18] Kalina WV, Woolums AR, Berghaus RD, Gershwin LJ. Formalin-inactivated bovine RSV vaccine enhances a Th2 mediated immune response in infected cattle. Vaccine 2004;22:1465–74.
- [19] Schreiber P, Matheise JP, Dessy F, Heimann M, Letesson J-J, Coppe P, et al. High mortality rate associated with bovine respiratory syncytial virus (BRSV) infection in Belgian white blue calves previously vaccinated with an inactivated BRSV vaccine. J Vet Med B Infect Dis Vet Public Health 2000;47:535–50.
- [20] Larsen LE, Tegtmeier C, Pedersen E. Bovine respiratory syncytial virus (BRSV) pneumonia in beef calf herds despite vaccination. Acta Vet Scand 2001;42:113–21.

- [21] Taylor G, Thomas LH, Furze JM, Cook RS, Wyld SG, Lerch R, et al. Recombinant vaccinia viruses expressing the F, G or N, but not the M2, protein of bovine respiratory syncytial virus (BRSV) induce resistance to BRSV challenge in the calf and protect against the development of pneumonic lesions. J Gen Virol 1997;78:3195–206.
- [22] Thomas LH, Cook RS, Wyld SG, Furze JM, Taylor G. Passive protection of gnotobiotic caves using monoclonal antibodies directed at different epitopes on the fusion protein of bovine respiratory syncytial virus. J Infect Dis 1998;177:874–80.
- [23] Taylor G, Thomas LH, Wyld SG, Furze JM, Sopp P, Howard CJ. Role of Tlymphocyte subsets in recovery from respiratory syncytial virus infection in calves. J Virol 1995;69:6658–64.
- [24] Braciale TJ. Respiratory Syncytial virus and T cells Interplay between the virus and the host adaptive immune system. Proc Am Thorac Soc 2005;2:141–6.
- [25] Openshaw PJ, Tregoning JS. Immune responses and disease enhancement during respiratory syncytial virus infection. Clin Microbiol Rev 2005;18:541–55.
- [26] Bangham CR, Openshaw PJ, Ball LA, King AM, Wertz GW, Askonas BA. Human and murine cytotoxic T cells specific to respiratory syncytial virus recognize the viral nucleoprotein (N), but not the major glycoprotein (G), expressed by vaccinia virus recombinants. J Immunol 1986;137:3973–7.
- [27] Connors M, Kulkarni AB, Collins PL, Firestone C-Y, Holmes KL, Morse HC, et al. Resistance to respiratory syncytial virus (RSV) challenge induced by infection with a vaccinia virus recombinant expressing the RSV M2 protein is mediated by CD8+T cells, while that induced by Vac-F or Vac-G recombinants is mediated by antibodies. J Virol 1992;66:1277–81.
- [28] Gaddum RM, Cook RS, Furze JM, Ellis SA, Taylor G. Recognition of bovine respiratory syncytial virus proteins by bovine CD8+ T lymphocytes. Immunology 2003;108:220–9.
- [29] Antonis AFG, Claassen EAW, Hensen EJ, de Groot RJ, de Groot-Mijnes JDF, Schrijver RS, et al. Kinetics of antiviral CD8 T cell responses during primary and post-vaccination secondary bovine respiratory syncytial virus infection. Vaccine 2006;24:1551–61.
- [30] Gaddum RM, Cook RS, Thomas LH, Taylor G. Primary cytotoxic T-cell responses to bovine respiratory syncytial virus in calves. Immunology 1996;88:421–7.
- [31] McInnes E, Sopp P, Howard CJ, Taylor G. Phenotypic analysis of local cellular responses in calves infected with bovine respiratory syncytial virus. Immunology 1999;96:396–403.
- [32] Rutigliano JA, Johnson TR, Hollinger TN, Fischer JE, Aung S, Graham BS. Treatment with anti-LFA-1 delays the CD8+ cytotoxic-T-lymphocyte response and viral clearance in mice with primary respiratory syncytial virus infection. J Virol 2004;78:3014–23.
- [33] Srikiatkhachorn A, Braciale TJ. Virus-specific CD8+ T lymphocytes downregulate T helper cell type 2 cytokine secretion and pulmonary eosinophilia during experimental murine respiratory syncytial virus infection. J Exp Med 1997;186:421–32.
- [34] Robinson HL, Pertmer TM. DNA vaccines for viral infections: basic studies and applications. Adv Virus Res 2000;55:1–74.
- [35] Bembridge GP, Rodriguez N, Garcia-Beato R, Nicolson C, Melero JA, Taylor G. DNA encoding the attachment (G) or fusion (F) protein of respiratory syncytial virus induces protection in the absence of pulmonary inflammation. J Gen Virol 2000;81:2519–23.
- [36] Brady RP, Topliff CL, Kelling CL. In vitro expression of full-length and truncated bovine respiratory syncytial virus G proteins and their antibody responses in BALB/c mice. Vaccine 2004;22:3762–8.
- [37] Babiuk LA, Pontarollo R, Babiuk S, Loehr B, van Drunen Littel-van den Hurk S. Induction of immune response by DNA vaccines in large animals. Vaccine 2003;21:649–58.
- [38] Toussaint JF, Letellier C, Paquet D, Dispas M, Kerkhofs P. Prime-boost strategies combining DNA and inactivated vaccines confer high immunity and protection in cattle against bovine herpesvirus-1. Vaccine 2005;23:5073–81.
- [39] Boxus M, Tignon M, Toussaint J-F, Roels S, Benoit MA, Walravens K, et al. DNA immunization against fusion and nucleocapsid proteins protects calves against BRSV. J Virol 2007;81:6879–89.
- [40] Martinez X, Li X, Kovarik J, Klein M, Lambert P-H, Siegrist C-A. Combining DNA and protein vaccination for early life immunization against RSV in mice. Eur J Immunol 1999;29:3390–400.
- [41] Samal SK, Zamora M, McPhillips TH, Mohanty SB. Molecular cloning and sequence analysis of Bovine Respiratory Syncytial Virus mRNA encoding the major nucleocapsid protein. Virology 1991;180:453–6.
- [42] Tjornehoj K, Uttenthal A, Viuff B, Larsen LE, Rontved C, Ronsholt L. An experimental infection model for reproduction of calf pneumonia with bovine respiratory syncytial virus (BRSV) based on one combined exposure of calves. Res Vet Sci 2003;74:55–65.
- [43] Wellemans G, Leunen J. La Rhino-trachéite infectieuse des bovins et sa sérologie. Ann Méd Vét 1973;117:507518.
- [44] Uttenthal A, Larsen LE, Philipsen JS, Tjornehoj K, Viuff B, Nielsen KH, et al. Antibidy dynamics in BRSV-infected Danish dairy herds as determined by isotype-specific immunoglobulins. Vet Microbiol 2000;76:329–41.
- [45] Boxus M, Letellier C, Kerkhofs P. Real Time RT-PCR for the detection and quantitation of bovine respiratory syncytial virus. J Virol Methods 2005;125:125– 30.
- [46] Gurunathan S, Klinman DM, Seder RA. DNA vaccines: immunology, application, and optimization. Annu Rev Immunol 2000;18:927–74.
- [47] Van Drunen Little-van den Hurk S, Braun RP, Lewis PJ, Karvonen BC, Babiuk LA, Griebel PJ. Immunization of neonates with DNA encoding a bovine herpesvirus

glycoprotein is effective in the presence of maternal antibodies. Viral Immunol 1999;12:67–77.

- [48] Martinez X, Brandt C, Saddallah F, Tougne C, Barrios C, Wild F, et al. DNA immunization circumvents deficient induction of TH1 and cytotoxic T cells lymphocyte responses in neonates and during early life. PNAS 1997;94:8726–31.
- [49] Morein B, Abusugra I, Blomqvist G. Immunity in neonates. Vet Immunol Immunopathol 2002;87:207–13.
- [50] Li X, Sambhara S, Li CX, Ewasyshyn M, Parrington M, Caterini J, et al. Protection against respiratory syncytial virus infection by DNA immunization. J Exp Med 1998;188:681–8.
- [51] Li X, Sambhara S, Li CX, Ettorre L, Switzer I, Cates G, et al. Plasmid DNA encoding the respiratory syncytial virus G protein is a promising vaccine candidate. Virology 2000;269:54–65.
- [52] Schrijver RS, Langedijk JP, Keil GM, Middel WGJ, Maris-Veldhuis M, Van Oirschot JT, et al. Immunization of cattle with a BHV1 vector vaccine or a DNA vaccine both coding for the G protein of BRSV. Vaccine 1997;15:1908–16.
- [53] Schrijver RS, Langedijk JP, Keil GM, Middel WGJ, Maris-Veldhuis M, Van Oirschot JT, et al. Comparison of DNA application methods to reduce BRSV shedding in cattle. Vaccine 1998;16:130-4.
- [54] Taylor G, Bruce C, Barbet AF, Wyld SG, Thomas LH. DNA vaccination against respiratory syncytial virus in young calves. Vaccine 2005;23:1242–50.
- [55] Vaughan K, Rhodes GH, Gershwin LJ. DNA immunization against respiratory syncytial virus (RSV) in infant rhesus monkeys. Vaccine 2005;23:2928–42.
- [56] King AMQ, Stott EJ, Langer SJ, Young KK-Y, Ball LA, Wertz GW. Recombinant vaccinia virus carrying the N gene of Human respiratory syncytial virus: studies of gene expression in cell culture and immune response in mice. J Virol 1987;61:2885–90.
- [57] Connors M, Collins PL, Firestone C-Y, Murphy B. Respiratory syncytial virus (RSV) F, G, M2, and N proteins each induce resistance to RSV challenge, but resistance induced by M2 and N proteins is relatively short-lived. | Virol 1991;65:1634–7.
- [58] Grell SN, Riber U, Tjornehoj K, Larsen LE, Heegaard PMH. Age-dependent differences in cytokine and antibody responses after experimental RSV infection in a bovine model. Vaccine 2005;23:3412–23.
- [59] Viuff B, Tjornehoj K, Larsen LE, Rontved C, Uttenthal A, Ronsholt L, et al. Replication and clearance of respiratory syncytial virus apoptosis is an important pathway of virus clearance after experimental infection with BRSV. Am J Pathology 2002; 161:2195–207.
- [60] Estes DM, Brown WC. Type 1 and type 2 responses in regulation of Ig isotype expression in cattle. Vet Immunol Immunopathol 2002;90:1–10.
- [61] Cannon MJ, Openshaw PJ, Askonas BA. Cytotoxic T cells clear virus but augment lung pathology in mice infected with respiratory syncytial virus. J Exp Med 1988;168:1163–8.
- [62] Ostler T, Davidson W, Ehl S. Virus clearance and immunopathology by CD8(+) T cells during infection with respiratory syncytial virus are mediated by IFNgamma. Eur J Immunol 2002;32:2117–23.

- [63] Aung S, Tang YW, Graham BS. Interleukin-4 diminishes CD8(+) respiratory syncytial virus-specific cytotoxic T lymphocyte activity in vivo. J Virol 1999;73:8944–9.
- [64] Woolums AR, Singer RS, Boyle GA, Gershwin LJ. Interferon gamma production during bovine respiratory syncytial virus (BRSV) infection is diminished in calves vaccinated with formalin-inactivated BRSV. Vaccine 1999;17:1293– 7.
- [65] Oumouna M, Mapletoft JW, Karvonen BC, Babiuk LA, van Drunen Littel-van den Hurk S. Formulation with CpG oligodeoxynucleotides prevents induction of pulmonary immunopathology following priming with formalin-inactivated or commercial killed bovine respiratory syncytial virus vaccine. J Virol 2005;79:2024–32.
- [66] Mapletoft JW, Oumouna M, Townsend HG, Gomis S, Babiuk LA, van Drunen Littel-van den Hurk S. Formulation with CpG oligodeoxynucleotides increases cellular immunity and protection induced by vaccination of calves with formalin-inactivated bovine respiratory syncytial virus. Virology 2006;353:316–23.
- [67] West K, Petrie L, Konoby C, Haines DM, Cortese V, Ellis JA. The efficacy of modified-live bovine respiratory syncytial virus vaccines in experimentally infected calves. Vaccine 2000;18:907–19.
- [68] Sharma AK, Woldehiwet Z, Walravens K, Letesson J. Immune response of lambs to the fusion glycoprotein of BRSV expressed on insect cells infected with a recombinant baculovirus. Vaccine 1996;14:773–9.
- [69] Abe T, Takahashi H, Hamazaki H, Miyano-kurosaki N, Matsuura Y, Takaku H. Baculovirus induces an innate immune response and confers protection from lethal Influenza virus infection in mice. J Immunol 2003;171:1133–9.
- [70] Liang R, van den Hurk JV, Babiuk LA, van Drunen Littel-van den Hurk S. Priming with DNA encoding E2 and boosting with E2 protein formulated with CpG oligodeoxynucleotides induces strong immune responses and protection from Bovine viral diarrhea virus in cattle. J Gen Virol 2006;87:2971–82.
 [71] Estcourt MJ, Ramsay AJ, Brooks A, Thomson SA, Medveckzy CJ, Ramshaw IA.
- [71] Estcourt MJ, Ramsay AJ, Brooks A, Thomson SA, Medveckzy CJ, Ramshaw IA. Prime-boost immunization generates a high frequency, high-avidity CD8(+) cytotoxic T lymphocyte population. Int Immunol 2002;14:31–7.
- [72] Loehr BI, Pontarollo R, Rankin R, Latimer L, Willson P, Babiuk LA, et al. Priming by DNA immunization augments T-cell responses induced by modified live bovine herpesvirus vaccine. J Gen Virol 2001;82:3035–43.
- [73] Kong W-p, Xu I, Stadler K, Ulmer JB, Abrignani S, Rappuoli R, et al. Modulation of the immune response to the severe acute respiratory syndrome spike glycoprotein by gene-based and inactivated virus immunization. J Virol 2005;79:13915–23.
- [74] Cristillo AD, Wang S, Caskey MS, Unangst T, Hocker L, He L, et al. Preclinical evaluation of cellular immune responses elicited by a polyvalent DNA prime/protein boost HIV-1 vaccine. Virology 2006;346:151–68.