

## RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

### **Indole-pyridinyl-ethanones as novel inhibitors of indoleamine-2,3 dioxygenase (IDO), a promising target for anti-cancer immunotherapy**

Dolusic, Eduard; Larrieu, Pierre; Blanc, Sébastien; Moineaux, Laurence; Sapunarcic, Frédéric; Pouyez, Jenny; Colette, Delphine; Fraser, Graeme; Stroobant, Vincent; Pilotte, Luc; Colau, Didier; Frère, Jean-Marie; Masereel, Bernard; Van den Eynde, Benoît; Wouters, Johan; Frédérick, Raphaël

*Publication date:*  
2011

*Document Version*  
Early version, also known as pre-print

[Link to publication](#)

*Citation for published version (HARVARD):*

Dolusic, E, Larrieu, P, Blanc, S, Moineaux, L, Sapunarcic, F, Pouyez, J, Colette, D, Fraser, G, Stroobant, V, Pilotte, L, Colau, D, Frère, J-M, Masereel, B, Van den Eynde, B, Wouters, J & Frédérick, R 2011, 'Indole-pyridinyl-ethanones as novel inhibitors of indoleamine-2,3 dioxygenase (IDO), a promising target for anti-cancer immunotherapy', 47th RICT: Drug Discovery and Selection., Lyon, France, 6/07/11 pp. Book of Abstracts, 47èmes RICT, 6-8 juillet 2011, Lyon, France, IS08, p. 136.

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Indole-pyridinyl-ethanones as Novel Inhibitors of Indoleamine 2,3-Dioxygenase (IDO), a Promising Target for Anti-Cancer Immunotherapy



Eduard Dolušić<sup>a</sup>, Sébastien Blanc<sup>b</sup>, Pierre Larriue<sup>c</sup>, Laurence Moineaux<sup>a</sup>, Delphine Colette<sup>b</sup>, Graeme Fraser<sup>b</sup>, Vincent Stroobant<sup>c</sup>, Luc Pilotte<sup>c</sup>, Didier Colau<sup>c</sup>, Johan Wouters<sup>a</sup>, Bernard Masereel<sup>a</sup>, Benoît Van den Eynde<sup>c</sup> and Raphaël Frédérick<sup>a</sup>

<sup>a</sup>Drug Design and Discovery Center, University of Namur (FUNDP), 61 Rue de Bruxelles, B-5000 Namur, Belgium;


<sup>b</sup>Euroscreen SA, 47 Rue Adrienne Boland, B-6041 Gosselies, Belgium;

<sup>c</sup>Ludwig Institute for Cancer Research, Université Catholique de Louvain, 74 Avenue Hippocrate, B-1200 Brussels, Belgium  
edolusic@fundp.ac.be

**Introduction.** Immunotherapy is a promising novel strategy for cancer therapy. However, this approach showed a limited efficacy *in vivo* because cancer cells can develop mechanisms allowing tumors to resist or escape immune rejection.

IDO (EC 1.13.11.52), a heme dioxygenase, is expressed constitutively in many human tumors and its role in a tumoral immune resistance mechanism has been proved,<sup>1</sup> justifying the interest in IDO inhibitors.<sup>2</sup>


**Aim of the Work.** We sought to develop a novel series of IDO inhibitors starting with a virtual screening of a database of commercially-available compounds.




**Virtual screening.** Based on recent results such as structural findings<sup>3</sup> and rational design of IDO inhibitors,<sup>4</sup> we applied virtual screening of the ZINC database (<http://zinc.docking.org>) for the discovery of new inhibitors (Figure 1). The most promising candidates were purchased and tested *in vitro*. 1-(1*H*-Indol-2-yl)-2-pyridin-3-yl-ethanone (**7a**; IC<sub>50</sub> = 65 μM) was selected for pharmacomodulation (Schemes 1 and 2 and Table 1).

**Figure 1.** IDO Virtual screening flowchart. (i) fragment library, (ii) goldscore > 50, (iii) Cscore ≥ 4, (iv) visual analysis and selection, (v) really commercially-available.

## Synthesis



**Scheme 1.** General synthetic scheme for indol-2-yl ethanones.<sup>5</sup> Reagents and conditions: (i), LDA, THF / hexanes, -78 to 0°C, 1 h; (ii) THF / hexanes, 0°C to r.t., 16-24 h; (iii) SOCl<sub>2</sub>, Δ, 15 min; (iv), hexane, 200°C (μW), 5 min; (v), N<sub>3</sub>CH<sub>2</sub>CO<sub>2</sub>Et, NaOEt, EtOH, -10 to 4°C, 1.5 - 20 h; (vi), MeI, K<sub>2</sub>CO<sub>3</sub>, DMF, 80°C, 5 days; (vii), AlCl<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, 0°C to r. t., 24 h; (viii), HCO<sub>2</sub>NH<sub>4</sub>, Pd black, MeOH, r. t., 1 h; (ix) H<sub>2</sub> (1 atm), 3% Pd/C, EtOH, r. t., 45 min; (x) PhB(OH)<sub>2</sub>, Pd(PPh<sub>3</sub>)<sub>4</sub>, K<sub>2</sub>CO<sub>3</sub>, EtOH/toluene 1/1, Δ, 20 h.



**Scheme 2.** Synthesis of compounds bearing different linker groups. Reagents and conditions: (i) HCO<sub>2</sub>NH<sub>4</sub>, Pd black, MeOH, r. t., 3 days; (ii), H<sub>2</sub>NNH<sub>2</sub>, KOH, (CH<sub>2</sub>OH)<sub>2</sub>, μW, 1h then aq. NH<sub>4</sub>Cl; (iii), NH<sub>2</sub>OH·HCl, pyridine, EtOH, μW (120°C), 30 min; (iv), SOCl<sub>2</sub>, Δ, 15 min then 3-aminopyridine, DIPEA, THF, 0°C → r. t., 30 min.


**Conclusion.** The synthesis and SAR of a novel series of IDO inhibitors are described.

Starting from the lead compound **7a** (IC<sub>50</sub> = 65 μM) identified through a virtual screening procedure, up to a 5-fold improvement in *in vitro* potency could be achieved by introducing small substituents in the 5- and 6-positions of the indole nucleus. Most modifications of the aromatic moieties are tolerated. On the contrary, the presence of an iron chelating group on the linker seems to be mandatory, as corroborated by the docking experiments (Fig. 2). A number of compounds are also moderately active in the *in vivo* assay, thus opening possibilities for further pharmacological evaluation.

**Table 1. Biological evaluation**

compound	R	R'	enzymatic assay cell assay inh. %	
			IC <sub>50</sub> (μM)	@ 20 μM
7a	H	H	65	13
7b	3-Br	H	>100	NI
7c	4-F	H	153	12
7d	4-OCH <sub>3</sub>	H	58	NI
7e	4-OH	H	83	NI
7f	5-F	H	36	24
7g	5-Cl	H	25	24
7h	5-Br	H	18	NI
7i	5-CH <sub>3</sub>	H	87	21
7j	5-OCH <sub>3</sub>	H	49	-
7k	5-NO <sub>2</sub>	H	>100	NI
7l	5-Ph	H	96	NI
7m	5-OCF <sub>3</sub>	H	13	12
7o	5-OH	H	37	NI
7p	6-F	H	43	NI
7q	7-OCH <sub>3</sub>	H	82	-
7r	4-OBn, 5-OCH <sub>3</sub>	H	>100	NI
7s	4-OH, 5-OCH <sub>3</sub>	H	63	11
7t	4,6-diCl	H	139	NI
7u	4,6-di(OCH <sub>3</sub> )	H	45	toxic
7v	H	CH <sub>3</sub>	141	NI
8a	H	H	37	10
9a	H	H	>100	NI
10a	H	H	26	NI
11a	H	H	29	NI
12a	H	CH <sub>3</sub>	34	NI
13	H	H	>100	NI
14	H	H	>100	NI
15	H	H	>100	25
16	H	-	94	NI

NI = no inhibition



**Figure 2.** Docking of compound **7a** inside the IDO active site showing interaction of the carbonyl oxygen with the heme iron

**References.** [1] Uyttenhove, C. *et al*, *J. Nat Med* **2003**, *9*, 1269-1274; [2] Macchiarulo *et al*, *Inflamm. Res.* **2009**, *37*, 219-229; [3] Sugimoto H. *et al*, *PNAS* **2006**, *103*, 2611-2616. [4] Röhrig, U. *et al*, *J. Med. Chem.* **2010**, *53*, 1172-1189; [5] Sundberg, R. *et al*, *J. Org. Chem.* **1978**, *43*, 4859-4865. This work is supported in part by the FNRS and Biowin (CANTOL : Convention n° 5678).