

## RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

### Towards more predictive and interdisciplinary climate change ecosystem experiments

Rineau, Francois; Malina, Robert; Beenaerts, Natalie; Arnauts, Natascha; Bardgett, Richard D.; Berg, Matty P.; Boerema, Annelies; Bruckers, Liesbeth; Clerinx, Jan; Davin, Edouard L.; De Boeck, Hans J.; De Dobbelaer, Tom; Dondini, Marta; De Laender, Frederik; Ellers, Jacintha; Franken, Oscar; Gilbert, Lucy; Gudmundsson, Lukas; Janssens, Ivan A.; Johnson, David; Lizin, Sebastien; Longdoz, Bernard; Meire, Patrick; Meremans, Dominique; Milbau, Ann; Moretti, Michele; Nijs, Ivan; Nobel, Anne; Pop, Sorin; Puetz, Thomas; Reyns, Wouter; Roy, Jacques; Schuetz, Jochen; Seneviratne, Sonia I.; Smith, Pete; Solmi, Francesca; Staes, Jan; Thiery, Wim; Thijs, Sofie; Vanderkelen, Inne; Van Landuyt, Wouter; Verbruggen, Erik; Witters, Nele; Zscheischler, Jakob; Vangronsveld, Jaco

*Published in:*

Nature Climate Change

*DOI:*

[10.1038/s41558-019-0609-3](https://doi.org/10.1038/s41558-019-0609-3)

*Publication date:*

2019

*Document Version*

Peer reviewed version

[Link to publication](#)

*Citation for published version (HARVARD):*

Rineau, F, Malina, R, Beenaerts, N, Arnauts, N, Bardgett, RD, Berg, MP, Boerema, A, Bruckers, L, Clerinx, J, Davin, EL, De Boeck, HJ, De Dobbelaer, T, Dondini, M, De Laender, F, Ellers, J, Franken, O, Gilbert, L, Gudmundsson, L, Janssens, IA, Johnson, D, Lizin, S, Longdoz, B, Meire, P, Meremans, D, Milbau, A, Moretti, M, Nijs, I, Nobel, A, Pop, S, Puetz, T, Reyns, W, Roy, J, Schuetz, J, Seneviratne, SI, Smith, P, Solmi, F, Staes, J, Thiery, W, Thijs, S, Vanderkelen, I, Van Landuyt, W, Verbruggen, E, Witters, N, Zscheischler, J & Vangronsveld, J 2019, 'Towards more predictive and interdisciplinary climate change ecosystem experiments', *Nature Climate Change*, vol. 9, no. 11, pp. 809-816. <https://doi.org/10.1038/s41558-019-0609-3>

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

1 **Improving the predictive power and interdisciplinarity of climate change experiments**

2

3 Francois Rineau 1\*, Robert Malina 2, Natalie Beenaerts 1, Natascha Arnauts 1, Richard D. Bardgett 3,  
4 Matty P. Berg 4,5, Annelies Boerema 6, Liesbeth Bruckers 7, Jan Clerinx 1, Edouard L. Davin 8, Hans J  
5 De Boeck 9, Tom De Dobbelaer 10, Marta Dondini 11, Frederik De Laender 12, Jacintha Ellers 4, Oscar  
6 Franken 4, Lucy Gilbert 13, Lukas Gudmundsson 8, Ivan A. Janssens 9, David Johnson 3, Sebastien  
7 Lizin 2, Bernard Longdoz 14, Patrick Meire 6, Dominique Meremans 6, Ann Milbau 10, Michele  
8 Moretti 2,15, Ivan Nijs 9, Anne Nobel 2, Sorin Pop 16, Thomas Puetz 17, Wouter Reyns 1,12, Jacques  
9 Roy 18, Jochen Schuetz 16, Sonia I. Seneviratne 8, Pete Smith 11, Francesca Solmi 7, Jan Staes 6, Wim  
10 Thierry 8,19, **Sofie Thijs 1**, Inne Vanderkelen 19, Wouter Van Landuyt 10, Erik Verbruggen 9, Nele  
11 Witters 2, Jakob Zscheischler 8,20,21, Jaco Vangronsveld 1, 22

12 \* Corresponding author: Francois Rineau, Environmental Biology, Centre for Environmental Sciences,  
13 Hasselt University, Diepenbeek, Belgium. Email: francois.rineau@uhasselt.be

14

15 1 Hasselt University, Environmental Biology, Centre for Environmental Sciences, Agoralaan Gebouw  
16 D, 3590 Diepenbeek, Belgium

17 2 Hasselt University, Environmental Economics, Centre for Environmental Sciences, Agoralaan  
18 Gebouw D, 3590 Diepenbeek, Belgium

19 3 The University of Manchester, School of Earth and Environmental Sciences, Michael Smith Building,  
20 Manchester M13 9PT, UK

21 4 Vrije Universiteit Amsterdam, Department of Ecological Science, De Boelelaan 1085, 1081HV  
22 Amsterdam, The Netherlands

23 5 Groningen University, Groningen Institute for Evolutionary Life Science, Community and  
24 Conservation Ecology Group, Nijenborgh 7, 9747AG Groningen, The Netherlands

25 6 University of Antwerp, Ecosystem Management research group (ECOBE), Universiteitsplein 1C,  
26 2610 Antwerp, Belgium

27 7 Hasselt University, Interuniversity Institute for Biostatistics and Statistical Bioinformatics, Agoralaan  
28 Gebouw D, 3590 Diepenbeek, Belgium

29 8 ETH Zurich, Institute for Atmospheric and Climate Science, Universitaetstrasse 16, 8092 Zurich,  
30 Switzerland

31 9 University of Antwerp, Plants and ecosystems (PLECO), Universiteitsplein 1, 2610 Wilrijk, Belgium

32 10 Research Institute for Nature and Forest (INBO), Herman Teirlinckgebouw, Havenlaan 88 box 73,  
33 B-1000 Brussels, Belgium

34 11 University of Aberdeen, Institute of Biological and Environmental Sciences, 23 St Machar Drive,  
35 Aberdeen, AB24 3UU, UK

36 12 Namur University, Namur Institute of Complex Systems, and the Institute of Life, Earth, and  
37 Environment, Research Unit in Environmental and Evolutionary Biology, Rempart de la Vierge 8,  
38 5000 Namur , Belgium

39 13 University of Glasgow, Institute of Biodiversity, Animal Health and Comparative Medicine, Graham  
40 Kerr Building, G12 8QQ, UK.

41 14 University of Liege, Gembloux Agrobio Tech, TERRA, Res Ctr Atmosphere Ecosyst Exchanges, B-  
42 5030 Gembloux, Belgium

43 15 University of Liege, Gembloux Agrobio Tech, Department of Agricultural Economics and Rural  
44 Development, B-5030 Gembloux, Belgium

45 16 Hasselt University, Faculty of sciences, Agoralaan Gebouw D, 3590 Diepenbeek, Belgium

46 17 Forschungszentrum Juelich, Inst Bio & Geosci IBG-3: Agrosphere, Juelich, Germany

47 18 Ecotron Européen de Montpellier, Centre National de la Recherche Scientifique, 34980  
48 Montferrier-sur-Lez, France

49 19 Vrije Universiteit Brussel, Department of Hydrology and Hydraulic Engineering, Pleinlaan 2, 1050  
50 Brussels, Belgium

51 20 University of Bern, Climate and Environmental Physics, Sidlerstrasse 5, 3012 Bern, Switzerland

52 21 University of Bern, Oeschger Centre for Climate Change Research, Switzerland

53 22 Department of Plant Physiology, Faculty of Biology and Biotechnology, Maria Curie-Skłodowska  
54 University, Lublin, Poland

55

56 **Preface (100 words)**

57 While the general *direction* of ecosystems' responses to a variety of climate change scenarios has  
58 been well investigated, insights in the potential *amplitude* and *dynamics* of this response are scarce  
59 and the societal impacts often remain unquantified. Drawing on the expertise of researchers from a  
60 variety of disciplines, this paper outlines how methodological and technological advancements can  
61 help design climate experiments that better capture the dynamics and amplitude of ecosystem  
62 responses provoked by climate change and translate these responses into societal impacts.

63

64 **1. Introduction**

65 Climate change is expected to impact ecosystem communities and ecosystem functioning<sup>1</sup>. Crop  
66 yields<sup>2</sup>, carbon (C) sequestration in soil<sup>3</sup>, and pollination rate<sup>4</sup> are generally predicted to decrease,  
67 while land evapotranspiration<sup>5</sup> and tree mortality, especially in the Boreal region, are expected to  
68 increase<sup>6</sup>. At the same time, the redistribution of species will increase opportunities for pest and  
69 pathogen emergence<sup>1</sup>.

70 These functions are crucial for human well-being through their contribution to ecosystem services,  
71 and so impacting them will have important consequences for society<sup>7</sup>. However, refining the societal  
72 cost estimations remains a challenge, partly because large knowledge gaps regarding the amplitude  
73 and dynamics of these responses that make it difficult to plan for climate adaptation. Specifically  
74 designed climate change experiments are necessary to address these issues. The goal of this  
75 perspective paper is fourfold. First, while acknowledging the great advances achieved by climate  
76 change-ecosystem responses experiments so far, we also identify the challenges that many of them  
77 currently face: high complexity of climate change in terms of environmental variables, constraints in  
78 the number and amplitude of climate treatment levels, and the limited scope with regard to

79 responses and interactions covered (Section 2). Second, to overcome these challenges we propose  
80 an experimental design that can leverage the increased computational and technological capabilities  
81 to more accurately capture the complexity of climate change in experiments; increase the number  
82 and range of climate treatment levels, and employ an interdisciplinary approach to broaden the  
83 range of responses and interactions covered (Section 3). Third, we outline an experiment that applies  
84 these design recommendations to demonstrate how it can enhance our capacity to understand and  
85 predict ecosystem responses to climate change. We describe the technical infrastructure used in this  
86 experiment, the climate manipulations, and the analysis pathway all the way to the valuation of the  
87 changes in ecosystem services (Section 4). Fourth, we place this design within the larger context of  
88 climate change experiments and pinpoint its complementarity to other designs (Section 5).

89

## 90 2. Challenges of climate change experiments

### 91 *The complexity of climate change*

92 The first challenge for research on climate change-ecosystem responses lies in the complex manner  
93 in which global climate change will affect local weather. To mimic a future climate, factors such as air  
94 temperature, atmospheric CO<sub>2</sub>, and precipitation need to be manipulated in combination, which can  
95 be both conceptually and technologically challenging<sup>8</sup>. Therefore, a significant proportion of climate  
96 change experiments have focused on measuring the effects of specific combinations of climate  
97 factors (such as warming plus drought), manipulated using technology that was available or  
98 affordable at that time (such as passive night-time warming and rain exclusion curtains)<sup>9</sup>. Although  
99 these experiments have led to many invaluable outcomes, such approaches cannot fully cover the  
100 complexity of climate projections or the covariance of meteorological variables. As such, they may,  
101 for example, under- or overestimate the effects on ecosystem functioning of changes in the  
102 frequencies of frosts and heat waves, drought-heat-wave reinforcements<sup>10</sup>, interactions between soil  
103 moisture conditions and subsequent precipitation occurrence<sup>11</sup>, increased frequencies of mild  
104 droughts (including in spring and autumn), and increased frequency of heavy precipitation events<sup>12</sup>.

105 These climate alterations can have a strong influence on ecosystem functioning: for example,  
106 decreased frost frequency may have a significant impact on plant mortality<sup>13</sup> and more frequent mild  
107 droughts can trigger plant acclimation and hence resistance to drought stress<sup>14</sup>. Therefore, many  
108 climate change experiments did not simulate (i) an extreme event instead of a change in the mean  
109 for a given single factor, (ii) regimes of events instead of a single event for a given single factor, and  
110 (iii) complex coupling between multiple factors. This lack of refinement in climate manipulations  
111 likely compromised the reliability of the estimation of ecosystem responses. Some steps have already  
112 been taken to address this, by applying treatments of precipitation regime or heatwaves as observed  
113 in the field<sup>15,16</sup> and by using translocation experiments, where macrocosms are displaced across  
114 geographic gradients in order to expose them to other climates that match possible future conditions  
115 at the location of origin (space for time approach)<sup>17</sup>. However, such an issue cannot be solved by  
116 modelling alone, because it requires testing too many possible interactions between factors, as well  
117 as changing regimes of single factors.

118

119 *Climate treatment levels: number and range*

120 Because of the cost of specialized infrastructure, scientists are often limited in the number of  
121 experimental units they can set up within a given experiment. Hence, climate factors are often  
122 applied at only two levels: ambient and future projections<sup>9</sup>. This provides useful estimations on the  
123 direction of ecosystem responses but does not provide insights into the shape of the responses to  
124 these factors or how far away current conditions are from potential tipping points to alternative  
125 stable states<sup>18</sup>. Moreover, ecosystem responses to multifactor global change drivers are regulated by  
126 complex, nonlinear processes<sup>19</sup>, which makes modeling difficult with experimental data that comes  
127 only from the two-level manipulation of environmental factors<sup>20</sup>.

128 Also stemming from high equipment costs is the narrow range of climate treatments. Most  
129 experiments have kept this range within conservative boundaries<sup>21</sup>, presumably because more  
130 drastic (though realistic) climate treatments may have a catastrophic impact on a studied ecosystem,

131 potentially leading to the loss of expensively equipped replicates. The truncation of more extreme  
132 climate conditions has, in turn, led to a lack of evidence regarding their effects on ecosystem  
133 functioning.

134 Finally, low temporal resolution is also an issue. Because it requires an extensive and high frequency  
135 monitoring of ecosystem functions, a substantial proportion of climate change experiments have  
136 only measured the ecosystem dynamics or trajectories annually or seasonally. Such experiments may  
137 fail to detect short-term dynamics of ecosystem responses<sup>22</sup> or trajectories leading to a transition to  
138 an alternative stable state<sup>23,24</sup>. However, trends related to ecosystem dynamics often appear on  
139 decadal time scales, because of the time needed to alter biogeochemical cycles and the properties of  
140 soil organic matter. Therefore the duration of the monitoring should be prioritized over its frequency  
141 if the setup does not allow a good coverage of both.

142

#### 143 *Integration among disciplines*

144 The very nature of climate change and its impacts is discipline-spanning and therefore requires an  
145 integrated approach<sup>25</sup>. Although the number of interdisciplinary studies related to climate change is  
146 increasing steadily<sup>26</sup>, there are still many challenges related to interdisciplinary research. These  
147 include establishing common terminology, concepts and metrics<sup>25,27,28</sup>, a consistently lower funding  
148 success for interdisciplinary research projects<sup>29</sup>, and a general lack of interdisciplinary research  
149 positions<sup>25</sup>. The barriers depend largely on the purpose, forms and extent of knowledge integration,  
150 and their combination<sup>30</sup>. Although climate change research developed from multidisciplinary to  
151 interdisciplinarity, and further to transdisciplinarity<sup>31</sup>, most collaborative work in environmental  
152 research is small-scale rather than large-scale interdisciplinary work<sup>30</sup>. Small-scale integration refers  
153 to collaborations between similar partners (for example, different natural science disciplines), while  
154 large-scale integration crosses broader boundaries (such as between natural and social science)<sup>30</sup>.  
155 Currently, ecosystem services studies are mostly limited to either the natural science aspects or the  
156 socio-economic science aspects and rarely cover the entire ecosystem services cascade<sup>32</sup>. This lack of

157 large-scale knowledge integration results in errors along this cascade; both when moving from  
158 biodiversity and ecosystem functions to ecosystem services, and when moving from ecosystem  
159 services to societal values.

160

### 161 3. Recommendations

162 *Using climate model outputs and technology to refine climate change treatments*

163 A first option to prescribe a projected change in weather dynamics is to alter specific characteristics  
164 (such as drought duration, heat wave intensity) in isolation using high-frequency data of ambient  
165 weather conditions so that they match future projections. The advantage of this method is that  
166 atmospheric conditions can be modified with high-quality field data instead of relying upon less  
167 precise regional climate model outputs with lower spatial and temporal resolution. Moreover, if used  
168 to manipulate one climate factor at a time, such an approach facilitates a mechanistic understanding  
169 of ecosystem responses that can be further extrapolated through modeling. This design may  
170 combine two or more factors to provide information about interactions between climate  
171 parameters.

172 Incorporating the complexity of projected changes can also be achieved by using outputs of state-of-  
173 the-art climate models. Due to model biases, the appropriate model must be selected very carefully.  
174 Global climate models (GCMs) are useful tools for assessing climate variability and change on global  
175 to continental scales, typically with a spatial resolution of 100–250 km. To estimate climate variability  
176 at more local scales, GCMs are dynamically downscaled using regional climate models (RCMs), which  
177 resolve the climate at higher resolutions (typically 10–50 km). The GCM/RCM combinations can then  
178 be chosen based on (i) how well models perform against local climate and weather characteristics in  
179 the studied ecosystem and (ii) how representative future projections are to the multi-model mean. In  
180 this case, one can simulate an ecosystem response to a given climate setup with higher accuracy.  
181 However, unlike with a full factorial experiment, it is not possible to attribute an ecosystem response  
182 to a given climate factor. Nevertheless, the model-output approach does facilitate the application of



183 increasingly high warming levels by using a global mean temperature gradient (see Section 4). It also  
184 addresses the issues of covarying variables, and it can be directly linked with a scenario from the  
185 Intergovernmental Panel on Climate Change which would represent a major step towards bridging  
186 the gap between climate and ecosystem science.

187

188 However, to implement these options it is necessary to control climate conditions and atmospheric  
189 composition with high frequency and high accuracy. This can be achieved only with dedicated and  
190 advanced equipment. Ecotron infrastructures, which consist of a set of replicated experimental units  
191 where environmental conditions are tightly controlled and where multiple ecosystem processes are  
192 automatically monitored, are well-suited to fulfill these needs<sup>33</sup>. Such infrastructures have been  
193 historically limited to a handful across the world<sup>9</sup>, but are becoming increasingly widespread<sup>34–36</sup>.  
194 They also offer the opportunity to monitor ecosystem responses at sub-hourly frequencies, making it  
195 possible to simultaneously discriminate between short- and long-term ecosystem responses.

196

197 *Increasing the number and range of climate treatment levels*

198 A gradient design, in which one or several climate factors are applied at increasingly high levels, can  
199 substantially increase the resolution of a climate change experiment. This is better suited to  
200 quantitatively describing the relationship between a response variable and a continuous climate  
201 factor than the more traditional approach of testing ambient versus a single future projection, and  
202 allows the collection of quantitative data for ecological models<sup>37</sup>. It also makes it possible to detect  
203 nonlinearity, thresholds, and tipping points, and to interpolate and extrapolate ecosystem  
204 responses<sup>18</sup>. While such gradient designs should ideally be replicated, unreplicated regression  
205 designs can be a statistically powerful way of detecting response patterns to continuous and  
206 interacting environmental drivers, provided that the number of levels in the gradient is large  
207 enough<sup>37</sup>.

208 To ensure appraisal of the largest possible range of ecosystem responses, the gradient should be as  
209 long as possible, even extending beyond the most extreme conditions. Broader treatment modalities  
210 can also inform how far a specific ecosystem response is situated relative to its upper or lower  
211 tolerance limit. In addition, the levels of the gradient may be spread in a non-linear manner to  
212 achieve the highest resolution in the range where the strongest ecosystem responses are expected.

213

214 *Employing an interdisciplinary approach to better capture responses and interactions*

215 We argue that an overarching objective of climate change experiments is to contribute to the  
216 understanding of the impacts that climate change has on nature and society as well as to enlarge our  
217 potential for climate adaptation. However, as outlined in Section 2, the lack of large-scale knowledge  
218 integration can result in errors along the ecosystem services cascade; first in the step from  
219 biodiversity and ecosystem functions to ecosystem services and second from ecosystem services to  
220 societal values.

221 Regarding the first step, thorough quantification of ecosystem services should be based on specific  
222 data regarding how the ecosystem is functioning. Many ecosystem service studies use land use as an  
223 indicator of ecosystem service delivery<sup>32</sup>, but often land use classification cannot capture differences  
224 between abiotic conditions and ecological processes that explain differences in service delivery<sup>38</sup>.  
225 Therefore, using land use as a simple indicator will result in inappropriate management decisions<sup>38</sup>.

226 Regarding the second step, economists need to be involved early in the process. Although there are  
227 many ways in which ecosystem function changes can affect the provision of ecosystem services to  
228 society<sup>39</sup>. However, budget constraints necessitate the selection of those ecosystem functions and  
229 services that are considered most important to society. A common selection approach is to consider  
230 the potential impact of ecosystem changes in terms of human welfare endpoints, often by means of  
231 monetary valuation. Ecologists and economists must interact across disciplinary boundaries if  
232 ecological experiments are intended to predict these endpoints within an ecosystem services  
233 context<sup>40</sup>. Hence, economists need to be involved during the design of ecological experiments in

234 order to ensure that those ecosystem service changes that are most relevant for human welfare are  
235 measured and predicted.

236 We suggest that, the desired large-scale integration can be achieved in several steps, organized in a  
237 top-down approach. The first step is to identify the key ecosystem services to value based on welfare  
238 endpoints<sup>41</sup>. For most terrestrial ecosystems, this would imply assessing services from the following  
239 list: food and raw material production and quality, water supply and quality, C sequestration,  
240 depollution, erosion prevention, soil fertility, pest and pathogen control, pollination, maintenance of  
241 biodiversity and recreation. The second step consists of identifying the set of variables that best  
242 describes the ecosystem functions, processes and structures associated with these services. Based on  
243 the literature<sup>42</sup>, we suggest the following measures (see also Table 1): (i) vegetation variables (plant  
244 community structure, above/belowground biomass, litter quality), (ii) atmospheric parameters (net  
245 ecosystem exchange, greenhouse gas emissions), (iii) soil abiotic (pH, texture, electrical conductivity,  
246 macro-, micronutrient and pollutant content) and biotic (fauna and microbial community structure,  
247 respiration, and biomass) variables, and (iv) all parameters that describe movements of water in the  
248 soil-plant-atmosphere continuum (precipitation, leaching, air relative humidity, evapotranspiration,  
249 water potential). Air and soil temperatures should also be monitored, since they determine  
250 biogeochemical reaction rates. Finally, ecosystem processes, structures and functions need to be  
251 translated into services, and ultimately into societal value by expressing them in monetary and non-  
252 monetary terms. Measuring all of these variables, integrating them in an ecosystem service  
253 framework, and estimating the societal value of these services would require expertise from plant  
254 ecologists and ecophysiologicals, hydrologists, soil biogeochemists, animal ecologists, microbiologists,  
255 pedologists, climatologists, as well as modelers and environmental economists<sup>43</sup>.

256

#### 257 **4. An initial application of the recommendations: The UHasselt Ecotron experiment**

258 Here we describe our proposed interdisciplinary approach in the context of a climate change  
259 manipulation using the UHasselt Ecotron experiment.

260


### 261 *Ecotron infrastructure*

262 The UHasselt Ecotron facility consists of tightly controlled climate change manipulations of 12  
263 macrocosms (soil-canopy columns of 2 m in diameter and 1.5 m depth), extracted without significant  
264 disruption of the soil structure from a dry heathland plot in the 'Hoge Kempen' National Park (50° 59'  
265 02.1" N, 5° 37' 40.0" E) in November 2016. The plot was managed for restoration six years before the  
266 sampling. The design of this infrastructure benefited from exchanges through the AnaEE (Analysis  
267 and Experimentation on Ecosystems)/ESFRI (European Strategy Forum on Research Infrastructure)  
268 project. Some of the infrastructure's features were inspired by the Macrocosms platform of the CNRS  
269 Montpellier Ecotron<sup>16</sup>. Each UHasselt Ecotron unit consists of three compartments: the dome, the  
270 lysimeter, and the chamber. The dome consists of a shell-shaped dome made of highly PAR  
271 (photosynthetically active radiation) transparent material, where wind and precipitation are  
272 generated and measured and where the concentration of greenhouse gases (CO<sub>2</sub>, N<sub>2</sub>O, CH<sub>4</sub>), PPF  
273 (photosynthetic photon flux density) and difference between incoming and outgoing short- and long-  
274 wave radiation are measured. The lysimeter (equipment for measuring hydrological variations  
275 undergone by a body of soil under controlled conditions) contains the soil-canopy column, where  
276 soil-related parameters are controlled (including the vertical gradient of soil temperature and water  
277 tension) and measured, and is weighed every minute. Suction cups and soil sensors are installed  
278 following a triplicated 5 depth design (Fig. S1). The chamber is a gastight room that encloses the  
279 lysimeter, where air pressure, air temperature, relative humidity, and CO<sub>2</sub> concentration are  
280 controlled and key variables measured in each unit (Fig. S1). The UHasselt Ecotron is linked with a  
281 nearby Integrated Carbon Observation System (ICOS) ecosystem tower ([https://www.icos-](https://www.icos-ri.eu/home)  
282 [ri.eu/home](https://www.icos-ri.eu/home)), which provides real-time data on local weather and soil conditions, with a frequency of  
283 at least 30 minutes.

284

### 285 *Climate manipulations*

286 A double-gradient approach is adopted: one approach (six units) measures the effect of an altered  
 287 single factor (here, precipitation regime), while maintaining the natural variation of other abiotic  
 288 factors, and the other approach (six units) manipulates climate by jointly simulating all covarying  
 289 parameters, representing increasingly intense climate change. The two approaches are described  
 290 below. Because they sit isolated in an enclosed facility, it is possible that small initial differences in  
 291 the soil-canopy core in a given unit will increase with time to the point where it becomes statistically  
 292 different from the others. Therefore, the units were first distributed within the two gradients using a  
 293 cluster analysis to minimize the noise in ecosystem responses measured during a test period (see Fig.  
 294 S2) due to small-scale soil heterogeneity. This clustering was used to distribute the units according to  
 295 the pattern shown in Fig. 1.



296 *Figure 1. Overview of the two climate change gradient designs in the UHasselt Ecotron experiment.*  
 297

298 *The units have been redistributed to maximize statistical similarity within a gradient prior to the*


299 *treatment. Global mean temperature anomalies are computed with respect to the reference period*  
300 *1951-1955.*

301

302 Climate change projections for the NW Europe region predict higher probability of both heavier  
303 precipitation and longer droughts, without a significant change in yearly precipitation<sup>44</sup>. The  
304 precipitation regime gradient uses real-time input from the ecosystem tower nearby, and only alters  
305 precipitation events: across the gradient, increasingly long periods (2, 6, 11, 23, 45 and 90 days),  
306 based on local climate records from Maastricht, NL<sup>45</sup>) in which precipitation is withheld (dry period)  
307 are followed by increasingly long periods in which precipitation is increased (wet period), with the  
308 duration of the two periods kept equal within a unit (Fig. 1). Precipitation events during the wet  
309 period are increased twofold and are adjusted at the end of the period to avoid altering the yearly  
310 precipitation amount.

311 To drive the second gradient of the UHasselt Ecotron experiment, we use the climate variables  
312 produced by an RCM following Representative Concentration Pathway (RCP) 8.5, a high-emission  
313 scenario<sup>46</sup>. The gradient itself is determined based on global mean temperature anomalies. In the six  
314 units, climates corresponding to a +0 ° to +4 °C warmer world (projected for periods ranging from  
315 1951–1955 to 2080–2089) are simulated (Fig. 1, Fig. S3), by extracting local climate conditions from  
316 the RCM for periods consistent with these warming levels (Fig. S3)<sup>47</sup>. This set-up also facilitates  
317 comparison of the ‘present-day’ climate as simulated by the RCM (the +1 °C unit), to the unit driven  
318 by ICOS field observations. Moreover, the climate simulated in the +1.5° C unit is reasonably  
319 consistent with the lower end of the long-term temperature goals set by the Paris Agreement<sup>48</sup>.

320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333



334 Figure 2. Impact pathway showing the reasoning behind the integration of scientific disciplines in the  
335 UHasselt Ecotron experiment. The research hypotheses are given in italics and described in more  
336 detail in Fig. S4.

337

### 338 Integrating scientific disciplines for an interdisciplinary ecosystem service approach

339 As outlined in Section 3, climate change experiments require large-scale knowledge integration to  
340 enable more useful estimates of climate change effects on ecosystem functioning and on society. The  
341 UHasselt Ecotron facility makes it possible to extend the degree of interdisciplinarity by investigating  
342 the entire cascade from climate changes to ecosystem functions, ecosystem services, and, finally,  
343 societal values. As such, the ecotron facility contributes to the development towards large-scale  
344 knowledge integration on climate change. Consequently, the UHasselt Ecotron experiment brings  
345 together several disciplines in an interdisciplinary framework (Fig 2). With input from other involved

346 disciplines, climatologists design the protocols for climate manipulations and plant ecologists  
347 monitor plant communities in each ecotron unit. Numerical models for water movement within one  
348 unit are developed by mathematicians and hydrologists. Ecotron output on C cycling is fed into a soil  
349 C model<sup>49</sup>, both for calibration and prediction purposes. Community modelers improve the power of  
350 this model by accounting for the soil community structure and species interactions (food web). The  
351 specific role of soil organisms in soil biogeochemistry is investigated by microbial and soil fauna  
352 ecologists. This is inferred from variation in responses of different functional groups such as nitrogen  
353 fixers, mycorrhizal fungi and different feeding guilds of soil fauna, combined with additional separate  
354 experiments, both in the field and *in vitro*. The outputs of the measurements above (see Table 1)  
355 allow experts in ecosystem ecology to quantify ecosystem services. Environmental economists  
356 express the change in ecosystem services provided using best-practice monetization approaches<sup>50</sup>.  
357 For example, water quality regulation is assessed as the prevented cost of intensified water  
358 treatment or use of other water resources. Measurements of vegetation, soil abiotic parameters and  
359 the water balance make it possible to quantify this benefit. Carbon sequestration is assessed as the  
360 prevented cost from increased global temperature, which can be quantified based on vegetation, air  
361 parameters and soil abiotic parameters measurements. Maintenance of biodiversity and recreation  
362 can be assessed based on measurements of vegetation.

363 We note that (monetary) estimates from an individual study can often not be applied directly for  
364 generating policy-recommendations<sup>51</sup>, especially for complex and spatially heterogeneous problems  
365 such as climate change impacts on ecosystems. However, meta-analyses need to rely on data  
366 generated by primary studies that estimate the societal cost (or benefit) of changes in specific  
367 services provided by a specific ecosystem at specific location(s). In this regard, the UHasselt Ecotron  
368 experiment can also provide valuable input data for dedicated policy-guiding analyses<sup>52</sup>.

369



370 Table 1. Measured variables in the UHasselt Ecotron experiment and links with ecosystem functions,  
 371 services, and values. Left-hand side of the table: ecosystem services. Right-hand side: variables  
 372 measured in the ecotron experiment. Lower part of the table: illustration of how the societal value of  
 373 four of the ecosystems services will be assessed.

ECOSYSTEM SERVICES										MEASURED VARIABLES				
Food	Raw materials	Water quality	C sequestration	Erosion prevention	Maintenance of biodiversity	Recreation	Climate regulation	Water retention	Soil fertility	Depollution	Pathogen control	Variable category	Variable	Frequency of measurement
o	o	o	o	o	o	o						Vegetation	Plant community structure	6 months
o	o	o	o	o	o	o							Shoot & root biomass	6 months
		o					o					Air parameters	Net ecosystem exchange (NEE)	30 min
													Temperature	2 min
													GHG emissions (CH4, N2O)	2 min
			o					o	o			Soil abiotic parameters	Texture	1 year
								o	o		Temperature		2 min	
		o								o	Biochemical composition		1 year	
		o								o	Electrical conductivity		30 min	
		o								o	Soil pore water chemistry		2 weeks	
										o	Available pollutant concentration		1 year	
										o	o	Soil biotic parameters	Fauna community structure	6 months
			o							o	o		Microbial community structure	6 months
										o			Mineralization rate	1 year
	o		o					o				Water balance	Precipitation	30 min
	o							o	o	o	Leaching		30 min	
	o									o	Relative humidity		30 min	
	o									o	Evapotranspiration		30 min	
	o									o	Soil water potential		30 min	
ECONOMIC VALUATION														
												Prevented cost of intensified water treatment or use of other water resources		
												Prevented damage cost from increased global temperature		
												Non-use value of continued existence of biodiversity		
												Use value of recreational enjoyment		

374

375 **5. The place of the suggested design within the landscape of climate change experiments**

376 A comprehensive understanding of ecosystem responses to climate change can only be achieved  
 377 through the use of a broad range of different, complementary experimental designs, all of which can  
 378 be integrated through modeling. The experimental design suggested here exhibits a unique set of  
 379 advantages and drawbacks, which makes it suited to tackle specific needs within the climate change  
 380 experiments landscape.

381 **Strengths and limitations of the design**

382 The strengths of the suggested design comprises (1) high-performance microclimate conditioning,  
 383 both above- and belowground, which makes it possible to approximate field conditions while  
 384 maintaining control, (2) high-frequency automated measurements of ecosystem functions and thus

385 of treatment impact thereon, and (3) a large-scale interdisciplinary approach. The first two strengths  
386 are inherent to the ecotron research infrastructure, while the large-scale integration can  
387 theoretically be implemented in any climate change experiment. However, we consider ecotron  
388 infrastructures to be particularly suitable for such an interdisciplinary approach, because of the high-  
389 end climate control and the broad range of functions monitored at a high frequency.

390 With respect to (1), studies focusing on ecosystem functions, processes and structures that are highly  
391 sensitive to soil temperature and soil water potential would benefit most from being conducted in  
392 ecotrons (for example, soil CO<sub>2</sub> exchange and C sequestration, growth and activity of soil microbes  
393 and soil fauna), as the lysimeter component can generate very precise lower boundary conditions  
394 and thus realistic vertical soil profiles of temperature and soil water status. With respect to (2),  
395 studies in which the high-resolution temporal pattern of ecosystem functions and their coupling is  
396 important would also benefit from ecotron infrastructures, as it is difficult to measure these  
397 parameters manually across long time scales. For example, simultaneous automated measurement  
398 of the carbon, water and mineral nutrient cycles makes it possible to disentangle their interactions in  
399 a range of climate conditions, and to feed control mechanisms into models.

400 A first set of constraints in the usefulness of the experimental design described in this paper stems  
401 from the scale limitation of the experimental units. Ecotrons can accommodate plants only of small  
402 stature (less than two meters in height), which excludes forests and tall crops. For the same reason,  
403 the impact of megafauna such as grazers or top predators cannot be tested. Results obtained in  
404 macrocosms only integrate small-scale (less than one meter) variability, which leads to a lack of  
405 accuracy when scaling up to ecosystem.

406 Second, it may be difficult to financially support this type of experiment on the time scale of  
407 ecosystem responses (10 years or more)<sup>53</sup>. Ecosystem shifts to alternative stable states may remain  
408 undetected if the funding period is shorter than the period required for the ecosystem to shift. A  
409 partial solution for this would be to adopt a gradient design with increasingly late endpoints of

410 projected climate change; this would allow for some extrapolation of ecosystem response in time  
411 (trajectories), which is possibly enough to estimate ranges of this response in the longer term.

412 Third, macrocosms in ecotron facilities are isolated from their ecosystem of origin. Hence genetic  
413 input from propagules or pollination probably differ significantly from the field, which can be an  
414 issue, especially in long-term experiments. This could be mitigated in two ways. The first is by  
415 minimizing sampling disturbance, by sampling for soil microbes and soil fauna not more than twice a  
416 year, using 10 cm diameter soil cores, this would account for only 1.5% of total soil surface annually.  
417 The second way is by replacing soil sampling cores in the lysimeter by cores taken from the same  
418 ecosystem. This would also avoid holes at the soil surface that may alter water flow through the soil  
419 column. Furthermore, field traps to collect airborne propagules can be collected yearly and their  
420 content spread on the enclosed surface of the soil-canopy columns. These solutions would at least  
421 ensure fresh genetic input into the system, even though this input may be different in the field in  
422 future conditions.

423 Finally, radiation in ecotron enclosures sometimes differ than in the field. Artificial LED-lightning  
424 allows to control radiation precisely but is yet not able to reach the same radiation level as in the  
425 field, while ambient lightning can disrupt its synchronization with temperature or precipitation. This  
426 may be an issue while simulating heatwaves and droughts, which have more sunshine hours than  
427 wet periods<sup>54</sup>.

428

#### 429 *Complementarity with other climate change experiments*

430 The weaknesses of the proposed design (small spatial scale, potentially insufficient time-scale, lack of  
431 interaction with the surrounding environment) can be mitigated further through the use of  
432 complementary experiments, which might even be partially integrated into the overarching  
433 approach. For example, owing to small spatial scale, the results might have limited validity as a  
434 predictor of ecosystem responses at other sites and in other habitats. Running experiments in  
435 parallel across multiple climates and locations with the same methodology, also known as

436 “coordinated distributed experiments” (CDEs), would be better suited for this purpose as it allows  
437 extrapolation and generalization of results while correcting for effect size<sup>55</sup>. For example, such a  
438 design makes it possible to study plant response to nutrient addition and herbivore exclusion<sup>56</sup>; and  
439 ecological responses to global change factors across 20 eco-climate domains using a set of  
440 observatory sites<sup>57</sup>. In fact, a coordinated distributed experiment using the design presented in  
441 Section 4, and testing the same climate gradient in different ecosystems across several ecotron  
442 facilities would combine the high generalization potential of CDEs with the precision of ecotrons.  
443 A second area for potential complementarity and integration is translocation experiments. These  
444 experiments are well suited for long-term observations due to their relatively low funding  
445 requirements and relative ease of implementation, and the soil macrocosms used in these  
446 experiments are still connected to their surrounding environment<sup>17</sup>. However, the functioning of the  
447 ecosystem is monitored less comprehensively and frequently within these types of experiments and  
448 the influence of different climate factors on ecosystem functioning cannot be disentangled.  
449 Consequently, running an ecotron and a translocation experiment in parallel on the same ecosystem  
450 with similar climate treatments would make it possible to estimate the effect size of the connection  
451 with the surrounding environment on ecosystem response to climate change. This information can  
452 then, in turn, be used to correct the outputs of future ecotron experiments by accounting for the  
453 isolation factor.

454

#### 455 *Usefulness of the suggested design for modeling ecosystem response to climate change*

456 While ecosystem models can be evaluated and calibrated using a range of data sources, including  
457 sites in different climate zones and long-term experiments without climate manipulation<sup>58</sup>, data from  
458 well-controlled, replicated and highly instrumented facilities such as those described here are  
459 invaluable for testing the process understanding encapsulated in the models, and for testing model  
460 behavior against detailed, multi-parameter observations<sup>36</sup>. Models that are tested and, where  
461 necessary, calibrated against such data can then be evaluated against data from other sites. If the

462 outputs do not prove to be generalizable, the information derived from testing the model could be  
463 used to refine the experimental design and explain variation in the measured values. If the outputs  
464 prove generalizable, the models can be used across larger temporal and spatial scales to project  
465 potential impacts of future climate change<sup>59,60</sup>.

466

## 467 **6. Conclusion**

468 The effects of climate change on ecosystem functioning have far-reaching consequences for society.  
469 Here we present a type of experiment that is designed to estimate the amplitude and dynamics of  
470 ecosystem responses to climate change, and the consequences for ecosystem services. We have  
471 outlined that climate change experiments are facing three types of challenges: limitations in  
472 addressing the complexity of climate change in terms of control of environmental variables,  
473 constraints in the number and range of climate level treatments, and restrictions in scope. We have  
474 suggested ways to address these challenges: improving computational and technological capabilities,  
475 increasing the number and range of climate treatment levels, and employing an interdisciplinary  
476 approach. We illustrated these suggestions through a case study where they have been  
477 implemented, and outlined the place of this design in the broader landscape of climate change  
478 experiments.

479 We foresee that the holistic approach outlined in this perspective could yield more reliable,  
480 quantitative predictions of terrestrial ecosystem response to climate change, and could improve  
481 knowledge on the value of ecosystem services and their links with ecosystem processes. We expect  
482 these results to be of interest for society beyond just scientists: they provide nature managers with  
483 predictions on ecosystem responses to help them decide on ecosystem management practices in the  
484 mid- and long-term, and that they will explain to policymakers and the wider public the societal  
485 impact of ecosystem changes induced by climate change at a more detailed, ecosystem-specific level.

486

## 487 **Acknowledgments**

488 The authors thank the Flemish government (through Hercules Stichting big infrastructure and the  
489 Fund for Scientific Research Flanders project G0H4117N), LSM (Limburg Sterk Merk, project 271) for  
490 providing funds to build the UHasselt Ecotron; Hasselt University for both funding and policy support  
491 (project BOF12BR01 and Methusalem project 08M03VGRJ); and the ecotron research committee for  
492 useful comments on the experimental design. We also thank RLKM (Regional Landscape Kempen and  
493 Maasland) for its collaboration and support. Nele Witters, Sebastien Lizin, Anne Nobel, and Inne  
494 Vanderkelen are funded by Research Foundation-Flanders (FWO).

495

#### 496 **Authors' contributions**

497 FR and RM took the lead in writing the manuscript and received input from all co-authors. The initial  
498 conceptualization of this manuscript was discussed during a consortium meeting. All authors  
499 proofread and provided their input to different draft versions and gave their final approval for  
500 submission.

501

#### 502 **References**

- 503 1. Scheffers, B. R. *et al.* The broad footprint of climate change from genes to biomes to people.  
504 *Science (80-. ).* **354**, (2016).
- 505 2. Zhao, C. *et al.* Temperature increase reduces global yields of major crops in four independent  
506 estimates. 1–6 (2017). doi:10.1073/pnas.1701762114
- 507 3. Allen, C. D. *et al.* A global overview of drought and heat-induced tree mortality reveals  
508 emerging climate change risks for forests. *For. Ecol. Manage.* **259**, 660–684 (2010).
- 509 4. Hat, J. L. & Prueger, J. H. Temperature extremes : Effect on plant growth and development.  
510 **10**, 4–10 (2015).
- 511 5. Collins, M. *et al.* Long-term Climate Change: Projections, Commitments and Irreversibility. in  
512 *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth*  
513 *Assessment Report of the Intergovernmental Panel on Climate Change* (eds. Stocker, T. F. et

- 514 al.) 1029–1136 (Cambridge University Press, Cambridge, United Kingdom and New York, NY,  
515 USA, 2013). doi:10.1017/CBO9781107415324.024
- 516 6. Pecl, G. T. *et al.* Biodiversity redistribution under climate change: Impacts on ecosystems and  
517 human well-being. **9214**, (2017).
- 518 7. Millenium Ecosystem Assessment. *Ecosystems and human well-being: Synthesis*. Island Press,  
519 Washington, DC. (2005). doi:10.1196/annals.1439.003
- 520 8. Leuzinger, S. *et al.* Do global change experiments overestimate impacts on terrestrial  
521 ecosystems? *Trends Ecol. Evol.* **26**, 236–241 (2011).
- 522 9. Stewart, R. I. A. *et al.* Mesocosm Experiments as a Tool for Ecological Climate-Change  
523 Research. *Advances in Ecological Research* **48**, (Elsevier Ltd., 2013).
- 524 10. Zscheischler, J. & Seneviratne, S. I. Dependence of drivers affects risks associated with  
525 compound events. *Sci. Adv.* **3**, 1–11 (2017).
- 526 11. Guillod, B. P., Orlowsky, B., Miralles, D. G., Teuling, A. J. & Seneviratne, S. I. Reconciling spatial  
527 and temporal soil moisture effects on afternoon rainfall. *Nat. Commun.* **6**, 1–6 (2015).
- 528 12. Thiery, W. *et al.* Hazardous thunderstorm intensification over Lake Victoria. *Nat. Commun.* **7**,  
529 1–7 (2016).
- 530 13. Berendse, F., Schmitz, M. & Visser, W. De. Experimental Manipulation of Succession in  
531 Heathland Ecosystems. *Oecologia* **100**, 38–44 (1994).
- 532 14. Backhaus, S. *et al.* Recurrent Mild Drought Events Increase Resistance Toward Extreme  
533 Drought Stress. *Ecosystems* **17**, 1068–1081 (2014).
- 534 15. Verburg, P. S. J. *et al.* Impacts of an anomalously warm year on soil nitrogen availability in  
535 experimentally manipulated intact tallgrass prairie ecosystems. *Glob. Chang. Biol.* **15**, 888–  
536 900 (2009).
- 537 16. Roy, J. *et al.* Elevated CO<sub>2</sub> maintains grassland net carbon uptake under a future heat and  
538 drought extreme. *Proc. Natl. Acad. Sci.* **113**, 6224–6229 (2016).
- 539 17. Cantarel, A. M. & Bloor, J. M. G. Four years of simulated climate change reduces above-

- 540 ground productivity and alters functional diversity in a grassland ecosystem. **24**, 113–126  
541 (2013).
- 542 18. Kreyling, J. *et al.* To replicate, or not to replicate - that is the question: how to tackle nonlinear  
543 responses in ecological experiments. *Ecol. Lett.* (2018). doi:10.1111/ele.13134
- 544 19. Zhou, X., Weng, E. & Luo, Y. Modeling patterns of nonlinearity in ecosystem responses to  
545 temperature, Co<sub>2</sub>, and precipitation changes. *Ecol. Appl.* **18**, 453–466 (2008).
- 546 20. Luo, Y. *et al.* Modeled interactive effects of precipitation, temperature, and [CO<sub>2</sub>] on  
547 ecosystem carbon and water dynamics in different climatic zones. *Glob. Chang. Biol.* **14**,  
548 1986–1999 (2008).
- 549 21. Kayler, Z. E. *et al.* Experiments to confront the environmental extremes of climate change.  
550 (2015). doi:10.1890/140174
- 551 22. Svenning, J. C. & Sandel, B. Disequilibrium vegetation dynamics under future climate change.  
552 *Am. J. Bot.* **100**, 1266–1286 (2013).
- 553 23. Harris, R. M. B. *et al.* Biological responses to the press and pulse of climate trends and  
554 extreme events. *Nat. Clim. Chang.* **8**, 579–587 (2018).
- 555 24. Scheffer, M., Carpenter, S., Foley, J. a, Folke, C. & Walker, B. Catastrophic shifts in ecosystems.  
556 *Nature* **413**, 591–6 (2001).
- 557 25. Hein, C. J. *et al.* Overcoming early career barriers to interdisciplinary climate change research.  
558 *Wiley Interdiscip. Rev. Clim. Chang.* **9**, 1–18 (2018).
- 559 26. Xu, X., Goswami, S., Gullette, J., Wullschleger, S. D. & Thornton, P. E. Interdisciplinary  
560 research in climate and energy sciences. *Wiley Interdiscip. Rev. Energy Environ.* **5**, 49–56  
561 (2016).
- 562 27. Sievanen, L., Campbell, L. M. & Leslie, H. M. Challenges to Interdisciplinary Research in  
563 Ecosystem-Based Management. *Conserv. Biol.* **26**, 315–323 (2012).
- 564 28. Abiven, S. *et al.* Integrative research efforts at the boundary of biodiversity and global change  
565 research. *Curr. Opin. Environ. Sustain.* **29**, 215–222 (2017).



- 566 29. Bromham, L., Dinnage, R. & Hua, X. Interdisciplinary research has consistently lower funding  
567 success. *Nature* **534**, 684–687 (2016).
- 568 30. Turner, L. M. *et al.* Transporting ideas between marine and social sciences: experiences from  
569 interdisciplinary research programs. *Elem Sci Anth* **5**, (2017).
- 570 31. Hellsten, I. & Leydesdorff, L. The construction of interdisciplinarity: the development of the  
571 knowledge base and programmatic focus of the journal *Climatic Change*, 1977-2013. *J. Assoc.*  
572 *Inf. Sci. technology* **67**, 2181–2193 (2016).
- 573 32. Boerema, A., Rebelo, A. J., Bodi, M. B., Esler, K. J. & Meire, P. Are ecosystem services  
574 adequately quantified? *J. Appl. Ecol.* **54**, 358–370 (2017).
- 575 33. Clobert, J. *et al.* How to Integrate Experimental Research Approaches in Ecological and  
576 Environmental Studies : AnaEE France as an Example. **6**, (2018).
- 577 34. Mougin, C. *et al.* A coordinated set of ecosystem research platforms open to international  
578 research in ecotoxicology, AnaEE-France. *Environ. Sci. Pollut. Res.* **22**, 16215–16228 (2015).
- 579 35. Eisenhauer, N. & Türke, M. From climate chambers to biodiversity chambers. *Front. Ecol.*  
580 *Environ.* **16**, 136–137 (2018).
- 581 36. Milcu, A. *et al.* Functional diversity of leaf nitrogen concentrations drives grassland carbon  
582 fluxes. *Ecol. Lett.* **17**, 435–444 (2014).
- 583 37. Cottingham, K. L., Lennon, J. T. & Brown, B. L. Knowing when to draw the line : designing more  
584 informative ecological experiments. *Front. Ecol. Environ.* **3**, 145–152 (2005).
- 585 38. Van der Biest, K. *et al.* Evaluation of the accuracy of land-use based ecosystem service  
586 assessments for different thematic resolutions. *J. Environ. Manage.* **156**, 41–51 (2015).
- 587 39. Polasky, S. & Segerson, K. Integrating Ecology and Economics in the Study of Ecosystem  
588 Services: Some Lessons Learned. *Annu. Rev. Resour. Econ.* **1**, 409–434 (2009).
- 589 40. Inkpen, S. A. & Desroches, C. T. When Ecology Needs Economics and Economics Needs  
590 Ecology: Interdisciplinary Exchange in the Age of Humans. **21** (2019).
- 591 41. Costanza, R. *et al.* Changes in the global value of ecosystem services. *Glob. Environ. Chang.* **26**,

- 592 152–158 (2014).
- 593 42. Braat, L. C. & de Groot, R. The ecosystem services agenda: bridging the worlds of natural  
594 science and economics, conservation and development, and public and private policy. *Ecosyst.*  
595 *Serv.* **1**, 4–15 (2012).
- 596 43. Plaas, E. *et al.* Towards valuation of biodiversity in agricultural soils: A case for earthworms.  
597 *Ecol. Econ.* **159**, 291–300 (2019).
- 598 44. Brouwers, J. *et al.* *MIRA Climate Report 2015, about observed and future climate changes in*  
599 *Flanders and Belgium.* (2015). doi:10.13140/RG.2.1.2055.8809
- 600 45. Klein Tank, A. M. G. *et al.* Daily dataset of 20th-century surface air temperature and  
601 precipitation series for the European Climate Assessment. *Int. J. Climatol.* **22**, 1441–1453  
602 (2002).
- 603 46. van Vuuren, D. P. *et al.* The representative concentration pathways: An overview. *Clim.*  
604 *Change* **109**, 5–31 (2011).
- 605 47. Seneviratne, S. I., Donat, M. G., Pitman, A. J., Knutti, R. & Wilby, R. L. Allowable CO<sub>2</sub> emissions  
606 based on regional and impact-related climate targets. *Nature* **529**, 477–483 (2016).
- 607 48. UNFCCC. Conference of the Parties (COP). Paris Climate Change Conference–November 2015,  
608 COP 21. *Adopt. Paris Agreement. Propos. by Pres.* **21932**, 32 (2015).
- 609 49. Smith, J. *et al.* Estimating changes in Scottish soil carbon stocks using ECOSSE. II. Application.  
610 *Clim. Res.* **45**, 193–205 (2010).
- 611 50. Schaubroeck, T. *et al.* Environmental impact assessment and monetary ecosystem service  
612 valuation of an ecosystem under different future environmental change and management  
613 scenarios ; a case study of a Scots pine forest. *J. Environ. Manage.* **173**, 79–94 (2016).
- 614 51. Hunter, J. E. & Schmidt, F. L. Cumulative research knowledge and social policy formulation:  
615 The Critical Role of Meta-Analysis. *Psychol. Public Policy, Law* **2**, 324–347 (1996).
- 616 52. Gerstner, K. *et al.* Will your paper be used in a meta-analysis? Make the reach of your  
617 research broader and longer lasting. *Methods Ecol. Evol.* **8**, 777–784 (2017).

- 618 53. Knapp, A. K. *et al.* Past , Present , and Future Roles of Long-Term Experiments in the LTER  
619 Network. **62**, 377–389 (2012).
- 620 54. De Boeck, H. ., Dreesen, F. E., Janssens, I. A. & Nijs, I. Climatic characteristics of heat waves  
621 and their simulation in plant experiments. *Glob. Chang. Biol.* **16**, 1992–2000 (2010).
- 622 55. Fraser, L. H. *et al.* Coordinated distributed experiments : an emerging tool for testing global  
623 hypotheses in ecology and environmental science. (2013). doi:10.1890/110279
- 624 56. Lind, E. M. *et al.* Life-history constraints in grassland plant species : a growth-defence trade-  
625 off is the norm. 513–521 (2013). doi:10.1111/ele.12078
- 626 57. Keller, M., Schimel, D. S., Hargrove, W. W. & Hoffman, F. M. A continental strategy for the  
627 National Ecological Observatory Network. *Front. Ecol. Enviornment* **6**, 282–284 (2008).
- 628 58. Smith, P. *et al.* Towards an integrated global framework to assess the impacts of land use and  
629 management change on soil carbon : current capability and future vision. 2089–2101 (2012).  
630 doi:10.1111/j.1365-2486.2012.02689.x
- 631 59. Richards, M. *et al.* High-resolution spatial modelling of greenhouse gas emissions from land-  
632 use change to energy crops in the United Kingdom. **44**, 627–644 (2017).
- 633 60. Song, J. *et al.* A meta-analysis of 1,119 manipulative experiments on terrestrial carbon-cycling  
634 responses to global change. *Nat. Ecol. Evol.* (2019). doi:10.1038/s41559-019-0958-3  
635