

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

Recherche d'une méthodologie pour la simulation des spectroscopies vibrationnelles exaltées par effet de surface de systèmes adsorbés

Matatu Mbengo, Christ

Publication date:
2022

[Link to publication](#)

Citation for published version (HARVARD):

Matatu Mbengo, C 2022, 'Recherche d'une méthodologie pour la simulation des spectroscopies vibrationnelles exaltées par effet de surface de systèmes adsorbés', 17^{eme} Rencontre des Chimistes Théoriciens Francophones, Bordeaux, France, 27/06/22 - 1/07/22.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.


- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Introduction

Vibrational spectroscopies are of great interest for the study of materials and molecules. Their simulation allows to have some insight about their structures. Several materials can be studied: nanoparticles, crystals or 2D materials. The spectroscopic signal of the sample can be enhanced by placing it near a nanostructured surface (Au or Ag) during the analysis. These are surface enhanced vibrational spectroscopies (SEVS). Their mechanisms are complex because of the coupling between the sample and the surface. A unified numerical approach would be useful for the simulation of SEVS. The SURFASCOPE (Surface Enhanced Spectroscopy by Second-Principles Calculations) project aims at designing and implementing a second-principles (2ndP) numerical approach to interpret and guide SEVS. 2ndP methods rely on access to 1stP data that can be used as parameters to study system responses, with a fixed topology. My methodological research is part of the current challenge of inhomogeneous systems treatment in a 2ndP approach. This treatment involves 1stP simulations on sufficiently small systems or those that benefit from a certain periodicity to deduce transposable parameters to larger structures.




Strategy

Study the adsorption of an alkyl chain on a surface using ONIOM QM/QM method as implemented in Gaussian. Previous results on the adsorption of n-octadecyl-trichlorosilane (OTS) adsorbed on a SiO₂ surface^[1] are used as a reference.

Optimization

Level of theory : ONIOM(wB97XD/6-311++G(2d,p):wB97XD/STO-3G



- ❖ The idea behind this strategy is to focus on a specific part of the structure. In this study, it is the interface between the surface and the molecule. The benefit is to diversify the data sources and focus on the most accurate way to describe each part of the large structure.
- ❖ The ONIOM models are designed to fulfill this objective. To deal with this 395 atoms structure, other options are needed. These options concern the choice of the SCF algorithm, convergence thresholds, and atom freezing.
- ❖ 2 model systems, with most of the surface in the low layer (blue) and the most relevant part of the surface and OTS in the high layer (red), are evaluated against the reference.
- ❖ For vibrational spectroscopy simulation, tight convergence criteria, which are the most difficult and challenging to achieve, are required.
- ❖ The default Gaussian optimization parameters and the short-chain ONIOM model consume fewer resources.



Vibrational frequencies

- ❖ Harmonic vibrational frequencies and normal modes are calculated analytically by setting atomic masses of 1000 amu to the atoms of the low layer and the H atoms saturating the dangling oxygen bonds.
- ❖ Frequencies are scaled (0.95) to account for missing anharmonicity effects as well as for the limitations of the XC functional.

RF				LCM		
vibration	FREQ (cm ⁻¹)	IR (Km/mol)	Raman (A ⁴ /AMU)	FREQ (cm ⁻¹)	IR (Km/mol)	Raman (A ⁴ /AMU)
(CH ₂) _{sym}	2884	271	30	2883	279	36
(CH ₃) _{sym}	2888	45	169	2886	43	224
(CH ₃) _{asym}	2955	87	31	2954	85	24
(CH ₃) _{asym}	2960	50	107	2959	49	105




Geometrical parameters

- ❖ The Z-axis (in dark blue) is defined as normal to the surface. The molecular z-axis (in light blue) is fitted using the least-squares approach as the best vector through the C atoms of the silane molecule.
 - ✓ θ : Tilt angle of the alkyl chain . The angle between the molecular z-axis and the laboratory Z-axis
- $$\theta = \cos^{-1}(Z \cdot z)$$
- ✓ α : Tilt angle of the terminal (CH₂ – CH₃) bond with respect to the Z-axis.
- $$\alpha = \cos^{-1}(Z \cdot CH_2 - CH_3)$$
- ✓ ξ : The rotation of the plane of the alkyl chain around the molecular z-axis.
- $$\xi = \text{atan2}(x \cdot Z, y \cdot Z)$$

- ❖ Analytical and experimental scales for those parameters from [1] :


$$\theta < 15^\circ$$

$$40^\circ < \alpha < 50^\circ$$




	Calculated from RF files	LCM	SCM
θ	11	15	16
ξ	-116	-171	-172
α	39	49	49

Conclusion



Perspectives



The calculations were performed on computers of the Consortium des Equipements de Calcul Intensif, including those of the Technological Platform of High-Performance Computing, supported by the FNRS-FRRC (Conventions No. 2.4.61707.F and 2.5020.11) and the University of Namur. Thanks to the ARC SURFASCOPE project for the founding and support on this work.

[1] C. G. Tetsassi Feugmo, V. Liégeois, Y. Caudano, F. Cecchet, and B. Champagne, "Probing alkylsilane molecular structure on amorphous silica surfaces by sum frequency generation vibrational spectroscopy: First-principles calculations," *The Journal of Chemical Physics*, vol. 150, no. 7, p. 074703, 2019, doi: [10.1063/1.5080007](https://doi.org/10.1063/1.5080007).