

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

SDPS-13 CELL LINE STUDY OF NUCLEOSOME-BASED BIOMARKERS IN THE DIAGNOSIS AND DETECTION OF RELAPSES IN GLIOBLASTOMA

Decarpentrie, Jonathan; Van den Ackerveken, Priscilla; Govaerts, Adrien; Lobbens, Alison; Beine, Edeline; Donis, Nathalie; Herzog, Marielle; Douxfils, Jonathan

Published in:
Neuro-Oncology Advances

DOI:
[10.1093/noajnl/vdad070.072](https://doi.org/10.1093/noajnl/vdad070.072)

Publication date:
2023

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (HARVARD):

Decarpentrie, J, Van den Ackerveken, P, Govaerts, A, Lobbens, A, Beine, E, Donis, N, Herzog, M & Douxfils, J 2023, 'SDPS-13 CELL LINE STUDY OF NUCLEOSOME-BASED BIOMARKERS IN THE DIAGNOSIS AND DETECTION OF RELAPSES IN GLIOBLASTOMA', *Neuro-Oncology Advances*, vol. 5, no. 3.
<https://doi.org/10.1093/noajnl/vdad070.072>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

; $p < 0.05$), phosphorylation of H3S10 (H3S10Ph) (control=2.31% ; GBM=5.57% ; $p < 0.05$), and a trend in H3K4Me2 elevation (control=0.43% ; GBM=1.07% ; $p = 0.066$) in GBM cell lines ($n = 4$). These results were confirmed by western blot. In addition, principal component analysis revealed a segregation between GBM and control cells, mainly driven by these three PTMs. Moreover, to demonstrate the capacity of the Nu.Q® assays to quantitatively monitor PTMs, GBM cell line SF-126 ($n = 3$) was treated with an EZH2 inhibitor (responsible of H3K27 methylation) (iEZH2). After 24 hours and 48 hours of exposure, the level of expression of H3K27Me3 – expressed as ratio of total nucleosomes was decreased by 32% (control=14.67%±0.58% ; iEZH2=10.00%±1.73% ; $p < 0.05$) and 41% (control=15.33%±1.53% ; iEZH2=9.00%±0.00% ; $p < 0.05$), respectively. Those results were also confirmed by western blot analysis. In conclusion, we identified three specific in vitro epigenetic-based marks of GBM and demonstrated that the Nu.Q® technology is a valuable tool to monitor the degree of PTMs expression.

ABSTRACT CITATION ID: vdad070.072

SDPS-13

CELL LINE STUDY OF NUCLEOSOME-BASED BIOMARKERS IN THE DIAGNOSIS AND DETECTION OF RELAPSES IN GLIOBLASTOMA

Jonathan Decarpentrie¹, Priscilla Van den Ackerveken², Adrien Govaerts², Alison Lobbens², Edeline Beine², Nathalie Donis³, Marielle Herzog², Jonathan Douxflis^{1,3}; ¹University of Namur, Department of Pharmacy, Namur Research Institute for Life Sciences, Namur, Belgium. ²Belgian Volition SRL, Isnes, Belgium. ³Qualiblood s.a., Namur, Belgium

Currently, glioblastoma (GBM) diagnosis and monitoring rely on neuroimaging and histopathological confirmation. However, overall survival has not improved in last decades due to therapeutic failure and to a lack of biomarkers for relapses' detection. Liquid biopsies (i.e. blood or cerebrospinal fluid) using nucleosomes-containing-histone post-translational modifications (PTMs) have the potential to become valuable biomarkers for diagnosis and monitoring GBM. Four glioblastoma cell lines (SF-126, U-87MG, U-118MG, and U-138MG) compared to a healthy microglia cell line (HMC3) and other solid cancer cell lines including pancreas (MIA PaCa-2), liver (HepG2), and cervix/uterus (HeLa) have been analyzed to identify their epigenetic profile. Nucleosomes were extracted and analyzed with the Nu.Q® Discover immunoassays platform (Belgian Volition) addressing 13 histone-PTMs. Quantitative results of PTMs expression were normalized on quantification of total nucleosomes (H3.1-nucleosomes). The immunoassay identified three candidate biomarkers compared to control cell lines ($n = 4$): citrullinated-histone H3 (H3R8Cit) (control=4.68% ; GBM=13.88%